

The International Lily Register and Checklist 2007

Second Supplement



© 2010 The Royal Horticultural Society
80 Vincent Square, London SW1P 2PE, United Kingdom
www.rhs.org.uk
RHS Registered Charity No. 222879/SC038262

International Lily Registrar 2006–2009:
Kate Donald

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781907057076

Printed and bound in the UK by Page Bros, Norwich

Cover: *Lilium* 'Willcroidii', drawing of Award of Merit plant by Winifred Walker, 1932.
Image courtesy of RHS Herbarium

The International Lily Register and Checklist 2007 Second Supplement

Introduction	page 1
Notes on the entries	page 1
Horticultural classification	page 2
Register and Checklist, Sep 2007–Aug 2009	page 4
Hybridizers, registrants and others	page 56

The lily epithets listed here were registered between 1 September 2007 and 31 August 2009. Details of lilies with unregistered names are published also.

Although registration is a voluntary procedure and does not confer any legal protection on the plant, as the International Cultivar Registration Authority for *Lilium*, The Royal Horticultural Society urges all hybridizers, raisers and introducers to register their lily names to reduce the potential confusion caused by new epithets which are the same as, or very similar to, existing names. Epithets which conform to the Articles (and, ideally, the Recommendations) of the latest edition of the *International Code of Nomenclature for Cultivated Plants* (ICNCP) are acceptable for registration. Registration, for which a certificate is issued on request, is free of charge, and should take place before a plant is released or described in a catalogue.

Registration forms are available as a download from RHS Online: www.rhs.org.uk; from Duncan Donald, International Lily Registrar, 16 Midtown of Inverasdale, Poolewe, Ross-shire IV22 2LW, UK; e-mail: lily@rhs.org.uk; tel: +44 (0)1445 781717; and from the Regional Representatives listed below.

The Royal Horticultural Society is indebted to the following Regional Representatives for their continued efforts to encourage the registration of lily names and for contributing amendments to *The International Lily Register and Checklist* and its *Supplements*:

• Australia:

Mr Neil Jordan, 27 Scarfe Street, Camdale,
Tasmania 7320
E-mail: neil.jordan@gotalk.net.au

• Czech Republic:

Ing. Břetislav Mičulka, 687 07 Velehrad 225
E-mail: miculka.b@tiscali.cz

• The Netherlands:

Mrs Saskia Bodegom, Koninklijke Algemeene
Vereeniging voor Bloembollencultuur,
Postbus 175, 2180 AD Hillegom
E-mail: bodegom@kavb.nl

• New Zealand:

Mrs Merle I. Lepper, P.O. Box 1394, Christchurch
E-mail: merle.lepper@xtra.co.nz

• North America:

Mrs Kathy Andersen, 2565 Charlestown Road,
Phoenixville, Pennsylvania 19460, USA
E-mail: ksa2006@verizon.net

Notes on the entries

The format of entries is similar to the layout adopted for *The International Lily Register and Checklist 2007*.

Parentage

The seed parentage is given first, preceding the larger multiplication sign.

Roles

The sequence and explanation of the abbreviations is as follows:

- S:** Selected by,
- H:** Hybridizer (reinstated in favour of the rather ambiguous term, Raiser),
- G:** Grown to first flowering by,
- N:** Named by,
- I:** Introduced by,
- REG:** Registrant.

Colour

Numbers and letters in parentheses refer to the *RHS Colour Chart* (1966 (reprinted 1986), 1995, 2001 and 2007). A colour name preceding a *Colour Chart* reference is taken from *A Contribution toward Standardization of Color Names in Horticulture: Application of the Universal Color Language to the Colors of the Royal Horticultural Society's Colour Chart* by RD Huse and KL Kelly, edited by DH Voss, The American Rhododendron Society, 1984.

Nomenclatural Standard

Sharp colour images, clearly showing the characteristics of the plant(s), and/or, if practicable, pressed specimens, are helpful. These, with the completed form, constitute a permanent, definitive record (nomenclatural standard) of the cultivar or Group. Those held in the RHS Herbarium at RHS Garden Wisley are under a designated WSY reference number.

Kate Donald (International Lily Registrar 2006–09) would like to thank colleagues for their support and advice – in particular Rupert Wilson, Dr Alan Leslie, Lawrie Springate, staff at the RHS Lindley Library (both at Wisley and Vincent Square), Mike Grant, Beatrice Somers (Modern Languages Dept, Gairloch High School) and, most especially, Duncan Donald.

• **Further information to augment existing lily records is always welcome.**

Horticultural classification

Division I Asiatic hybrids

Hybrids derived from the following species and interspecific hybrids: *amabile*, *bulbiferum*, *callosum*, *cernuum*, *concolor*, *dauricum*, *dauidii*, *L. × hollandicum*, *lancifolium* (syn. *tigrinum*), *lankongense*, *leichtlinii*, *L. × maculatum*, *pumilum*, *L. × scottiae*, *wardii* and *wilsonii*.

Flowers usually small to medium-sized. Flower colour often uniform, or with contrasting tepal tips and/or throat. Spots absent, or when present, well-defined and often rounded. Conspicuous brushmarks sometimes present. Tepal margins usually smooth or slightly ruffled. Flowers usually with little or no scent.

Examples: 'Ariadne' I(c/d), Citronella Group I(c/d), 'Mont Blanc' I(a/b-c).

Division II Martagon hybrids

Hybrids of martagon type derived from the following species and interspecific hybrids: *L. × dalhansonii*, *hansonii*, *martagon*, *medeoloides* and *tsingtauense*.

Flowers usually small, often numerous, mostly down-facing, typically with rather thick, recurved tepals and often of Turk's cap form. Spots usually present, numerous, often on at least three-quarters of each tepal. Flowers with little or an unpleasant scent. Buds often hairy. Bulbs often mauve or orange-yellow. Early flowering.

Examples: 'Cadense' II(b/d), 'Claude Shride' II(c/d), *L. × dalhansonii* 'Marhan' II(c/d).

Division III Euro-Caucasian hybrids

Hybrids derived from the following species and interspecific hybrids: *candidum*, *chalcedonicum*, *kesselringianum*, *monadelphum*, *pomponium*, *pyrenaicum* and *L. × testaceum*.

Flowers mostly small to medium-sized, often bell-shaped to Turk's cap-shaped, often down-facing. Flower colour often in rather pale muted shades. Spots absent to numerous. Tepal-margins smooth, often gently reflexed. Flowers often scented. Many are lime tolerant.

Examples: Caucasian Group III(c/d), 'Moonlight Madonna' III(-/a), 'Zeus' III(c/d).

Division IV American hybrids

Hybrids derived from the following American species and interspecific hybrids: *bolanderi*, *L. × burbankii*, *canadense*, *columbianum*, *grayi*, *bumboldtii*, *kellyanum*, *kelloggii*, *maritimum*, *michauxii*, *michiganense*, *occidentale*, *L. × pardaboldtii*, *pardalinum*, *parryi*, *parvum*, *philadelphicum*, *pitkinense*, *superbum*, *vollmeri*, *washingtonianum* and *wigginsii*.

Flowers small to medium-sized, mostly down-facing. Flower colour often strong yellow to orange or orange-red, often with contrasting centre and tepal tips. Spots often very conspicuous, over at least half of each tepal, often surrounded by a paler halo. Tepals rather narrow, margins smooth, usually gently to strongly reflexed. Flowers with little scent.

Examples: Bellingham Group IV(c/d), 'Lake Tulare' IV(c-c/d), 'Shuksan' IV(c/d).

Division V Longiflorum lilies

Hybrids or selections derived exclusively from *formosanum*, *longiflorum*, *philippinense* and *wallichianum*.

Flowers medium-sized to large, often few, trumpet-shaped. Flower colour typically uniform (white). Spots, papillae and brushmarks absent. Tepal margins smooth. Flowers usually scented.

Examples: Formolongi Group V(-/a), *longiflorum* 'Slocum's Ace' V(b/a).

Division VI Trumpet and Aurelian hybrids

Hybrids derived from the following species and interspecific hybrids: *L. × aurelianense*, *brownii*, *L. × centigale*, *henryi*, *L. × imperiale*, *L. × kewense*, *leucanthum*, *regale*, *rosthornii*, *sargentiae*, *sulphureum* and *L. × sulphurgale* (but excluding hybrids of *henryi* with all species listed in Division VII). Aurelian hybrids are derived from a combination of *henryi* and trumpet lilies.

Flowers medium-sized to large, in all flower forms. Flower colour white, cream, yellow to orange or pink, often with a contrasting star-shaped throat and/or strong bands of colour outside. Trumpets usually scented, without markings; other types often with spotting as small streaks in two bands at the base. Tepal-margins smooth or twisted with irregular ruffling, tips often reflexed.

Examples: Herald Angel Group VI(a/a), *L. × kewense* 'White Henryi' VI(b-c/c).

Division VII Oriental hybrids

Hybrids derived from the following species and interspecific hybrids: *auratum*, *japonicum*, *nobilissimum*, *L. × parkmanii*, *rubellum* and *speciosum* (but excluding all hybrids of these with *henryi*).

Flowers usually medium-sized to very large. Inner tepals often very broad and overlapping at the base to give a 'closed' centre. Flower colour mostly white to pink to purplish red, some golden yellow; ground colour often white, with a contrasting central ray. Spots absent to numerous, sometimes over half of each tepal. Papillae and nectaries often conspicuous. Flowers usually scented. Often late-flowering.

Examples: *L. × parkmanii* Imperial Silver Group VII(b/c), 'Woodriff's Memory' VII(a-b/b).

Division VIII Other hybrids

Hybrids not covered by any of the previous Divisions, I-VII. Includes all interdivisional hybrids, such as Asiatic/Trumpet (Asiapet or AT) hybrids, *longiflorum*/Asiatic (LA) hybrids, *longiflorum*/Oriental (LO) hybrids, *longiflorum*/Trumpet (Longipet or LT) hybrids, Oriental/Asiatic (OA) hybrids and Oriental/Trumpet (Orienpet or OT) hybrids. Hybrids of *henryi* with *auratum*, *japonicum*, *nobilissimum*, *L. × parkmanii*, *rubellum* and *speciosum* (excluded from Divisions VI and VII) are placed here.

Examples: 'Oberhof' VIII(b-c/-) AT, 'Salmon Classic' VIII(a-b/b) LA, 'Sky Treasure' VIII(b/a) LO, 'Sunny Crown' VIII(a/b) OA, 'Yelloween' VIII(a/b) OT.

Division IX Species and cultivars of species

Includes all species and their subspecies, varieties and forms, and cultivars selected therefrom (excluding those derived exclusively from *formosanum*, *longiflorum*, *philippinense* and *wallichianum*, which are placed in Div. V).

Examples: *auratum* var. *rubrovittatum* 'Crimson Beauty' IX(b/c); *canadense* var. *editorum* 'Chocolate Chips' IX(c/a); *lancifolium* 'Splendens' IX(c/d); *nepalense* IX(c/a).

Sub-divisions

If desired, it is possible to further classify lilies of all Divisions according to flower aspect and form. Flower aspect is given first and the two are separated by a forward slash (/).

Flower aspect:

- a** up-facing;
- b** out-facing;
- c** down-facing

Flower form:

- a** trumpet-shaped;
- b** bowl-shaped;
- c** flat (or with tepal tips recurved);
- d** tepals strongly recurved (with the Turk's cap form as the ultimate state).

Examples: 'Madison' I(a/b) indicates an Asiatic hybrid with up-facing, bowl-shaped flowers; 'Lightning Bug' II(b/d) a martagon hybrid with out-facing flowers with strongly recurved tepals; 'Snowdon' VII(a-b/-) an Oriental hybrid with flowers varying from up- to out-facing, but flower form unknown; *speciosum* 'Crimson Glory' IX(b-c/d) a selection from *speciosum* with out- to down-facing flowers with strongly recurved tepals.

Register and Checklist, Sep 2007–Aug 2009

‘Abbeville’s Pride’ I(a/b)

Parentage unknown; hybridized in the USA

N: Lily Company BV, 2008 **REG:** Lily Company BV, 2008

Inside of tepals strong orange (25A) to 26A), with vivid orange-yellow (23A) patch above throat; throat moderate red (185B). Outside of outer tepals dark pink (182C), flushed dark purplish pink (186C) towards tips; margins strong orange (25A); a small spot of moderate purplish red (186A) and strong yellow-green (144A) at tips; base strong yellow-green (144C). Outside of inner tepals strong orange-yellow (22A) to vivid orange-yellow (23A), shading to light orange-yellow (22B) at tips; margins strong orange (25A); base flushed moderate red (185B); midribs flushed moderate purplish red (186B). Spots dark red (183A); papillae light orange-yellow (22D); nectaries strong yellow-green (144A); pollen dark red (183A); stigma dark red (183A) to deep red (185A). Fls 165mm wide; tepals 86 × 52mm, margins smooth, tips recurved. Lvs 105 × 16mm, dark green. Stems 0.75m, green with darker markings, with up to 6 fls. Named for Abbeville Opera House, Abbeville, South Carolina, which opened in 1908.

‘Adios Noninja’ I(a-b/b-c)

N: Fa Q.J. Vink en Zn BV, c.2006

Tepals milk-white; midribs stained pale yellowish green. Spots forming a solid patch of dark burgundy-red in lower two-fifths of each tepal, becoming progressively less dense at margins towards apex and towards central portion of upper half of each tepal, producing a picotee or stippling effect; also present on the outside of each tepal; pollen orange-brown; stigma dull brownish purple. Tepals of medium width, margins smooth, tapering quite abruptly to slightly recurved tips. Stems 0.9m. Mid-season. Fl colouring similar to ‘Burgundy Splash’, in which the burgundy-red coloration is less solid, and ‘Black Spider’, in which there is less stippling beyond the solid patch of burgundy-red.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Alicja Michalska’ VI(b/b-c)

‘Eva Steinbrück’ × unknown

H: R. Widerski, pre-2008 N: R. Widerski, 2008

I: R. Widerski, 2008 **REG:** R. Widerski, 2008

Tepals cream-white with rays of orange in basal two-thirds; midribs green and purple. Papillae small; nectaries and pollen orange; stigma brown. Fls 140mm wide; inner tepals 120 × 40mm, margins slightly ruffled, tips recurved; with secondary buds. Lvs 120 × 25mm, glossy, mid-green. Stems 1.2m, spotted purple, infl. of medium length. Early July.

‘Alinka’ I(a-b/b-c)

Parentage unknown

H: Mak Breeding BV, pre-2007 N: Mak Breeding BV,

2007 **REG:** Mak Breeding Rights BV, 2009

Inside of tepals yellowish white (NN155A); base of midveins white flushed purple. Outside of inner tepals predominantly yellowish white (NN155A). Spots few; papillae present; nectaries green; pollen dark brown; stigma purple. Fls 130–140mm wide; tepals short and of narrow to medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length, and of narrow to medium width, dark green. Stems 1.05m, green with darker spots and stripes, with up to 8 fls. Early to mid-season.

‘Altavilla’ VIII(b-a/b) OT

Seedling PG 96-12 × seedling RH 99-07

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals brilliant greenish yellow (5B) in basal half, including throat; pale greenish yellow (1D) in upper half. Inside of outer tepals brilliant greenish yellow (6C), shading to brilliant yellow (13C) on one side of each tepal. Outside of outer tepals pale greenish yellow (2D); one side of each tepal spotted strong purplish red (58C); margins light greenish yellow (3C); midribs strong yellow-green (144A at base, 143A at top and 144C between). Outside of inner tepals pale greenish yellow (2D), shading to pale yellow-green (4D) beside midribs; midribs strong yellow-green (144B) at base. Spots absent; papillae brilliant greenish yellow (5B); nectaries strong yellow-green (144A); pollen greyish brown (166A); stigma dark purplish red (N79A). Fls 185mm wide, slightly scented; tepals 120 × 55mm, margins ruffled, tips recurved. Lvs 220 × 27mm. Stems 1.4m, green, with up to 6 fls.

‘Amateras’ VIII(b-a/b) LA

Seedling BJ 95-19 × seedling OR 00-25

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals vivid yellowish pink (28A); margins of outer tepals vivid reddish orange (N30C); throat moderate yellowish pink (31D). Outside of outer tepals brilliant orange (29A); margins vivid reddish orange (30A); midribs strong yellow-green (144C at base and 144B at top). Outside of inner tepals light orange (28C); midribs strong yellow-green (144C) at base, vivid reddish orange (30A) at top. Spots and papillae absent; nectaries dark green (136A); pollen strong reddish orange (169A); stigma deep yellowish pink (46D). Fls 195mm wide; tepals 115 × 52mm, margins smooth, tips slightly recurved. Lvs 145 × 25mm. Stems 1m, green, with up to 8 fls.

The name of a Japanese sun goddess.

‘American Sentinel’ VIII(b-c/b) OT

H: Van der Salm, pre-2008

Inside of tepals vibrant, strong lemon-yellow; a large, rounded, auburn-red zone dominating basal half, almost extending to margins at base, leaving a narrow margin of strong lemon-yellow, and feathering higher along midveins. Spots and papillae absent; nectary furrows apple-green; filaments and style yellowish green; pollen dark reddish brown; stigma pale green. Tepal margins slightly wavy, tips recurved. Stems 1.2 to 1.5m. Late.

Published in *The Lily Nook cat.*, 2009 as ‘American Sentinel’

AMFORA VIII(a/b) OT

Trade designation of ‘Zanotfora’

‘Anacapri’ VII(b-a/b)

Seedling RM 96-22 × seedling 00-30

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Tepals, including throat, greenish white (155C); midribs pale yellow-green (4D) towards top. Spots absent; papillae greenish white (155C); nectaries strong yellow (N144B); pollen greyish red (178A); stigma greyish yellow-green (195B). Fls 200mm wide; tepals 115 × 57mm, margins slightly ruffled, tips slightly recurved. Lvs 210 × 40mm, dark green. Stems 1.2m, pale green with some darker markings at base, with up to 6 fls.

The name of a community on the island of Capri.

‘Andrews’ VII(a/b)

Seedling RM 98-18 × seedling RM 96-52

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals dark purplish pink (186C), shading to deep purplish pink (N66C) towards margins of outer tepals; throat yellowish white (155D). Outside of outer tepals dark purplish pink (186C), shading to deep purplish pink (N66D) towards margins and yellowish white (155D) beside midribs at base; midribs strong yellow-green (144C) at base shading to light olive (152A) at top. Outside of inner tepals light purple (75B), shading to deep purplish pink (N66D) towards margins and yellowish white (155D) beside midribs at base; midribs light yellow-green (144D) at base. Spots and papillae absent; nectaries strong yellow-green (144B) edged light greenish yellow (3D); pollen dark orange (N163A); stigma very pale green (192B). Fls 240mm wide, scented; tepals 160 × 80mm, margins ruffled, tips recurved. Lvs 173 × 58mm. Stems 0.8–1.1m, green with darker markings, with 1 or 2 fls.

‘Angel Blush’ VI(b-c/b)

Seedling bred in New Zealand by McLaren

× ‘Stowport Big Pink’

H: L.C. Bligh, 2003 **G:** L.C. Bligh, 2005

N: L.C. Bligh, 2008 **REG:** L.C. Bligh, 2008

Inside of tepals white, shading to very light purple (75C) towards margins and brilliant yellow-green (154C) towards base; throat brilliant yellow-green (154C). Outside of tepals light purple (78C), shading to white at margins. Spots absent; nectaries green; pollen golden brown; stigma chocolate brown. Fls 150mm wide, slightly scented; tepals 125 × 55mm, margins smooth, tips slightly recurved. Lvs 150 × 11mm, scattered. Stems 1.5m, green, with up to 16 fls. Mid-January (Tasmania).

‘Anna Magdalena’ I(c/c)

‘Saartiger’ × ‘Saarturm’

H: R. Widerski, pre-2008 **N:** R. Widerski, 2008

I: R. Widerski, 2008 **REG:** R. Widerski, 2008

Inside of tepals dark red-purple. Outside of tepals red; midribs pale red-purple. Spots small, red-purple, fairly numerous, over half of each tepal; pollen orange; stigma red. Fls 90mm wide; inner tepals 70 × 35mm, tips strongly recurved; pedicels 110mm, stiff, spotted purple; with secondary buds. Lvs 110 × 10mm. Stems 0.8m, slightly felted, mid-green, infl. of medium length. Early July.

‘Antikvita’ I(a/b-c)

(‘Medulena’ × ((‘Connecticut Lemonglow’ × ‘Fuga’) × ‘Quinella’)) × unknown

H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007

I: B. Mičulka, 2007 **REG:** B. Mičulka, 2008

Inside of tepals bronze orange-brown; throat darker and striped. Outside of tepals yellowish in centre; midribs yellow-green. Brushmarks small, conspicuous, mahogany brown; spots few; nectaries frosted white; filaments brown-red; pollen chocolate brown; style pale brown; stigma brown. Fls 135mm wide; tepals 85 × 47mm, margins slightly ruffled, tips slightly recurved; pedicels 45mm, stiff, slightly spotted. Lvs dull, pale green. Stems 0.75m, with a few brown spots. Late June.

‘Antoni Widerski’ VI(c/d)

‘Louis XIV’ × unknown

H: R. Widerski, pre-2008 **N:** R. Widerski, 2008

I: R. Widerski, 2008 **REG:** R. Widerski, 2008

Inside of tepals yellow. Outside of tepals yellow-green; midribs yellow. Nectaries green; pollen orange; stigma pale green. Fls 160mm wide; inner tepals 100 × 40mm, tips recurved; with secondary buds. Lvs 110 × 35mm, glossy. Stems 1.2m, green, infl. of medium length. Mid-July.

'Apogee' VII(a/b)

Seedling 93-65 × seedling RW 98-12

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals deep purplish pink (N57C), shading to N57D towards margins; tips tinged yellowish white (155D); throat 155D. Outside of outer tepals moderate purplish red (186B); tips and base tinged yellowish white (155D); midribs strong yellow-green at base (144C) and top (143C); yellowish white (155D) between. Outside of inner tepals dark purplish pink (186C); tips and base tinged yellowish white (155D); midribs tinged strong yellow-green (143C) at base and top. Spots strong purplish red (60B); papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma light purplish grey (N187D). Fls 220mm wide, scented; tepals 130 × 65mm, margins slightly ruffled, tips recurved. Lvs 160 × 50mm. Stems 1.2m, green, with up to 5 fls.

'Appia' VIII(a/b) LA

Seedling BG 95-71 × seedling AJ 98-04

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005 **REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals strong orange (25B); margins and midveins 25A; throat pale yellow (20C). Outside of outer tepals light orange-yellow (23C), shading to strong orange (25A) towards margins and greyish red (178A) beside midribs at base; midribs brilliant yellow-green (142B), shading to strong yellow-green (143B) at top. Outside of inner tepals strong orange-yellow (24B), shading to strong orange (25B) towards margins; midribs strong yellow-green (143C) at base. Spots dark red (59A); papillae brilliant yellow (20A); nectaries moderate olive-green (137A); pollen strong brown (172A); stigma strong orange (30D). Buds hairy. Fls 225mm wide; tepals 130 × 63mm, margins smooth, tips recurved. Lvs 163 × 23mm. Stems 1.3m, green with darker markings, with up to 7 fls.

'Arcato' VII(a/b)

Seedling RH 99-15 × seedling RW 96-05

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals, including throat, white (NN155C); basal two-thirds of midveins pale greenish yellow (2D). Outside of tepals white (NN155D); midribs strong yellow-green (143B) at base, those on outer tepals shading to pale greenish yellow (2D) and strong yellow-green (143B) at top. Spots and papillae pale greenish yellow (2D); nectaries strong yellow-green (141D); pollen strong orange (N25A); stigma light greenish grey (188C). Fls 200mm wide, slightly scented; tepals 135 × 75mm, margins slightly ruffled, tips slightly recurved. Lvs 235 × 65mm, dark green. Stems 1.4m, green, with up to 6 fls.

'Ariane' VII(a-b/b)

Parentage unknown

N: M.A. van den Berg en Zn, V.o.f., pre-2009

Tepals greenish white (155C). Spots absent; papillae present; nectaries green; pollen red-brown; stigma speckled purple-red. Tepals of medium length to long and of medium width to broad, margins ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad to very broad. Stems 0.95m, green, with up to 6 fls. Mid-season.

An application for registration was turned down in 2009, as the cultivar epithet is too similar to, and likely to cause confusion with, 'Ariadne', which was in cultivation at the time (ICNCP, 2004: Art. 19.25). However, as this lily was granted Dutch Plant Breeders' Rights under the name 'Ariane' in 2008, it remains the accepted epithet (*ibid.*: Art. 24.4).

'Arlecchino' VII(b/b-c)

Seedling RM 96-65 × seedling RM 98-01

H: Vletter en Den Haan Beheer BV, pre-2008**N:** Vletter en Den Haan Beheer BV, 2008**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals vivid purplish red (61C), with brilliant greenish yellow (3A) spot above nectaries and light greenish yellow (3C) spots towards margins; midveins vivid reddish orange (33B); throat pale yellow-green (4D). Outside of outer tepals deep purplish pink (N57C); midribs strong yellow-green at base (144C) and top (144A); strong yellow (153D) between. Outside of inner tepals deep purplish pink (N57C), N57D flushed vivid reddish orange (33B) in basal third, shading to pale yellow-green (4D) beside midribs at base; midribs strong yellow-green (144C), stained deep purplish pink (N57C) towards top. Spots strong purplish red (61B); papillae brilliant greenish yellow (3A); nectaries strong yellow-green (144C); pollen moderate reddish orange (171A); stigma light greenish grey (191D). Fls 250mm wide, scented; tepals 140 × 75mm, margins slightly ruffled, tips recurved. Lvs 253 × 53mm. Stems 1.25m, green with darker markings, with up to 5 fls.

'Arlington' VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Tepals, including throat, yellowish white (155D); midveins light yellow-green (150D); midribs of outer tepals strong yellow-green at base (144C) and top (144B); light yellow-green (150D) between; midribs of inner tepals light yellow-green (144D) at base shading to 150D). Spots absent but some inner tepals with small purple-red stripes; papillae yellowish white (155D); nectaries light yellow-green (144D); pollen moderate reddish orange (171A); stigma light grey (191C). Fls 190mm wide, scented; tepals 125 × 75mm, margins slightly ruffled, tips slightly recurved. Lvs 195 × 50mm. Stems 1.3m, green with darker markings at base, with up to 6 fls.

'Arosa Jewel' I(a/b-c)

N: De Jong Lelies Holland BV, c.2008

Tepals sparkling, deep magenta-pink. Spots and papillae absent; pollen orange-brown; stigma pinkish.

Tepals rather narrow, margins slightly ruffled, tips

slightly recurved. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009External images: *The Lily Nook cat.*, 2009**'Arsenal' I(a/c)**

Parentage unknown

H: Mak Breeding BV, pre-2007 **N:** Mak Breeding BV, 2007 **REG:** Mak Breeding Rights BV, 2009

Inside of tepals strong red (53B/C), more blue with age, with a large, light orange-yellow (16C) to pale orange-yellow (16D) patch towards base, and strong red (53C) beneath; midveins orange-yellow at base. Outside of inner tepals strong red (53B/C). Spots few to moderately numerous, dark; papillae present; nectaries green shading to purple-red; pollen red-brown; stigma orange. Fls 150–200mm wide; tepals short and of medium width, margins slightly ruffled, tips (very) slightly recurved. Lvs of medium length, and of narrow to medium width, dark green. Stems 1.15m, green with darker spots and stripes, with up to 8 fls. Early to mid-season.

'Arzachena' VII(a/b)

Seedling RW 99-17 × seedling RW 96-05

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Tepals, throat and papillae, greenish white (155C); midrib with a stripe of pale greenish yellow (1D) towards top. Spots absent; nectaries strong yellow (N144B); pollen dark reddish orange (175B); stigma dark purple (79B). Fls 220mm wide; tepals 130 × 70mm, margins ruffled, tips recurved. Lvs 160 × 40mm, dark green. Stems 1.25m, pale green, with up to 5 fls. A town in Sardinia, Italy.

'Asinara' VIII(b-a/b) OT

Seedling PG 96-70 × seedling RH 99-24

H: Vletter en Den Haan Beheer BV, pre-2006**N:** Vletter en Den Haan Beheer BV, 2006**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals brilliant yellow (12B), shading to light greenish yellow (6D) towards tips and in throat. Inside of outer tepals vivid yellow (12A), shading to brilliant yellow (13C) towards tips. Outside of outer tepals pale yellow (8D) tinged strong purplish red (58B), shading to brilliant greenish yellow (6C) towards margins; midribs strong yellow-green (143C) at base, flushed strong purplish red (58B); 144C at top). Outside of inner tepals light greenish yellow (5D), shading to pale yellow (11D) beside midribs; base of midribs light yellow-green (144D) flushed strong purplish red (58B). Spots brilliant yellow (12B); papillae absent; nectaries strong yellow-green (143C); pollen moderate reddish brown (175A); stigma dark greyish purple (N92A). Fls 195mm wide, scented; tepals 132 × 70mm, margins slightly ruffled, tips recurved. Lvs 180 × 47mm, dark green. Stems 1.3m, green, with up to 6 fls.

AUBERGE VIII(b/b) LO

Trade designation of 'Zanloberge'

'Augusta' VII(b-a/b)

Seedling RW 96-05 × seedling 01-330

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Tepals and throat, yellowish white (155D); midveins light yellow-green from above nectaries to half-way up (150D on inner tepals, 144D on outer); midribs strong yellow-green (144C) at base shading to light yellow-green (145D on outer tepals, 145C on inner); midribs of outer tepals strong yellow-green (144A) at top. Spots and papillae yellowish white (155D); nectaries strong yellow-green (144B); pollen strong reddish orange (169A); stigma yellowish grey (c.198C). Fls 225mm wide, slightly scented; tepals 130 × 75mm, margins ruffled, tips slightly recurved. Lvs 175 × 42mm. Stems 1.2m, green, with up to 7 fls.

'Barletta' VII(a/b)

Seedling 93-14 × seedling RM 98-25

H: Vletter en Den Haan Beheer BV, pre-2006**N:** Vletter en Den Haan Beheer BV, 2006**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals light purplish pink (63D), with narrow, yellowish white (155D) margins at tips; midveins pale purplish pink (65C); throat yellowish white (155D). Inside of outer tepals strong purplish pink (63C), with narrow, yellowish white (155D) margins at tips; midveins yellowish white (155D) in basal half. Outside of tepals moderate purplish pink (186D), shading to yellowish white (155D) beside base of midribs, at margins (these are narrower in inner tepals) and at tips; midribs of outer tepals strong yellow-green (144C) at base and top, yellowish white (155D) between; midribs of inner tepals strong yellow-green (144C) at base shading to pale greenish yellow (2D). Spots and papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma dark purplish red (N79C). Fls 220mm wide, scented; tepals 140 × 80mm, margins ruffled, tips slightly recurved. Lvs 195 × 55mm. Stems 1m, green with darker markings at base and leaf nodes, with up to 3 fls.

'Baruta' VIII(a-b/b-c) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** World Breeding BV, 2003 **REG:** World Breeding BV, 2008

Inside of inner tepals brilliant yellow (11A), inside of outer tepals light yellow (11B), all shading to pale yellow-green (4D) towards margins and tips. Outside of inner tepals light yellow (11B) to pale yellow (11C). Spots absent; papillae present; nectaries green; pollen red-brown; stigma purple. Fls 240mm wide; tepals of medium length to long and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and of medium width to broad. Stems 1.4m, green with faint darker spots and stripes, with up to 3 fls. Late to very late.

‘Baywatch’ VIII(a-b/b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV,

2003 **REG:** World Breeding BV, 2008

Inside of tepals flushed deep purplish pink (N57C/D) on a lighter ground, paler towards tips and shading to pale yellow (8D) at base; becoming paler with age.

Outside of inner tepals light yellowish pink (36C) to pale yellowish pink (36D). Spots few; papillae present; nectaries green; pollen red-brown; stigma green with purple-red spots. Fls 220mm wide; tepals of medium length to long and of medium width, margins slightly ruffled, tips slightly recurved. Lvs of medium length to long, broad. Stems 1.1–1.35m, green with some darker markings, with up to 6 fls. Late to very late.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Belletti’ I(a/c)

Parentage unknown

H: Mak Breeding BV, pre-2006 **N:** Mak Breeding BV, 2006 **REG:** Mak Breeding Rights BV, 2008

Tepals greenish white (155C); inside margins dark at base. Spots and papillae present; nectaries green; pollen orange-brown; stigma grey-green purple.

Tepals short and of narrow to medium width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length and of narrow to medium width. Stems 1.3m, green with darker spots and stripes, with up to 6 fls. Mid-season.

‘Beppo’

Unregistered pollen parent of ‘Schweriner Morgen’. No further details known.

‘Beyoncé’ VIII(a/b) LA

Asiatic seedling × seedling LA 94027-2T

H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2008

Inside of tepals light purplish pink (65B), with a yellowish white (155D) spot above nectaries; midveins of outer tepals with a broad, light purplish pink (65B) band; throat yellowish white (155D). Outside of tepals dark purplish pink (186C); midribs of outer tepals strong yellow-green (143C) at base, 143B at top; midribs of inner tepals moderate yellow-green (139D) at base shading to moderate purplish red (186B). Spots and papillae absent; nectaries moderate olive-green (137A); pollen moderate reddish orange (N172A); stigma light yellow (18A). Fls 165mm wide; tepals 125 × 65mm, margins smooth, tips straight. Lvs 150 × 35mm. Stems 1.2m, dark in basal half, green with darker markings in upper half, with up to 6 fls.

‘Birmingham’ VIII(b-a/b) OT

Seedling PG 96-43 × seedling RH 99-07

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals vivid yellow (9B), shading to brilliant yellow (9C) towards margins; midveins vivid yellow (9A); throat light greenish yellow (8B). Inside

of outer tepals vivid yellow (9A) to brilliant yellow (9C); midveins flushed strong red (39A) at top. Outside of outer tepals pale yellow-green (4D), shading to brilliant yellow (8A) towards margins and flushed strong red (39A) beside midribs. Outside of inner tepals light greenish yellow (8B), shading to pale yellow (8D) beside midribs; midribs light yellow-green (144D) at base shading to light greenish yellow (8B). Spots absent; papillae vivid yellow (9B to 14A); nectaries strong yellow-green (143C); pollen greyish red (178A); stigma moderate reddish purple (N79D). Fls 210mm wide, scented; tepals 140 × 70mm, margins ruffled, tips slightly recurved. Lvs 250 × 40mm. Stems 1.3m, green, with up to 5 fls.

‘Black Spider’ I(a/b)

N: “The Originals” U.A., Coöp. Kwekersvereniging, c.2005

Tepals ivory-white, more yellowish on opening. Spots forming a solid patch of dark burgundy-red in lower two-fifths of each tepal, but less dense towards edges of each patch, producing a picotee or stippling effect; pollen orange-brown. Tepals rather narrow, margins slightly ruffled, tips slightly recurved. Stems 0.9m. Mid-season. Fl colouring similar to ‘Burgundy Splash’, in which the burgundy-red coloration is less solid but more extensive, and ‘Adios Noninja’, in which there is more stippling beyond the solid patch of burgundy-red. Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Boldini’ VIII(a/b) LA

Seedling BG 96-49 × seedling AG 97-05

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals brilliant greenish yellow (6A), shading to brilliant yellow (7A) towards margins; throat light greenish yellow (1C). Outside of outer tepals pale greenish yellow (2D), shading to brilliant yellow (7A) towards margins; midribs strong yellow-green (143C) at base shading to brilliant yellow-green (150B).

Outside of inner tepals light greenish yellow (5D), shading to brilliant yellow (7A) towards margins; midribs light yellow-green (144D). Spots deep red (60A); papillae brilliant greenish yellow (6A); nectaries moderate olive-green (137A); pollen moderate reddish brown (175A); stigma brilliant yellow-green (154B). Fls 210mm wide, slightly scented; tepals 120 × 60mm, margins smooth, tips recurved. Lvs 170 × 22mm. Stems 1m, green with darker markings, with up to 5 fls. An Italian portrait painter, 1842–1931.

‘Bonbini’ VIII(a-b/b) OT

H: World Breeding BV, pre-2008

Inside of tepals milk-white; in centre of basal two-fifths a conspicuous, pale creamy yellow diamond-shaped flare, with a soft purplish rose-pink, slightly feathered margin. Nectary furrows at base of flare apple-green. Spots few, small, round, mostly scattered near margin of flare on inner tepals, darker carmine-pink; papillae absent; filaments and style pale bright green; pollen warm brown; stigma blackish brown. Tepals quite broad near base but

rather slender in upper half, margins slightly wavy, tips slightly recurved. Stems 1.2m. Late.
Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

‘Bonviván’ VI(b-c/a-b)

((‘Damsón’ × Golden Splendor Group) × African Queen Group) × ((*regale* ‘Royal Gold’ × unknown) × Golden Splendor Group)

H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007

I: B. Mičulka, 2007 **REG:** B. Mičulka, 2007

Syn. ‘Slovačka’

Inside of tepals lemon-yellow, paler towards tips; margins marbled pink. Outside of tepals marbled ‘old rose’ pink; midribs mahogany-carmine. Spots absent; nectaries deep green; pollen apricot-brown; stigma yellow-green. Fls 115mm wide; tepals 110 × 68mm, tips recurved; pedicels 60mm, spotted dark brown. Lvs 140 × 13mm, dull, dark green. Stems 1.1m, with numerous black-brown spots. Early July.

‘Bracelet’ VIII(a-b/b) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006

N: Testcentrum voor Siergewassen BV, 2006

REG: Testcentrum voor Siergewassen BV, 2008

Inside of tepals more red than moderate purplish red (186B), paling to dark purplish pink (186C) towards tips; midveins white at base, shading to pink. Outside of tepals moderate purplish red (186B) to dark purplish pink (186C). Spots small or very small; papillae present; nectaries green shading to purple-red; pollen orange; stigma light orange with purple spots. Tepals short and of narrow to medium width, margins (very) slightly ruffled, tips (very) slightly recurved. Lvs of medium length and of narrow to medium width. Stems 1.3m, green with darker spots and stripes, with up to 7 fls. Mid-season to late.

‘Brilliant Star Delight’ VII(a/c)

‘Mero Star’ × ‘Expression’

H: P.D. Hoogland, pre-2008 **N:** Maveridge International BV, pre-2008 **REG:** Maveridge International BV, 2008

Inside of tepals strong purplish red (61B), shading to 64C towards margins; margins, tips and throat greenish white (155C). Outside of tepals strong purplish red (64B/C) on a paler ground; margins greenish white (155C); base flushed strong yellow-green (144C). Spots deep red (60A); papillae greenish white (155C) and strong purplish red (64C); nectaries strong yellow-green (144B); pollen dark reddish orange (175B); stigma greyish yellow-green (196A). Fls 240mm wide; tepals 125 × 69mm, margins smooth, tips recurved. Lvs 150 × 50mm, dark green. Stems 1.1m, pale green with some darker markings, with up to 5 fls.

‘Brilliant Star Magic’ VII(a/c)

‘Mero Star’ × ‘Tiber’

H: P.D. Hoogland, pre-2008 **N:** Maveridge International BV, pre-2008 **REG:** Maveridge International BV, 2008

Inside of tepals strong purplish red (67A), shading to

vivid purplish red (67B) towards margins; margins deep purplish pink (N66C); tips pinkish white (N155B); throat greenish white (155C). Outside of tepals strong purplish red (63B) on a paler ground; midribs greenish white (155C) at base shading to strong purplish red (63B) towards top; top pale yellow-green (4D). Spots deep red (60A); papillae deep red (60A) to strong purplish red (60B); nectaries light yellow-green (145B); pollen dark reddish orange (175B); stigma dark purple (79A). Fls 215mm wide; tepals 116 × 62mm, margins ruffled, tips recurved. Lvs 160 × 50mm, dark green. Stems 0.9m, pale green, with up to 5 fls.

‘Bronze Medallion’ II(c/d)

H: E. Robinson, pre-2008

Inside of tepals bronze; muted old-gold near spots in basal half of each tepal. Spots chocolate-brown, quite large, round to elliptic, evenly and densely distributed over basal half of each tepal; also along margins on outside of tepals. Papillae absent; filaments yellowish, pale; pollen dark orange; stigma pale orange. Buds slightly bronzed rose-pink. Pedicels long, arching, dark brownish green. Stems 1.5–1.8m, dark. Early. Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Bulgari’ VIII(a/b) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006

N: Testcentrum voor Siergewassen BV, 2006

REG: Testcentrum voor Siergewassen BV, 2007

Tepals more red than dark red (187C). Spots few; papillae inconspicuous; nectaries green; pollen brown; stigma yellow-green. Tepals of short to medium length and of medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length to long, and of medium width. Stems very tall, green with darker spots and stripes, with average no. to many fls. Late to very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

‘Burgundy Splash’ I(a-b/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2006 **N:** ‘The Originals’ U.A., Coöp. Kwekersvereniging, 2006 **REG:** Mak Breeding Rights BV, 2008

Inside of tepals pale yellow-green (155A) to yellowish white (155D); midveins flushed orange in young fls. Outside of tepals yellowish white (155D). Spots dark red (59A) or darker, coalescing in basal half in inside of each tepal and almost obliterating underlying colour; becoming progressively less dense along margins towards apex and towards central portion of upper half of each tepal; also present on the outside of each tepal; papillae present; nectaries green; pollen pale brown; stigma purple. Fls 140mm wide; tepals short or very short and of narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of short to medium length and of narrow to medium width. Stems 0.8–1.15m, green with darker spots and stripes, with few to average no. of fls. Early to mid-season. External web links: *The Lily Nook cat.*, 2009

'Bussana' VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals vivid purplish red (67B), shading to strong purplish red (64C) towards margins; throat pale yellow-green (155A) and brilliant greenish yellow (2B). Outside of tepals strong purplish red (64C); pale yellow-green (155A) at base; midribs light yellow-green (154D); strong greenish yellow (151A) at top. Spots dark red (59A); papillae moderate purplish red (59C); nectaries strong yellow-green (144A); pollen strong brown (172A); stigma very pale green (192B). Fls 210mm wide; tepals 137 × 84mm, margins slightly ruffled, tips recurved. Lvs 200 × 46mm. Stems 1.15–1.3m, pale green, with up to 4 fls. Old and New Bussana are towns in Liguria, Italy.

'Cadore' VII(a/b)

Seedling RM 96-24 × seedling RW 97-04

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals moderate purplish pink (186D); inside of outer tepals dark purplish pink (186C); margins, beside nectaries and throat, yellowish white (155D). Outside of outer tepals very pale purple (69D); margins and base yellowish white (155D); midribs strong yellow-green (144B) at base, flushed light purplish pink (63D) above. Outside of inner tepals yellowish white (155D) with light purplish pink (63D) stripes; midribs strong yellow-green (144C) at base. Spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen moderate reddish orange (171A); stigma dark purple (79A). Fls 250mm wide, scented; tepals 145 × 85mm, margins slightly ruffled, tips slightly recurved. Lvs 205 × 50mm. Stems 1.2m, dark, with up to 3 fls. A mountain community in the region of Veneto, Italy.

'Caldeira' VII(a/c-b)

Seedling RW 94-06 × seedling RW 98-06

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Tepals and throat, yellowish white (155D); midveins pale yellow-green (155A); midribs of outer tepals strong yellow-green at base (144A) and tinged 144B at top; light yellow-green (150D) between; midribs of inner tepals light yellow-green (144D) at base. Spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144C) to light yellow-green (144D); pollen dark orange (N163A); stigma greyish purplish red (N77D). Fls 230mm wide, slightly scented; tepals 125 × 72mm, margins slightly ruffled, tips slightly recurved. Lvs 225 × 55mm. Stems 1.1m, green, with up to 6 fls.

'Camaioire' VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals strong purplish pink (73B), yellowish white (155D) above nectaries, shading to deep purplish pink (64D) towards margins of outer tepals; midveins of inner tepals strong purplish red (63B) at top; midveins of outer tepals deep purplish pink (64D) at top; throat yellowish white (155D). Outside of tepals moderate purplish pink (186D), margins dark purplish pink (186C), yellowish white (155D) beside base of midribs; midribs strong yellow-green (144A) at base shading to strong yellow (153D). Spots strong purplish red (58B); papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma pale yellow-green (196C). Fls 240mm wide, scented; tepals 140 × 85mm, margins ruffled, tips recurved. Lvs 180 × 40mm. Stems 1.17m, green, with darker markings near leaf nodes, with up to 3 fls.

'Camengo' VII(a/c)

Parentage unknown

H: Mak 't Zand BV, pre-2007 **N:** Mak Breeding BV, 2007 **REG:** Mak 't Zand BV, 2009

Inside of tepals deep purplish pink (N66D), base and midveins white (NN155C). Outside of inner tepals predominantly deep purplish pink (N66D). Spots absent; papillae white and bluish pink; nectaries green; pollen red-brown; stigma grey-green. Fls 200–250mm wide; tepals of medium length to long and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length and broad, dark green. Stems of average height to tall, green with darker spots and stripes, with up to 8 fls. Mid-season.

'Canaletto' VII(b-a/c)

Seedling RW 96-05 × seedling RM 96-03

H: Vletter en Den Haan Beheer BV, pre-2006**N:** Vletter en Den Haan Beheer BV, 2006**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals strong purplish red (60D) to deep purplish pink (N66C), shading to dark purplish pink (186C) towards margins; throat white (NN155D). Inside of outer tepals deep purplish pink (N66C); margins and tops of midveins white (NN155C). Outside of outer tepals dark purplish pink (186C), shading to deep purplish pink (N57C) towards margins; midribs strong yellow-green (143C) at base shading to moderate yellow-green (146B). Outside of inner tepals dark purplish pink (186C), shading to moderate purplish pink (70D) towards margins; midribs light yellow-green (145B) at base. Spots few, strong purplish red (60B); papillae strong purplish red (60D) to deep purplish pink (N66C); nectaries strong yellow-green (144B); pollen strong orange (N25B); stigma pale yellow-green (192A). Fls 225mm wide, scented; tepals 125 × 83mm, margins slightly ruffled, tips recurved. Lvs 170 × 56mm. Stems 0.95m, green, with darker markings at base, with up to 5 fls.

‘Candy Club’ VIII(a-b/b-c) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2008
Inside of tepals strong purplish red (63A); margins and base yellowish white (155D). Outside of inner tepals light purplish pink (63D) on a paler ground. Spots few, darker than strong purplish red (63A); papillae present; nectaries green; pollen red-brown; stigma dark purple. Fls 270mm wide; tepals of medium length to long and of narrow to medium width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and broad. Stems 0.95–1.2m, green with darker markings, with up to 7 fls. Very late.

‘Candy Floss’ VII(b/-)

I: H.W. Hyde and Son, 2009

Inside of tepals white tinged blush-pink; outer tepals soft, pale, apple-green along and beside midveins in basal three-quarters. Spots carmine-pink rimmed rose-pink, few, quite large, sprinkled towards the centre of the basal third of outer tepals. Papillae absent; spots absent from smaller, inner tepals; a few stamens and a little russet-crimson pollen present; style apple-green; stigma blush-pink. Buds apple-green. Flowers fully double; outer tepals well-formed, margins slightly wavy; inner tepals shorter, narrower and heavily ruffled; most tips recurved. Leaves abundant. Stems sturdy, with up to 15 fls. More vigorous and easier to grow than some double Orientals.

Exhibited by H.W. Hyde and Son at RHS Chelsea Flower Show, 2009. It is not possible to register this cultivar epithet as it is in use already for another lily (ICNCP, 2004: Art. 27.1).

‘Capuchino’ I(a-b/c)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003

N: Fa Q.J. Vink en Zn BV, 2003

REG: Mak Breeding Rights BV, 2008

Inside of tepals greenish white (155C); margins and base dark red (59A); midveins purple-red at base. Outside of tepals greenish white (155C), with spots darker than dark red (59A). Papillae present; nectaries green; pollen red-brown; stigma purple. Buds very hairy. Fls 150mm wide; tepals short to very short and of narrow to medium width, margins slightly to very slightly ruffled, tips slightly recurved. Lvs of medium length, and of narrow to medium width. Stems 0.8–1.1m, green with much darker markings, with few to average no. of fls. Early to mid-season.

Published in *The Lily Nook cat.*, 2009

‘Carracci’ VII(a/c-b)

Seedling 00-210 X seedling 01-31

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals strong purplish red (61B), shading to deep purplish red (61A) towards margins; outer tepals yellowish white (155D) at base and brilliant greenish yellow (3B) above nectaries; throat yellowish white (155D) shading to light greenish yellow (3D). Outside

of outer tepals deep purplish red (61A); yellowish white (155D) at base; midribs strong yellow-green (144A) at base, shading to light olive (152B), and dark green (132A) tinged yellowish white (155D) at top. Outside of inner tepals strong purplish red (64B/C); base yellowish white (155D); midribs light yellow-green (144D) at base shading to dark greenish yellow (152D). Spots and papillae dark red (59A); nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma pale yellow-green (193D). Fls 230mm wide; tepals 130 × 80mm, margins smooth, tips recurved. Lvs 20 × 70mm. Stems 1.2m, green with darker markings, with up to 5 fls.

Family of 16th-century Bolognese painters.

‘Cartesio’ VIII(a/b)

Seedling BG 97-29 X seedling AG 97-15

H: Vletter en Den Haan Beheer BV, 2002 **N:** Vletter en Den Haan Beheer BV, 2006 **REG:** Vletter en Den Haan Beheer BV, 2010

Inside of tepals greenish yellow on opening, maturing to brilliant yellow (7A); margins at tips vivid yellow (12A); throat pale greenish yellow (1D). Outside of tepals light greenish yellow (5C), shading to vivid yellow (12A) towards margins; midribs of outer tepals light yellow-green (144D) at base and top; midribs of inner tepals strong yellow-green (144B) at base. Spots and papillae absent; nectaries strong yellow-green (144B); pollen brownish orange (N170A) to moderate reddish orange (N172A); stigma deep purplish red (59B). Fls 180mm wide; tepals 120 × 65mm, margins smooth, tips slightly recurved. Lvs 130 × 28mm. Stems 1.4m, green, with up to 4 fls.

‘Cascade’ VII(a-b/b)

‘Star Gazer’ X un-named seedling

H: J.A. Vletter en Gebr. and Floris, pre-1979

G: J.A. Vletter en Gebr. and Floris, 1979

N: J.A. Vletter en Gebr., pre-1991

Tepals strong purplish pink (63C), shading to strong purplish red (63A) beside and along midveins; margins and tips white; midribs strong purplish red (63A). Spots numerous, dark pink or red-purple, elliptical, 2.5 × 1mm; papillae present; filaments white at base, pale green at top, 80mm long; anthers dark pink; pollen dark brown; style up to 100mm, pale to mid-green; stigma purple. Buds strong purplish pink (63C), 140mm long. Fls 200mm wide, strongly scented; tepal margins ruffled; pedicels mid-green, 100mm long, slightly curved. Lvs 180 × 55mm, moderate olive-green (137A). Stems 1.4m, light yellow-green (142C) with darker markings.

This cultivar epithet cannot be registered as it is in use for another lily (ICNCP, 2004: Art. 27.1). However, as this lily was granted a U.S. Plant Patent under the name ‘Cascade’ in 1993, this epithet remains the accepted name (*ibid.*: Art. 24.4).

‘Caselle’ I(a/b-c)

Seedling AM 96-18 × seedling RS 00-14

H: Vletter en Den Haan Beheer BV, pre-2006**N:** Vletter en Den Haan Beheer BV, 2006**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals whiter than yellowish white (NN155A); throat yellowish white (NN155A).

Outside of tepals yellowish white (NN155A); midribs light yellow-green (145C). Spots and papillae absent; nectaries strong yellow-green (144A); pollen dark orange (N163A); stigma light yellow-green (145D) with deep purplish red (59B). Fls 190mm wide; tepals 100 × 65mm, margins smooth, tips straight. Lvs 145 × 20mm. Stems 1.3m, green, with up to 7 fls.

‘Catania’ VII(b-a/b)

Seedling RM 98-28 × seedling 01-203

H: Vletter en Den Haan Beheer BV, pre-2004**N:** Vletter en Den Haan Beheer BV, 2004**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals strong purplish red (60D), shading to dark purplish pink (186C) towards margins; midveins strong purplish red (60B); throat yellowish white (155D). Inside of outer tepals strong purplish red (60D/61B), shading to vivid purplish red (61C) towards margins; midveins deep red (53A). Outside of outer tepals deep purplish red (61A), tinged yellowish white (155D) beside base of midribs; midribs light olive (152A). Outside of inner tepals strong purplish red (60D) in upper two-thirds, white (NN155C) in basal third; midribs light yellow-green (142C), tinged yellowish white (155D) at top. Spots dark red (59A); papillae strong purplish red (61B); nectaries strong yellow (N144B) rimmed brilliant greenish yellow (4A); pollen strong orange (N25A); stigma pale green (190A). Fls 215mm wide, scented; tepals 130 × 70mm, margins ruffled, tips slightly recurved. Lvs 210 × 50mm. Stems 1m, green, with up to 4 fls.

‘Catina’ VIII(a-b/b) OT

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006**N:** Testcentrum voor Siergewassen BV, 2006**REG:** Testcentrum voor Siergewassen BV, 2008

Inside of tepals vivid yellow (12A) to brilliant yellow (12B), shading to pale greenish yellow (9D) towards tips; younger fls paler. Outside of inner tepals light greenish yellow (8C). Spots absent; papillae present; nectaries green; pollen reddish brown; stigma purple. Tepals of medium length and width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length and medium width. Stems 1.4m, green with faint darker spots and stripes, with up to 5 fls. Late to very late.

‘Ceres’ I(a/b)

Tepals rose-pink. Spots and papillae absent; anthers blackish brown. Fls double; tepal margins mostly smooth, sometimes slightly wavy and nicked; tips slightly recurved. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009External images: *The Lily Nook cat.*, 2009**‘Cereza’ VIII(a/b) LA**

Asiatic seedling × Div. VIII LA seedling

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2008

Inside of inner tepals moderate reddish orange (171A) in upper half, moderate red (180A) in basal half; throat moderate yellowish pink (179D). Inside of outer tepals slightly more red than moderate red (180A), shading to moderate reddish orange (171A) towards margins. Outside of outer tepals moderate red (181B), shading to 179B towards margins; margins pale yellow (8D) at base; midribs strong yellow-green (143C) at base, tinged at top with brilliant yellow (8A) and strong orange (N25A). Outside of inner tepals moderate red (180B), shading to 179B towards margins; margins moderate yellowish pink (N170D) at base; midribs strong yellow-green (144C) at base. Spots and papillae dark red (59A); nectaries moderate olive-green (146A); pollen moderate reddish brown (176B); stigma vivid reddish orange (N30A). Fls 200mm wide; tepals 120 × 60mm, margins smooth, tips recurved. Lvs 170 × 20mm. Stems 1.3m, green, with darker markings at base, with up to 8 fls.

‘Checkpoint’ VI(c/b)

Parentage unknown

H: B. Mićulka, pre-2007 **N:** B. Mićulka, pre-2007**I:** B. Mićulka, 2007 **REG:** B. Mićulka, 2007

Inside of tepals copper, with paler tips and midveins; throat brownish apricot with carmine veining.

Outside of tepals conspicuously veined carmine; midribs darker and brownish. Spots absent; nectaries green; pollen brown; stigma green. Fls 120mm wide; tepals 115 × 40mm, margins slightly ruffled, tips recurved; pedicels 80mm, spotted red-brown, slender. Lvs 125 × 19mm, dull, mid-green. Stems 1.25m, with numerous red-brown spots. Mid-July.

‘China Pink’ VII(a/b)

A temporary name for ‘Pascoli’ prior to registration.

‘China Red’ VII(b/b)

‘Con Amore’ × ‘Acapulco’

H: S. Zheng, 2002 **G:** S. Zheng, Mrs J. Huang, F. Luo, Mrs L. Shi, H. Zhang, 2006 **N:** S. Zheng, Mrs J. Huang, F. Luo, Mrs L. Shi, H. Zhang, 2007 **REG:** S. Zheng, Mrs J. Huang, F. Luo, Mrs L. Shi, H. Zhang, 2009

Inside of tepals pinkish red; darker along and beside midveins; throat yellow and green. Outside of tepals pink; midribs whitish in basal two-thirds, pinkish red in upper third. Spots and papillae red; spots fairly large, evenly distributed over basal third to half of each tepal; nectaries green; pollen brown; stigma purple. Fls 220mm wide, strongly scented; tepals 142 × 59mm, margins slightly ruffled, tips slightly recurved. Lvs 143 × 32mm, whorled. Stems 1.05m, green, with up to 5 fls. Early (May). Diploid, through reduction from triploid.

The colour red in China represents success, good luck and happiness.

‘Circus Crowd’ I(b-c/c)

N: Mak-Leek, Inc., c.2008

Tepals bronze-orange, semi-reflexed; inside of each tepal with a golden yellow ray. Stems 1.2m. Mid-season.
Published in *The Lily Nook cat.*, 2009

‘Coastline’ I(a/b-c)

N: Fa Q.J. Vink en Zn BV, c.2008

Tepals pale yellowish green. Spots dark burgundy-red, coalescing in basal two-fifths on inside of each tepal, almost obliterating underlying colour; becoming progressively less dense along margins towards apex and towards central portion of upper half of each tepal. Pollen orange; stigma dull purplish pink. Tepals broad, margins very slightly ruffled, tips slightly recurved. Stems 0.9m. Mid-season. Distribution and colour of spots similar to ‘Burgundy Splash’.
Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009**‘Color Cosmetics’ VII(a/b)**

‘Sorbonne’ × ‘Acapulco’

H: Q.F. Cheng, 2004 G: Q.F. Cheng, 2008

N: Q.F. Cheng, 2008 REG: Yongkang Zhejiang Jiangnan Lily Breeding Co. Ltd, 2008

Syn. ‘Rouge’

Inside of tepals purplish pink in centre and at base, and along and beside midveins, shading to rose-pink towards margins and tips; throat pale yellow. Outside of tepals pink; midribs pale green. Spots few, large, sometimes slightly raised, dark purplish pink, between darker centre and paler margins in basal third to half of each tepal; papillae present over basal quarter of each tepal; nectaries green; pollen reddish brown; stigma white. Fls 200mm wide, strongly scented; tepals 100 × 55mm, margins slightly ruffled, tips slightly recurved. Lvs 125 × 35mm, scattered. Stems 1.05m, pale green.

External web links: www.lanlilium.com (image)**‘Contigo’ VII(a-b/b-c)**

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006

N: Testcentrum voor Siergewassen BV, 2006

REG: Testcentrum voor Siergewassen BV, 2008

Tepals greenish white (155C); midveins creamy white at base. Spots absent; papillae present; nectaries green; stigma grey with small purple spots. Fls sterile; tepals of medium length and of medium width to broad, margins ruffled, tips slightly recurved. Lvs of medium length and broad to very broad. Stems 1.05m, green with darker spots and stripes, with up to 5 fls. Mid-season.

‘Cosimo’ VIII(a/b) LA

Seedling BG 97-29 × seedling AW 98-10

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals pale yellow-green (155A), basal half of midveins 4D from above nectaries; young fls darker with outer tepals tinged red-brown at base and along midveins; throat pale yellow-green (155A). Outside of outer tepals yellowish white (155D); margins pale

greenish yellow (2D); midribs light yellow-green (144D) at base shading to pale greenish yellow (2D). Outside of inner tepals pale yellow-green (155A); midribs strong yellow-green, 144B at base shading to N144D. Spots and papillae absent; nectaries strong yellow-green (143A); pollen moderate reddish brown (175A); stigma deep purplish red (59B). Fls 200mm wide, scented; tepals 120 × 60mm, margins smooth, tips slightly recurved. Lvs 120 × 28mm. Stems 1.4m, green, with up to 4 fls.

‘Cunera’ VII(a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2006

N: Vletter en Den Haan Beheer BV, 2006

REG: Vletter en Den Haan Beheer BV, 2008

Tepals, including throat, white (NN155D); basal two-thirds of midveins light yellow-green (150D); midribs of outer tepals strong yellow-green at base (143C) and top (143A/C), light yellow-green (150D) between; midribs of inner tepals strong yellow-green at base (143C) and top (144C), pale yellow-green (155A) between. Spots numerous, light yellow-green (150D) and white (NN155D); papillae white (NN155D); nectaries light yellow-green (144D); pollen dark orange (N163A); stigma dark purplish red (N79B). Fls 270mm wide, scented; tepals 165 × 83mm, margins smooth, tips recurved. Lvs 233 × 50mm. Stems 1.3m, green, with up to 7 fls.

‘Dalila’ I(b-a/b)

Un-named seedling × (‘Selma’ × ‘Paola’)

H: De Jong Beheer BV, pre-2006 N: De Jong Lelies

Holland BV, 2006 REG: De Jong Lelies Holland BV, 2008

Inside of tepals strong purplish red (63A); midveins moderate purplish red (58A); throat strong purplish red (64C). Outside of outer tepals moderate purplish red (58A); midribs strong yellow-green (143C) at base, shading to moderate olive-brown (199A), and moderate olive-green (146A) at top. Outside of inner tepals strong purplish red (59D); midribs strong yellow-green (143C) at base. Spots and papillae absent; nectaries strong yellow-green (143A); pollen dark reddish orange (175C); stigma dark red (59A). Fls 160mm wide; tepals 93 × 50mm, margins smooth, tips slightly recurved. Lvs 142 × 20mm. Stems 1.15m, dark, with up to 15 fls.

‘Dark Sensation’ VII(a/-)

Parentage unknown

N: H.P. Imanse BV, 2007 REG: H.P. Imanse BV, 2008

Inside of tepals deep red (60A), shading to strong purplish red (60B) towards margins and to almost white at tips, becoming more blue and paler with age. Outside of inner tepals dark purplish pink (186C); midribs more red. Spots darker than deep red (60A); papillae present; nectaries dark purple-red; pollen orange-brown; stigma grey-green. Tepals of medium length and of medium width to broad, margins ruffled, tips recurved. Lvs of medium length and broad. Stems of average height, green with some darker markings, with few to average no. of fls. Mid-season to late.

‘Diena’ I(b/c)

‘Dzintars’ × seedling 170-F

H: Ā. Zorgevics, 1990 **G:** Ā. Zorgevics, 1993 **N:** Ā. Zorgevics, 1994 **I:** University of Latvia Botanical Garden, 2000 **REG:** Ā. Zorgevics and University of Latvia Botanical Garden, 2008

Inside of tepals vivid yellow (14A); throat yellow-orange. Outside of tepals yellow-orange; midribs green. Spots and papillae absent; nectaries clear; pollen brown; stigma pale brown. Fls 180mm wide, not scented; tepals 90 × 33–36mm, margins smooth, tips strongly recurved. Lvs 100 × 20mm. Stems 1m, green, with up to 7 fls. Late July.

Published in *Lilium Balticum. Latvijas liliju selekcionāru izaudzētās lilijas*, No. 2 (1995)

The name means “day” in Latvian.

‘Dimension’ I(a/c)

N: Fa Q.J. Vink en Zn BV, c.2005

Tepals brownish burgundy-red with a black-cherry sheen; midveins slightly paler, shaded pinkish red. Spots and/or short brushmarks sparse and inconspicuous, almost black, on basal third of each tepal; pollen orange; stigma brownish black. Tepals of medium width, margins very slightly ruffled, tips becoming strongly recurved. Stems 0.9m. Mid-season. External images: *The Lily Nook cat.*, 2009

‘Dimples’ I(a/-)

N: Columbia-Platte Lilies, Inc., c.2008

Tepals dark rose-pink; inside of each tepal heavily spotted creamy yellow towards centre. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009

‘Disegno’ I(a-b/b-c)

Parentage unknown

H: Mak Breeding BV, pre-2007 **N:** Mak Breeding BV, 2007 **REG:** Mak Breeding Rights BV, 2009

Inside of tepals moderate red (N34A) on an orange ground, paler towards base; midveins purple-red at base shading to red. Outside of inner tepals moderate red (N34A). Spots and papillae few; nectaries purple-red; pollen red-brown; stigma purple. Fls 140–150mm wide; tepals short and of narrow to medium width, margins (very) slightly ruffled, tips (very) slightly recurved. Lvs of medium length, and of narrow to medium width, dark green. Stems 1.1m, green, with up to 7 fls. Early to mid-season.

‘Divoženka’ VI(b-c/b)

(Pink-flowered seedling hybridized by Freimann × First Love Group) × unknown

H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007

I: B. Mičulka, 2007 **REG:** B. Mičulka, 2007

Syn. ‘Veletoč’

Inside of tepals creamy white; margins conspicuously veined pink; throat pale golden yellow. Outside of tepals pink-veined; midribs richer pink. Spots absent; nectaries greyed olive-green; pollen ochre-apricot; stigma dark brown. Fls 130mm wide; tepals 120 × 80mm, tips recurved; pedicels 95mm, with numerous dark brown spots. Lvs 135 × 12mm, dull, pale green. Stems 1.25m, stout, with a slight, speckled bluish tinge. Early July.

‘Double Sensation’ I(a/b)

Parentage unknown

N: W. van Lierop en Zn BV, 2008 **REG:** W. van Lierop en Zn BV, 2009

Tepals dark red (59A) to deep purplish red (59B), inside of tepals darker towards margins and with pale yellow-green (155A) patch towards base. Spots few; papillae present; nectaries green; pollen lacking; stigma green. Fls double; tepals short and of medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length and breadth. Stems tall, green with darker markings, with few fls. Mid-season to late.

‘Dragon’s Gowns’ VIII(b/a)

regale × ‘Yelloween’

H: Q.F. Cheng, 2004 **G:** Q.F. Cheng, 2008

N: Q.F. Cheng, 2008 **REG:** Yongkang Zhejiang Jiangnan Lily Breeding Co. Ltd, 2008

Inside of tepals, including throat, yellow. Outside of tepals, including midribs, pale yellow. Spots and papillae absent; nectaries green; pollen saffron-yellow; stigma green. Fls 180mm wide, slightly scented; tepals 110 × 50mm, margins slightly ruffled, tips slightly recurved. Lvs 115 × 9mm, scattered. Stems 1.8m, green.

External web links: www.lanlilium.com (image)

‘Dream Julius’ I(a/c)

Un-named, yellow-flowered, pollen-free seedling

× ‘Mint Fudge’

H: T. Zimmerman, 2003 **G:** T. Zimmerman, 2005

N: T. Zimmerman, 2008 **REG:** T. Zimmerman, 2008

Inside of tepals golden yellow with large, dark burgundy-red patch with feathered margins below centre; throat yellow. Outside of tepals brilliant greenish yellow (7C). Spots large, fairly sparse, dark purple (83A), mostly in basal fifth of each tepal between throat and base of patch; nectaries yellow; pollen absent; stigma pale yellowish beige. Fls 110 mm wide, not scented; tepals 65 × 30mm, margins smooth, tips straight. Lvs 110 × 15mm, scattered. Stems 1m, green, with up to 6 fls. Early July.

‘Dreamweaver’ VIII(b-a/b) LO

Div. VII (Oriental hybrid) seedling × seedling

LO 93003 ZB T

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2008

Inside of tepals strong purplish red (inner 64C, outer 60D); midveins of inner tepals 64B; those of outer 61B; margins and throat yellowish white (155D). Outside of outer tepals dark purplish pink (186C); margins and base yellowish white (155D); midribs strong yellow-green at base (144A) and top (143A). Outside of inner tepals deep purplish pink (64D); margins yellowish white (155D); midribs strong yellow-green (143C) at base. Spots absent; papillae strong purplish red (64B); nectaries strong yellow-green (144B); pollen dark orange (N163A); stigma greenish white (192D). Fls 215mm wide, scented; tepals 137 × 63mm, margins slightly ruffled, tips recurved. Lvs 220 × 32mm. Stems 1.3m, green, with up to 4 fls.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

'Dreamworks' VIII(a/b) LA

Seedling BW 96-19 × seedling AW 99-10

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Tepals yellowish white (155D), outside of outer tepals strong yellow-green at base (141D) and tips (143C); tips tinged strong purplish red (60B); outside of inner tepals tinged strong yellow-green (143B) at base; throat light yellow-green (145C). Spots and papillae absent; nectaries strong yellow-green (143A); pollen brownish orange (N167B); stigma moderate yellow-green (139D) with moderate purplish red (58A) stripes. Fls 160mm wide; tepals 110 × 57mm, margins smooth, tips slightly recurved. Lvs 135 × 25mm. Stems 1m, green, with up to 4 fls.

'Easter Charm' VIII(b/a) LT**H:** L.M. Collicutt, pre-2008

Tepals purplish rose-pink; towards tips fading to pinkish white and tinged pale yellowish green. Spots and papillae absent; filaments greenish yellow; pollen rich orange. Buds dark purple. Tepal margins smooth, tips recurved. Stems 0.9–1.2m. Mid-season to late. Published in *The Lily Nook cat.*, 2009

'Easter Morn' VIII(b-c/a) LT**H:** L.M. Collicutt, pre-2008

Inside of tepals white, shading to purplish pink towards margins and along midveins; basal one-third flushed pink overlaid pale greenish yellow. Outside of tepals darker purplish pink at base, colour becoming paler and less solid towards tips; midribs dark violet. Spots and papillae absent; filaments yellowish green; pollen rich, dark brown. Fls of strong substance; tepal margins smooth, tips recurved. Stems 1.07m. Mid-season to late. Disease resistant. Tetraploid. Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

'East of Eden' II(c/d)

Un-named seedling × unknown

H: D. Hercbergs, pre-2003 **G:** D. Hercbergs, 2003**N:** D. Hercbergs, 2004 **I:** D. Hercbergs, 2007**REG:** D. Hercbergs, 2008

Tepals deep purplish red (61A). Spots deep purplish red (61A) rimmed yellowish white (155D); pollen vivid yellowish pink (28A); stigma dark purplish red (N79A). Fls 48mm wide, slightly scented; tepals 35 × 14mm, margins smooth, tips strongly recurved. Lvs 140 × 40mm, whorled. Stems 1.2m, dark green to nearly black, with up to 18 fls. Mid-season.

'East Red' VII(b/b)

'Con Amore' × 'Acapulco'

H: S. Zheng, 2002 **G:** S. Zheng, 2006 **N:** S. Zheng, 2007 **REG:** S. Zheng, 2009

Tepals rose-pink; midveins darker; midribs yellowish green; throat green. Spots and papillae red, evenly distributed over basal half of each tepal; nectaries green; pollen orange-brown; stigma purple. Fls 198mm wide, strongly scented; tepals 125 × 48mm, margins slightly ruffled, tips straight. Lvs 126 × 32mm, whorled. Stems 1.05m, green, with up to 5 fls.

Early (May). Diploid, through reduction from triploid. The colour red in China represents success, good luck and happiness.

'Edinburgh' VII(b-a/b)

Seedling 02-308 × seedling 93-14

H: Vletter en Den Haan Beheer BV, pre-2006**N:** Vletter en Den Haan Beheer BV, 2006**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals, including throat, yellowish white (155D); upper half flushed strong purplish pink (63C). Outside of outer tepals yellowish white (155D), half of each tepal flushed pale purplish pink (65D); midribs strong yellow-green at base (144C) and top (144A/B), light yellow-green (145D) between. Outside of inner tepals yellowish white (155D), tinged pale purplish pink (65C) at tips; midribs strong yellow-green (144C) at base shading to strong yellow (153D). Spots and papillae yellowish white (155D); nectaries light yellow-green (145B); stigma light grey (188D). Fls 250mm wide, scented; tepals 150 × 78mm, margins ruffled, tips recurved. Lvs 230 × 55mm. Stems 1.15m, green, with up to 4 fls.

'El Divo' VIII(a-b/b) LA

Parentage unknown

H: Mak 't Zand BV, pre-2006 **N:** Mak Breeding BV, 2006 **REG:** Mak 't Zand BV, 2008

Inside of inner tepals more orange than vivid yellow (12A). Inside of outer tepals vivid yellow (12A/13A). Outside of inner tepals vivid yellow (12A). Spots few, along basal margin of each tepal; papillae few; nectaries yellow-green; pollen pale brown; stigma purple-red with small spots. Tepals of medium length and of medium width to broad, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length to long, and of medium width. Stems 1.25m, green with faint darker spots and stripes, with up to 10 fls. Mid-season.

'Electric Yellow' I(a/b-c)**N:** Gebr. N. & B. Laan en Zn, c.2008

Inside of tepals bright golden yellow extending to tips; broad white margin either side of tip in upper half of each inner tepal, but extending from near base of each outer tepal. Spots dark brown, rather large, elliptic, evenly distributed in basal half of each tepal, a few scattered on or near midvein in middle third of each tepal; pollen brownish yellow; stigma dull, slightly greenish yellow. Tepals of narrow to medium width, margins smooth, tips slightly recurved. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009External images: *The Lily Nook cat.*, 2009

'Elise' (Mak Leek) VIII(c/-)**H:** Mak Leek, Inc., pre-2008

Inside of tepals mango-orange passing to apricot-yellow with age; midveins darker orange. Brushstrokes darker orange but faint, delicate, short, curving away from either side of midveins, mostly distributed over basal half of inner tepals. Outside of tepals paler, soft apricot orange-yellow. Filaments slender, pale greenish yellow; style yellowish green; stigma apple-green; pollen blackish brown. Tepal margins smooth. Leaves broad, dark green. Stems 1.2–1.5m. Late July to early August (Washington, USA).
Published in *The Lily Nook cat.*, 2009

It is not possible to register either Weinreich's 'Elise' (hybridized pre-1985), or Mak Leek's cultivar epithet as it is too similar to, and likely to cause confusion with, 'Elisa', which was registered by J.G. Smith in 1984 (ICNCP, 2004: Art. 19.25).

'Ellington' VII(b-a/b)**Div.** VII (Oriental hybrid) seedling × **Div.** VII seedling**H:** De Jong Beheer BV, pre-2005 **N:** De Jong Lelies Holland BV, 2005 **REG:** De Jong Lelies Holland BV, 2008

Tepals, including throat, whiter than white (NN155D). Outside of outer tepals tinged moderate red (181A) beside base of midribs; midribs light yellow-green (144D) at base and top, top tinged strong yellow-green (144A). Midribs of inner tepals moderate red (181A), strong yellow-green (144C) at base, light yellow-green (150D) at top. Spots and papillae white (NN155D); nectaries strong yellow-green (144C); pollen strong orange (25A); stigma light grey (191C). Fls 215mm wide, scented; tepals 130 × 70mm, margins slightly ruffled, tips slightly recurved. Lvs 200 × 50mm. Stems 1.2m, green, with up to 4 fls.

'Energetic' VII(a/b-c)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007**N:** Testcentrum voor Siergewassen BV, 2007**REG:** Testcentrum voor Siergewassen BV, 2008

Inside of tepals deep purplish pink (N66D) on a pale ground, shading to white towards base. Outside of inner tepals slightly paler than deep purplish pink (N66D). Spots white on a pink ground; papillae present; nectaries green; pollen reddish brown; stigma grey-green. Tepals of medium length to long and of medium width to broad, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and broad. Stems of average height to tall, green with darker stripes and spots, with few to average no. of fls. Mid-season to late.

'Enigma' I(a/b)

Unknown × 'Tango Nocturno'

H: A. Krūmiņš, 1997

Tepals greenish cream towards tips; throat pale greenish yellow; midribs beige. Spots dark mahogany-brown, very numerous, closely arranged and coalescing in places over lower two-thirds of each tepal, becoming progressively more scattered and solitary towards tips; nectaries pale violet, frosted; pollen and stigma brown. Fls 120mm wide; tepals 70 × 26mm, margins slightly

ruffled, tips recurved; pedicels 45mm, slender, spotted brown. Lvs 80 × 8mm, dull, pale green. Stems 0.85m, green; infl. short. Early July.

It is not possible to register this cultivar epithet as it is in use already for another lily (ICNCP, 2004: Art. 27.1).

'Etchings' I(a/b-c)

('Sterling Star' × 'Connecticut King') × ('Connecticut Queen' × (Pastel Group: upright, white-flowered × ('Connecticut King' × brushmarked seedling)))

H: J. Freeman-McRae, pre-1988 **N:** J. Freeman-McRae, c.1988 **I:** Columbia-Platte Lilies, Inc., 1988

Inside of inner tepals glossy white, sometimes pale greenish yellow (9D) on opening. Basal one-third of each tepal marked with 2–4, 10–20 × 10–30mm, deep purplish red (71A) to strong purplish red (71B) brushmarks in an etched pattern; papillae very small, deep purplish red (71A) to strong purplish red (71B), 10–20 at base of each tepal parallel to nectaries; nectaries green; filaments pale greenish yellow (9D), 50mm; anthers and pollen strong brown (172A); style 50mm; stigma pale green tinged with pale plum. Buds up to 90mm long, obtuse, ovoid. Fls 150–190mm wide, not scented; tepals 25–35mm wide, tips recurved; pedicels 80–100mm, stout; occasionally with secondary buds. Lvs 80–120 × 10–20mm, glossy, leathery. Stems 0.9m, green with darker markings, with up to 8 fls in a racemose infl. 200–250mm long. Mid-season.

'Eton' VII(b-a/c-b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2006**N:** Vletter en Den Haan Beheer BV, 2006**REG:** Vletter en Den Haan Beheer BV, 2008

Tepals, including throat, yellowish white (155D); midribs of outer tepals strong yellow-green, 144C at base and 144A at top; midribs of inner tepals light yellow-green (144D) at base and strong yellow-green (144C) at top. Spots absent; papillae yellowish white (155D); nectaries light yellow-green (144D); pollen dark orange (N163A); stigma greyish purplish red (N77D). Fls 240mm wide, scented; tepals 145 × 83mm, margins ruffled, tips recurved. Lvs 203 × 53mm. Stems 1.15m, green with some darker markings, with 1 or 2 fls.

'Etrusk' I(a/b-c)

((('Heureka' × 'Sterling Star') × unknown) × unknown

H: M. Tlach, pre-2007 **N:** M. Tlach, pre-2007**I:** M. Tlach, 2007 **REG:** M. Tlach, 2007

Inside of tepals creamy white shading to pale pink at centre; margins yellowish; throat white. Outside of tepals brownish pink. Spots large, oval, violet, moderately dense over two-thirds of each tepal; papillae absent; nectaries brownish green, frosted white; pollen orange-brown; stigma violet. Fls 115mm wide; tepals 70 × 35mm, tips slightly recurved; pedicels 60mm, stout. Lvs 95 × 14mm, dull, mid-green. Stems 0.85m, with few violet spots; infl. long. Late June.

'Eudoxia' VIII(b-a/b) OT

Seedling PJ 96-08 × seedling RH 97-07

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals pale orange-yellow (23D) in upper half, light orange-yellow (23C) in basal half; midveins vivid yellow (15A) in basal half from above nectaries; throat pale orange-yellow (24D). Inside of outer tepals brilliant orange (25C), shading to light orange-yellow (23C) towards margins; midveins strong orange-yellow (17A) above nectaries. Outside of outer tepals light yellow (18B); midribs strong yellow-green at base (144A) and top (143B), strong yellow (N144B) between. Outside of inner tepals light orange-yellow (22D), shading to pale yellow (19C) beside midribs; midribs strong yellow-green (144A) at base, shading to brilliant yellow-green (150C). Spots absent; papillae vivid reddish orange (30B); nectaries strong yellow-green (144B); pollen strong reddish orange (169A); stigma pale green (189B). Fls 230mm wide, scented; tepals 135 × 65mm, margins ruffled, tips recurved. Lvs 220 × 30mm. Stems 1.3–1.5m, dark green, with up to 5 fls.

eupetes J.M.H. Shaw IX(c/b)**N:** J.M.H. Shaw, 2008 **I:** B. and S. Wynn-Jones, 2006

Inside of tepals maroon-red; outside of outer tepals greenish towards tips; midribs of inner tepals green. Spots absent; papillae and nectaries greenish yellow; filaments bright red; pollen orange; style maroon; stigma white. Inner tepals 40 × c.10mm, outer tepals 40 × 16mm, margins smooth, tips very slightly recurved; pedicels maroon. Lvs (3-) 5–6.5 × 10–16mm, scattered, light green with reddish midrib, axils of upper lvs with 1–3 small, green, conical bulbils. Stems to 0.45m, hairless, green, usually with solitary fl. Bulb yellowish white. Vietnam, growing epiphytically in mosses on large forest trees at altitudes of 1,900–2,000m.

Description of this new species collected by B. and S. Wynn-Jones in 2006, published in *The Plantsman*, New Series Vol. 7, Pt 1, pp. 39–41, RHS, March 2008. Name derived from the Greek word meaning "flying well", which refers to the apparently unique mode of dispersal of the bulbils which develop in the leaf axils. A leaf, with a bulbil attached, withers to a circular shape which spins through the air when the leaf detaches from the dying stem.

'Eva Steinbrück' VI(b/a)

Unknown × ((Green Magic Group × 'Bright Star') × mixed pollen from an unknown lily, 'Damson' and 'Erfordia')

H: R. Widerski, pre-2007 **N:** R. Widerski, pre-2007**I:** R. Widerski, 2007 **REG:** R. Widerski, 2007

Tepals yellow-orange; tips orange; throat pale yellow-orange. Nectaries green; pollen brown; stigma green. Fls 140mm wide; tepals 145 × 56mm, tips recurved; pedicels 70mm, stout; with secondary buds. Lvs glossy, mid-green. Stems 1.2m, green, with average no. of fls. Early July.

'Evidenza' VIII(b/a) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006**N:** Testcentrum voor Siergewassen BV, 2006**REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals and outside of inner tepals strong purplish red (60C/D); midveins greenish at base. Spots and papillae absent; nectaries pale green; pollen pale orange; stigma orange. Tepals of medium length to long and of medium width to broad, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length to long and of medium width. Stems 1.3m, green, with up to 3 fls. Late to very late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

'Falanghina' VII(a/b-c)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2003**N:** Marklily BV, 2003 **REG:** Testcentrum voor

Siergewassen BV, 2008

Inside of tepals deep red (53A/60A), paler towards margins, tips and base. Outside of inner tepals strong purplish red (60C). Spots numerous, darker than tepals; papillae present; nectaries dark purple; pollen red-brown; stigma grey-green. Tepals of medium length to long and of medium width, margins slightly ruffled to ruffled, tips recurved to strongly recurved. Lvs of medium length to long, and broad. Stems 1.45m, green with darker spots and stripes, with up to 6 fls. Late.

'Fire Mountain' I(a)**N:** Cebeco Lilies, Inc., c.2008

Tepals flame-red; inside of tepals shading to darker red towards central portion. Fls large. Stems 1.2m.

Mid-season. Tetraploid.

Published in *The Lily Nook cat.*, 2009**'Flashpoint' VIII(a-b/b) OT****H:** Mak 't Zand BV, pre-2006 **N:** Mak Breeding BV,2006 **REG:** Mak 't Zand BV, 2009

Inside of tepals deep red (60A/53A), shading to pale yellow (8D) at base, along margins and towards tips; midveins white at base shading to purple-red. Outside of inner tepals deep purplish pink (N66D), becoming paler towards pale yellow margins. Spots few, dark; papillae few; nectaries pale green; filaments greenish white; pollen pale brown; style pale apple-green; stigma grey-green speckled purple. Fls 200–250mm wide; tepals of medium length and width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and of medium width to broad, dark green. Stems 1.4m, green with darker spots and stripes, with up to 8 fls. Late. Prefers alkaline soil. Exhibited by H.W. Hyde and Son at RHS Chelsea Flower Show, 2009.

'Fleurise' VII(a/c-b)

Parentage unknown

H: R.W.J. van der Salm, pre-2007 **N:** Fa Van der Salm-De Goede, V.o.f., 2007 **REG:** Fa. Van der Salm-De Goede, V.o.f., 2009

Inside of tepals strong purplish red (61B) to vivid purplish red (61C), paler with age. Outside of inner tepals strong purplish red (63B). Spots and papillae present; nectaries green; pollen orange-brown; stigma grey-green. Fls 150–200mm wide; tepals of medium length and width, margins slightly ruffled, tips slightly recurved. Lvs of medium length, and of medium width to broad. Stems 1.15m, green with darker markings, with up to 6 fls. Early to mid-season.

'Frederique's Choice' VII(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2005 **N:** Liliande BV, 2005 **REG:** World Breeding BV, 2008

Inside of tepals white flushed deep purplish pink (N66D), greenish white (155C) towards base. Outside of inner tepals white flushed deep purplish pink (N66D). Spots numerous, white; papillae present; nectaries green; pollen orange-brown; stigma grey-green. Fls 280mm wide; tepals of medium length to long and of medium width to broad, margins ruffled, tips recurved. Lvs of medium length to long, and broad. Stems 1.05–1.15m, green with faint darker spots and stripes, with up to 7 fls. Mid-season to late. Named after the Dutch supermodel Frederique van der Wal.

'Zorina' was used as a temporary name prior to registration.

'Friso' VIII(b/b) OT

Parentage unknown

H: R.J. Griesbach, pre-2005 **N:** Lily Company BV, 2005 **REG:** Lily Company BV, 2008

Inside of tepals deep purplish red (59B) in an arc covering basal one-third to margins, rising to basal two-thirds towards centre; upper margins and tips pale yellow-green (155A); throat vivid yellow (15A) beside nectaries and pale greenish yellow (1D) above. Outside of outer tepals moderate purplish red (186B); midribs dark red (187B) at base shading to moderate purplish red (186A), strong yellow-green (144B) at top. Outside of inner tepals pale yellow-green (155A), shading to strong yellow-green (144C) beside midribs in basal one-quarter, dark purplish pink (186C) to moderate purplish red (186B) beside midribs in second basal quarter and extending to basal two-thirds, shading paler towards margins, upper third of midribs moderate purplish red (186B). Spots and papillae absent; nectaries strong yellowish green (N144A); pollen moderate reddish orange (N172A); stigma greyish reddish purple (N77C). Fls 190mm wide; tepals 140–150 × 40–60mm, margins smooth to slightly ruffled, tips recurved. Lvs 140–170 × 20–30mm, dark green. Stems 1.3m, green with darker markings, with 1 or 2 fls.

Named after the grandson, Friso Tjittes, of P. Groot of Lily Company BV

'Frosty Star' I(a/-)**N:** Columbia-Platte Lilies, Inc., c.2008

Tepals dark red; inside of each tepal with a central white flare, creating a six-pointed star. Spots black, sparse, near the edge of each white flare. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009

'Gambetta' VIII(b/c-b) OT

Seedling PG 94-43 × seedling RH 99-07

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals light greenish yellow (7D), tinged 6D at tips; midveins vivid reddish orange (30A); throat pale yellow-green (4D). Inside of outer tepals light greenish yellow (6D), at base beside nectaries yellowish white (155D) shading above nectaries from brilliant greenish yellow (6C) to vivid reddish orange (30A). Outside of outer tepals yellowish white (155B), shading to light greenish yellow (4C) towards margins; midribs strong yellow-green (144C) at base. Outside of inner tepals light greenish yellow (4C), yellowish white (155B) beside midribs; midribs strong yellow-green (144C) at base shading to pale yellow-green (4D). Spots absent; papillae vivid reddish orange (30A); nectaries strong yellow-green (143C); pollen greyish brown (166A); stigma greyish purplish red (N77B). Fls 260mm wide, scented; tepals 160 × 80mm, margins ruffled, tips recurved. Lvs 260 × 50mm. Stems 1.1m, green, with darker markings at base, with 1 or 2 fls.

'Garden Society' IV(c/d)*occidentale* × Div. IV (American hybrid)**H:** R.T. Whiteley, 2003 **G:** R.T. Whiteley, 2005**N:** R.T. Whiteley, 2006 **REG:** R.T. Whiteley, 2009

Inside of tepals brilliant orange-yellow (21B) at base, shading to vivid red (45A) at tips; throat strong yellow-green (141D). Outside of tepals strong orange-yellow (163B) at base, shading to strong orange (169C) at tips. Spots dark greyish purple (N186B), at base of each tepal; papillae absent; nectaries strong yellow-green (141D); pollen brownish orange (N170A); stigma dark greyish reddish brown (200A). Fls 70mm wide, not scented; tepals 75 × 18mm, margins smooth, tips strongly recurved; pedicels 200mm. Lvs 180 × 20mm, scattered, with single midvein. Stems 1.2m, moderate olive-green (137A), with up to 5 fls. July.

The Garden Society is a dining club for garden owners in England, Wales and the Channel Islands.

'Gaybird' II(c/d)**H:** E. Robinson, pre-2008

Inside of tepals raspberry-wine red towards tips and cream towards base. Spots dark maroon surrounded by a cream halo, numerous on basal half of each tepal, also along margins on outside of tepals. Stems 1.5m. Early.

Published in *The Lily Nook cat.*, 2009

‘Gerlach’ VI(b-c/d)

‘Mimosa Star’ × ‘Caramba’

H: A. Mego, pre-2008 **N:** A. Mego, 2008 **I:** A. Mego, 2008 **REG:** A. Mego, 2008

Inside of tepals apricot-orange. Outside of tepals apricot-green; midribs pale green. Raised, papillary, reddish brown dashes over one-third of throat; nectaries green; filaments pale green; pollen brown; style and stigma pale green. Fls 120mm wide; inner tepals 95 × 35mm; pedicels 100mm, green with inconspicuous brown spots; with secondary buds. Lvs 200 × 45mm, glossy, pale green. Stems 1.8m, pale green, with infl. of medium length. Mid-July.

‘Giacometti’ VIII(b-a/b) OT

Seedling PG 96-43 × seedling RH 94-01

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals brilliant yellow (10A); midveins 11A; throat pale yellow (11D). Inside of outer tepals light yellow (10B); midveins brilliant yellow (10A). Outside of outer tepals yellowish white (155B), shading to pale greenish yellow (10D) towards margins; midribs strong yellow-green (144C) at base and brilliant yellow-green (150C) at top, pale yellow (11C) between. Outside of inner tepals light greenish yellow (8C), shading to pale yellow (11D) towards tips and beside midribs; midribs tinged pale yellow-green (149D) at base. Spots absent; papillae vivid reddish orange (30A); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma greyish purple (N77A). Fls 180mm wide, scented; tepals 115 × 65mm, margins smooth, tips slightly recurved. Lvs 215 × 30mm. Stems 1.2m, green with some darker markings, with up to 4 fls. Named after the Swiss surrealist painter and sculptor who lived 1901–1966.

‘Gizmo’ VIII(b-a/b) LO

Seedling LO 97-11 × seedling RW 94-03

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Tepals, including throat, yellowish white (155D); midveins light greenish yellow (5D) above nectaries; midribs of outer tepals strong yellow-green at base (144C) and top (143A), light yellow-green (150D) between; midribs of inner tepals strong yellow-green (144C) at base. Spots absent; papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma pale yellow-green (157B). Fls 265mm wide, scented; tepals 160 × 70mm, margins slightly ruffled, tips recurved. Lvs 245 × 40mm. Stems 1.05–1.4m, green, with up to 4 fls. Prefers alkaline soil.

‘Glamour’ VII(a/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2008

Inside of tepals deep red (53A/60A), paler towards tips. Outside of inner tepals strong purplish red (60C/D). Spots and papillae present; nectaries green

to yellow-green; pollen orange-brown; stigma green, purple-spotted. Fls 250mm wide; tepals of medium length and breadth, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad. Stems 0.95–1.25m, green with darker markings, with up to 8 fls. Late.

‘Golden Belles’ VIII(b/b) AT

H: L.M. Collicutt, pre-2008

Inside of tepals greenish lemon-yellow; inside of outer tepals flushed pale brownish purple towards margins at first. Outside of outer tepals brownish purple with narrow margins of yellowish white; outside of inner tepals flushed pale brownish purple beside midribs in basal half, and with a narrow smudge of pale purplish brown beside midribs in upper half; midribs dark purplish brown. Spots blackish brown, tiny, round, sprinkled evenly over basal third of each tepal; papillae greenish sulphur white, very sparse and small; anthers greyed purple; stigma and upper half of style blackish brown. Buds dark chocolate-brown. Tepal margins smooth, tips recurved. Stems 0.9m. Mid-season to late. Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

It is not clear whether this lily is the same cultivar as ‘Golden Bell’ registered by C. Yerex in 1983, though that is described as having no spots.

‘Golden Drop’ VI(c/a)

Parentage unknown

G: L.C. Bligh, 2003 **N:** L.C. Bligh, 2007

REG: L.C. Bligh, 2008

Tepals, including throat, strong orange-yellow (24B), outside of tepals becoming paler, inside margins narrow, yellowish white (158C); midribs dark red (183B). Spots absent; nectaries green; pollen dark russet-brown; stigma pale green. Fls 150mm wide, strongly scented; tepals 150 × 55mm, margins smooth, tips slightly recurved. Lvs 150 × 11mm, scattered. Stems 1.57m, maroon with green flecks, with up to 15 fls. Late December (Tasmania).

‘Golden Stone’ VIII(a-b/b) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

N: Testcentrum voor Siergewassen BV, 2007

REG: Testcentrum voor Siergewassen BV, 2008

Inside of tepals brilliant yellow (7B); base of midveins yellow. Outside of inner tepals brilliant greenish yellow (7C). Spots darker than dark red (187B), along all margins and base of each tepal, including base of midveins; papillae present; nectaries yellow-green; pollen red-brown; stigma dark purple. Tepals of short to medium length and of medium width, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long and of medium width. Stems 1.75m, green, with up to 6 fls. Mid-season.

‘Good Question’ I(a/c-d)

Parentage unknown

H: B. Strohmman, pre-2008 **G:** B. Strohmman, pre-2008 **N:** B. Strohmman, c.2008 **I:** The Lily Nook, 2009
Upper half of tepals raspberry-pink, basal half of inside of each tepal milk-white, feathering to narrow margins of raspberry-pink and tapering towards midveins to form a large dome-shaped patch; midveins flushed pale creamy yellow in basal half. Spots and papillae absent; nectaries green; filaments pinkish, pollen rich russet-brown; stigma pale greenish yellow. Tepal margins smooth, tips strongly recurved. Stems 0.9m. Mid-season.
Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

‘Graffity’ I(a-b/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** Vink, 2003 **REG:** Mak Breeding Rights BV, 2008
Inside of tepals brilliant yellow (7A/B); margins and base dark red (59A) to deep purplish red (59B); base of midveins yellow-green with purple-red spots. Outside of tepals brilliant greenish yellow (7C) to light greenish yellow (7D). Spots darker than dark red (59A) to deep purplish red (59B); papillae present; nectaries yellow-green; pollen pale brown; stigma dark purple. Fls 180mm wide; tepals short and narrow, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs of medium length, and of narrow to medium width. Stems 0.8–1.1m, green with darker spots and stripes, with average no. of fls. Early to mid-season.
Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

‘Grandview’ I(a/c)

Un-named, small-, white-flowered, pollen-free sport of ‘Sunkissed’ O.P. × un-named, tall, exceptionally vigorous, large-, orange-flowered seedling of ‘Chinook’ O.P.

H: H. Mantel, pre-1986 **G:** H. Mantel, 1986

I: Mount Hood Lilies, Inc., c.1994

Tepals light orange-yellow (23C) to pale orange-yellow (23D), shading towards base to brilliant orange (25C) to light orange-yellow (25D); more peachy orange with age. Spots small, deep magenta, very lightly scattered over area parallel to nectaries; nectaries pale green to ivory, even pale orange in bright light, downy with short white hairs; filaments 40–70mm, light yellowish pink (27B); anthers tiny, 1–4mm, pale grey, indehiscent; pollen absent; style 40–90mm, pale yellowish pink (27C/D); stigma large, moderate purplish red (185C) to deep pink (185D). Fls 150mm wide, shiny, star-shaped, not scented; tepals 30mm wide, tips slightly recurved; pedicels 70–100mm, ascending, dark green flushed deep plum, appearing almost black; infls from large bulbs bear numerous secondary buds on vertical pedicels. Lvs 70–120 × 20–40mm, glossy, mid- to dark green. Stems 1.25–1.75m, with up to 12 fls. Mid-season.

Griesbach Tetra Gold Group VI(c/a)

Tetraploid Trumpets hybridized by R.J. Griesbach
× tetraploid Trumpets hybridized by R.J. Griesbach
Tepals pale golden yellow. Inside of tepals flushed peach-pink at margins, tips and midveins. Outside of tepals overlaid with shades of brownish and purplish pinks. Spots and papillae absent; filaments pale green; pollen dark russet-brown. Tepal margins slightly wavy, tips recurved. Stems 1.2–1.5m. Late. Tetraploid.
Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

Griesbach Tetra Pink Group VI(c/a)

Tetraploid Trumpets hybridized by R.J. Griesbach
× tetraploid Trumpets hybridized by R.J. Griesbach
Inside of tepals milk-white tinged rose-pink at margins and flushed pale creamy lemon in basal half; midveins soft rose-pink. Outside of tepals milk-white, flushed pale creamy lemon-yellow beside midribs, overlaid extensively with richer rose-pink colouring; midribs slightly duller, rose-pink. Spots absent; filaments pale green; pollen mahogany-brown. Tepal margins slightly wavy, tips recurved. Stems 1.2–1.5m. Late. Tetraploid.
Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

‘Gui Fei’ VII(b/b)

‘Bernini’ × ‘Siberia’

H: W. Dong, 2001 **G:** W. Dong, pre-2008 **N:** W. Dong, 2008 **I:** H.P. Zhang, 2006 **REG:** Wei Dong and Cheng du United Bio-technology Co. Ltd, 2008
Tepals, including throat, clear, slightly lilac, pink. Spots and papillae rich purplish pink-red, raised, quite large, fairly evenly distributed over basal half of each tepal, avoiding margins and midveins; nectaries green; pollen orange-brown; stigma apple-green. Fls 240mm wide, strongly scented; tepals 130 × 60mm, margins strongly ruffled, tips strongly recurved. Lvs 120 × 44mm, scattered. Stems 1.25m, green, with up to 7 fls in a racemose infl. June.
The epithet is a transcription of Mandarin Chinese meaning “dignity and beauty”.

‘Haley’ VII(a-b/b)

Parentage unknown

S: E.T. Kirsch, pre-1985 **I:** Melridge Inc., c.1988

Tepals strong purplish red (63A/B), with a strong red (53B) band 2–3mm wide straddling midveins, shading to strong purplish pink (63C) towards margins; margins white, 4–5mm wide at tips narrowing to 1mm at base; tepals becoming more brown with age. Papillae strong purplish red (63B), widely distributed over reddish part of tepals; nectaries green, with 1–2mm yellow margin; filaments 70–80mm, pale green to white; anthers 20mm, moderate red (179A/B); style pale green to white, sometimes tinged pink; stigma pale purple-grey. Fls 180–200mm wide, strongly scented; tepals ovate, inner 60mm, outer 30mm wide, margins slightly ruffled, tips slightly recurved; pedicels 70–80mm, dark green, ascending at 45° angle; infl. a compact raceme. Lvs 50–120 × 20–30mm, glossy, leathery, dark green. Stems 0.7–0.9m, with up to 8 fls. Mid-season.

‘Helen Skinner’ II(c/d)

H: F.L. Skinner, pre-1967

Inside of tepals dark purplish pink; midveins blackish purple in basal third. Spots and papillae absent; pollen dark orange. Fls dainty; borne on long pedicels. Stems 1.2m, greyish. Early.
Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

‘Helios’ I(a/b-c)

Seedling A-080-78 × seedling A-034-84

H: D.L. Egger, pre-1995 **I:** Cebeco Lilies, Inc., c.1998

Tepals vivid yellow (9A), outside of tepals shading to light greenish yellow (8B) beside midribs. Spots absent; papillae few, as roughened patches near nectaries; nectaries 22mm, strong yellow-green (144A); pollen strong orange (26A); stigma vivid yellow (9B) with faint greyish red (176A) spots. Fls 150mm wide, not scented; outer tepals 100 × 35mm, inner 100 × 48mm, ovate, waxy, with a satin sheen, tips slightly recurved; pedicels 60–140mm, strong yellow-green (144A), curving upwards. Lvs 115 × 18mm, leathery, thick, glossy, moderate olive-green (137A/147A). Stems 0.86m, strong yellow-green (144A); infl. an umbel-like raceme with up to 4 fls. or a branching raceme with 5 or more fls. Early.

‘Hitch’ VII(a/b)

Un-named Div. VII (Oriental hybrid) seedling
× un-named Div. VII seedling

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2008

Inside of inner tepals deep purplish pink (N66D); margins light purplish pink (65B) at base; midveins vivid purplish pink (N66B); throat yellowish white (155D). Inside of outer tepals deep purplish pink (N66C); margins and midveins vivid purplish pink (N66B), margins shading to deep purplish pink (N66D) at base. Outside of outer tepals dark purplish pink (186C), yellowish white (155D) beside midribs at base; margins deep purplish pink (N66C); midribs strong yellow-green (144C) at base and top [tip itself 144A], yellowish white (155D) between. Outside of inner tepals dark purplish pink (186C) with yellowish white (155D) veins, yellowish white (155D) beside midribs at base; midribs light yellow-green (144D) at base, shading to yellowish white (155D) tinged dark purplish pink (186C), yellowish white (155D) at top. Spots strong purplish red (60B); papillae yellowish white (155D) tinged strong purplish red (60B); nectaries strong yellow-green (144C); pollen strong orange (N25A); stigma greyish purplish red (N77D). Fls 220mm wide, scented; tepals 135 × 75mm, margins smooth, tips slightly recurved. Lvs 180 × 51mm. Stems 0.9m, green, with darker markings near leaf nodes, with up to 4 fls.

‘Honeymoon’ VII(a/b-c)

‘Sorbonne’ × ‘Simplon’

H: Q.F. Cheng, 2004 **G:** Q.F. Cheng, 2008

N: Q.F. Cheng, 2008

Inside of tepals palest flesh-pink, central portion of basal third honeyed golden yellow shading to closely

stippled colouring above, along and beside midveins, peachy pinkish orange in inner tepals and stronger reddish pink in outer tepals; throat green. Outside of tepals whitish; midribs purplish red. Spots few, dark pinkish red, sometimes raised, varying in size, scattered mostly towards edge of central yellow patch; papillae few, small, honeyed golden yellow, towards throat; nectary furrows apple-green; pollen dark russet-brown; stigma whitish, frosted. Fls 201mm wide, strongly scented; tepals 105 × 60mm, margins of outer tepals strongly ruffled, margins of inner tepals only slightly so, tips strongly recurved. Lvs 149 × 37mm, scattered, pale green. Stems 1.2m (under glass), green.

External web links: www.lanlilium.com (image)

It is not possible to register this cultivar epithet as it is in use already for another lily (ICNCP, 2004: Art. 27.1).

‘Hongyun’ VII(b/b)

‘Con Amore’ × ‘Acapulco’

H: Mrs Q. Lin, 2003 **G:** Mrs Q. Lin & Q. Zhou, 2006

N: Mrs Q. Lin & Q. Zhou, 2007 **REG:** Mrs Q. Lin & Q. Zhou, 2009

Tepals pale pink, inside of tepals with yellow flare at base of midvein; throat yellow and green. Spots sparse, conspicuous, slightly raised, pinkish red, evenly distributed either side of flare at base of each tepal; papillae white, beside nectary furrows; nectaries green; pollen orange-brown; stigma pale greyed purple. Fls 223mm wide, strongly scented; tepals 134 × 54mm, margins slightly ruffled, tips slightly recurved. Lvs 171 × 54mm, whorled. Stems 1.08m, green with purple spots, with up to 5 fls. Early (May). Diploid, through reduction from triploid. The epithet is a transcription of Mandarin Chinese meaning “great luck”.

‘Hyde Park’ VIII(a/b) LA

Seedling OR 02-11 × seedling BJ 496-06

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals strong orange (N25B), margins N25A; midveins of inner tepals N25A; throat pale greenish yellow (2D). Outside of outer tepals brilliant orange-yellow (23B), shading to vivid yellowish pink (28A) towards margins; midribs strong yellow-green (144B) at base, tinged dark red (59A) at top. Outside of inner tepals strong orange (25B), shading to N25B towards margins, closely spotted N25A; midribs strong yellow-green (144C) at base. Spots dark red (59A); papillae strong orange (N25A); nectaries dark bluish green (131A); pollen greyish red (178A); stigma strong reddish orange (169A). Fls 175mm wide; tepals 123 × 60mm, margins smooth, tips slightly recurved. Lvs 115 × 23mm. Stems 1.1–1.3m, green with darker markings, especially at base, with up to 4 fls. Lime-tolerant.

'Ice Bear' VII(a/b-c)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007**N:** Testcentrum voor Siergewassen BV, 2007**REG:** Testcentrum voor Siergewassen BV, 2009

Inside of tepals white (NN155C); outside of inner tepals predominantly NN155C. Spots absent; papillae conspicuous; nectaries green; pollen orange-brown; stigma dark purple. Tepals of medium length and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length and (very) broad, dark green. Stems of average height to tall, green with faint darker spots and stripes, with up to 8 fls. Mid-season.

'Ice Crystal' VIII(a/b) LA**'Esra' X seedling LA 93096-1 T**

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2008

Tepals, including throat, yellowish white (155D). Midribs of outer tepals strong yellow-green (143A) at base shading to light yellow-green (144D) at top, all flushed deep purplish pink (61D). Midribs of inner tepals strong yellow-green (144C) at base, shading to deep purplish pink (61D), upper half pale yellow (11D). Spots few, dark red (59A); papillae white; nectaries strong yellow-green (143A); pollen brownish orange (N167B); stigma light yellow-green (154D). Fls 215mm wide, slightly scented; tepals 125 × 60mm, margins smooth, tips slightly recurved. Lvs 160 × 24mm. Stems 1.2m, green, with up to 5 fls.

'Indian Summer' VIII(c/b-d) OT**H:** Mak Leek, Inc., pre-2008

Inside of tepals clear yellow; outside of tepals greenish yellow. Filaments and style pale lime-green, all very exserted; pollen dark russet-brown; stigma green.

Buds lime-green. Fls large; tepals slender, margins smooth. Stems 1.5m, dark. Late.

Published in *The Lily Nook cat.*, 2009

It is not possible to register this cultivar epithet as it, and Indian Summer Group, are in use already (ICNCP, 2004: Art. 27.1).

'Indiras' VIII(b/a) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007**N:** Testcentrum voor Siergewassen BV, 2007**REG:** Testcentrum voor Siergewassen BV, 2008

Inside of tepals strong reddish purple (70B) to moderate purplish red (186B); midveins greenish at base shading to purple-red. Outside of tepals moderate purplish red (70A) to strong reddish purple (70B). Spots and papillae absent; nectaries green; pollen orange-yellow; stigma pale grey. Tepals of medium length and width, margins slightly ruffled, tips slightly recurved. Lvs of medium length to long, and of medium width to broad. Stems of average height to tall, green, with few to average no. of fls. Late to very late.

'Intuition' VII(a/b)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2004

N: Marklily BV, 2004 **REG:** Testcentrum voor Siergewassen BV, 2007

Inside of tepals greenish white (155C) flushed strong purplish pink (73B); margins, tips and base paler; midveins white at base. Outside of inner tepals very pale purple (73D). Spots absent; papillae present; nectaries pale green; pollen reddish brown; stigma grey-green. Tepals of medium length and width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and broad to very broad. Stems of average height to tall, green with darker spots and stripes, with few to average no. of fls. Mid-season to late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)**'Invasion' VIII(a-b/b) OT**

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2008

Inside of inner tepals strong red (46A) to deep red (53A), becoming paler with age. Inside of outer tepals deep red (53A) to deep purplish red (187D); all margins light yellow (10C); midveins yellow-green at base. Outside of inner tepals light yellow (10C) to pale greenish yellow (10D), flushed purple-red at base. Spots few, darker than deep red (53A); papillae present; nectaries green; pollen red-brown; stigma dark purple. Fls 230mm wide; tepals of medium length and width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and of medium width to broad. Stems 1–1.25m, green with darker markings, with up to 7 fls. Late to very late.

'Ipenama' VII(a-b/c)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003 **REG:** World Breeding BV, 2008

Inside of tepals deep red (53A/60A), more blue with age; margins, tips and base a little paler. Outside of inner tepals strong purplish red (60C). Spots darker than deep red (53A/60A); papillae present; nectaries green shading to yellow; pollen red-brown; stigma grey-green spotted purple. Tepals of medium length to long and of medium width, margins ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and very broad. Stems 1.55m, green with some darker markings, with up to 4 fls. Mid-season.

'Jaunie Varti' II(c/d)

Un-named seedling X unknown

S: D. Hercbergs, c.2003 **G:** D. Hercbergs, 2003**N:** D. Hercbergs, 2004 **I:** D. Hercbergs, 2007**REG:** D. Hercbergs, 2008

Tepals light greenish yellow (4C); throat greenish; midribs strong purplish pink (68B). Spots few, small, red-purple; nectaries greenish; pollen greyed orange; stigma dark purple. Fls 70mm wide, slightly scented; tepals 38 × 14mm, margins smooth, tips strongly recurved. Lvs 200 × 50mm, whorled. Stems 1.4m,

green with darker markings, with up to 30 fls. Mid-season.

The name means "the new gates" in Latvian.

'Jerzy' VI(c/d)

'Rogalo' × unknown

H: R. Widerski, pre-2008 **N:** R. Widerski, 2008

I: R. Widerski, 2008 **REG:** R. Widerski, 2008
Inside of tepals golden yellow. Outside of tepals yellow; midribs pale green. Brushstrokes small, purple, fairly numerous, over one-third of throat; nectaries green; pollen orange; stigma green. Fls 150mm wide; tepals 130 × 40mm, margins slightly ruffled, tips recurved; pedicels 120mm, green; with secondary buds. Lvs 100 × 35mm, glossy. Stems 1.2m, green spotted brown-purple, with infl. of medium length. Mid-July.

'Jessie Metcalf' I(a/b)

'Merryspring' × 'Sutherland Peach Melody'

H: D.J. Hay, 1996 **G:** D.J. Hay, 2000 **N:** D.J. Hay, 2008
REG: D.J. Hay, 2008

Tepals cantaloupe-melon orange; throat white and green. Spots few, quite large, darker orange, scattered in an arc in central half of lowermost fifth of each tepal; nectary furrows white; pollen dark rust; stigma pink. Fls 160mm wide, not scented; tepals 78 × 38mm, margins smooth, tips straight. Lvs 68 × 25mm. Stems 0.82m, pale green, with up to 6 fls. Mid-season (July).

Named after an aunt of the registrant's husband.

'Jixiang' VII(b/b)

'Con Amore' × 'Acapulco'

H: S. Zheng, 2002 **G:** S. Zheng, 2006 **N:** S. Zheng, 2007
REG: S. Zheng, 2009

Tepals pinkish red; midribs greenish; throat yellow and green. Spots and papillae red, fairly large, numerous, over basal half of each tepal; nectaries green; pollen orange-brown; stigma milky white. Fls 230mm wide, strongly scented; tepals 145 × 65mm, margins slightly ruffled, tips slightly recurved. Lvs 136 × 25mm, whorled. Stems 1.12m, purple, with up to 6 fls. Early (May). Triploid.

The epithet is a transcription of Mandarin Chinese meaning "good luck".

'Johnathon Hay' I(b/c)

'Catherine Laburi' × 'Sutherland Fascination'

H: D.J. Hay, 2000 **G:** D.J. Hay, 2002 **N:** D.J. Hay, 2007
REG: D.J. Hay, 2008

Inside of tepals pale yellow fading to creamy white; throat white and green. Outside of tepals bronze-yellow. Spots few, small, round to elliptic, dark brown, in halo around throat; nectaries white; anthers very slender, pollen orange; stigma green flushed pink. Fls 155mm wide, not scented; tepals 72 × 40mm, margins slightly ruffled, tips slightly recurved. Lvs 56 × 18mm, scattered, glossy. Stems 0.72m, green, with up to 6 fls. Mid-season (July).

'Joyous Event' VII(a/b)

'Acapulco' × 'Miami'

H: Q.F. Cheng, 2004 **G:** Q.F. Cheng, 2008

N: Q.F. Cheng, 2008 **REG:** Yongkang Zhejiang Jiangnan Lily Breeding Co. Ltd, 2008

Inside of tepals purple-pink, breaking up to patchy white margins of irregular width and length, in upper half or less; tips and throat white. Outside of tepals pink; midribs green. Spots and papillae small, red-purple, only moderately numerous over basal half of each tepal, avoiding margins and midveins; nectaries pale green; pollen orange-brown; stigma mauve. Fls 170mm wide, strongly scented; tepals 85 × 45mm, margins slightly ruffled, tips slightly recurved. Lvs 95 × 30mm, scattered, glossy. Stems 0.4m (under glass), green, with up to 6 fls in a racemose infl. External web links: www.lanlilium.com (image)

'Kabana' VII(a-b/b-c)

Parentage unknown

H: Mak 't Zand BV, pre-2005 **N:** Mak 't Zand BV, 2005
REG: Mak 't Zand BV, 2008

Inside of tepals dark red (59A) to deep red (60A), more blue with age. Outside of inner tepals deep red (60A) to strong purplish red (60B). Spots darker red; papillae present; nectaries green shading to yellow; pollen orange-brown; stigma grey-green with small, pale purple spots. Tepals of medium length and width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and broad. Stems 1.25m, green with darker spots and stripes, with up to 5 fls. Mid-season.

'Kalahari' VIII(a/b) LA

Seedling AJ 95-32 × seedling BJ 96-24

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals strong orange (25A); midveins N25B; throat light orange-yellow (25D). Outside of outer tepals brilliant orange (25C), shading to strong orange (25A) towards margins; midribs light yellow (163D) at base, strong yellow-green (144A) at top, light orange-yellow (16B) between. Outside of inner tepals brilliant orange (25C); midribs brilliant greenish yellow (1B). Spots few, deep red (60A); papillae absent; nectaries vivid yellowish pink (28A); pollen strong reddish orange (169A); stigma moderate purplish red (58A). Fls 200mm wide; tepals 120 × 54mm, margins smooth, tips recurved. Lvs 160 × 18mm. Stems 1.4m, dark, with up to 9 fls.

‘Kampala’ VII(a/c-b)

Seedling RM 96-23 × seedling RM 96-24

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals deep purplish pink (N57C), shading towards margins through strong purplish red (63B) and strong purplish pink (63C) to pale purplish pink (65C); midveins vivid purplish red (N57B); throat pale yellow-green (4D). Inside of outer tepals deep purplish pink (N57C); margins yellowish white (155D); midveins vivid purplish red (N57A). Outside of outer tepals deep purplish pink (N57C) to dark purplish pink (186C); upper half of margins and base yellowish white (155D); midribs strong yellow-green (144B) at base, shading to light yellow-green (144D). Outside of inner tepals dark purplish pink (186C); tips and base yellowish white (155D); midribs strong yellow-green (144C) at base shading to strong greenish yellow (151A). Spots absent; papillae vivid purplish red (61C); nectaries strong yellow (N144B) with pale yellow-green (4D) margins; pollen strong orange (N163B); stigma light greenish grey (190C). Fls 215mm wide, scented; tepals 125 × 75mm, margins slightly ruffled, tips recurved. Lvs 200 × 65mm. Stems 1.15m, green with darker markings, with up to 5 fls.

‘Key Biscayne’ VII(a/b)

Seedling RM 96-17 × seedling RM 96-52

H: Vletter en Den Haan Beheer BV, pre-2006**N:** Vletter en Den Haan Beheer BV, 2006**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals deep purplish pink (N74C), shading to strong reddish purple (70B) towards margins; throat yellowish white (155D). Inside of outer tepals deep purplish pink (N74C); tips and beside nectaries yellowish white (155D). Outside of outer tepals deep purplish pink (N74C) to light purple (75B); base yellowish white (155D); midribs strong yellow-green (144B) at base and light olive (152B) at top, yellowish white (155D) between. Outside of inner tepals light purple (75A); base yellowish white (155D); midribs strong yellow-green (144C) at base. Spots and papillae deep purplish pink (N57C); nectaries strong yellow-green (144C); pollen deep orange-yellow (163A); stigma light grey (188D). Fls 230mm wide, scented; tepals 135 × 76mm, margins ruffled, tips straight. Lvs 185 × 75mm. Stems 1m, green with some darker markings, with up to 3 fls. Named after the most southerly barrier island off the Atlantic coast of Florida.

‘Kóbe’ I(a/b)

Parentage unknown

H: M. Tlach, pre-2007 **N:** M. Tlach, pre-2007**I:** M. Tlach, 2007 **REG:** M. Tlach, 2007

Inside of tepals pink, shading to yellowish centres; throat pinkish violet. Outside of tepals brown-pink; midribs green and pink. Spots absent; nectaries pale green, frosted pink; pollen yellow-orange; stigma pale pink. Fls 130mm wide; tepals 75 × 35mm, tips slightly recurved; pedicels 60mm, faintly spotted brown; infl. long, with secondary buds. Lvs 100 × 15mm, pale green. Stems 1m, with some faint brown spotting. Early July.

‘Kushi Maya’ VIII(c/b-c) OA*nepalense* × un-named, white-flowered, Div. VII (Oriental hybrid) seedling**S:** H.W. Hyde and Son, 2006 **G:** H.W. Hyde and Son, 2004 **N:** H.W. Hyde and Son, 2008 **I:** H.W. Hyde and Son, 2008 **REG:** H.W. Hyde and Son, 2008

Inside of tepals with broad, central band in upper half darker than pale yellow-green (149D), shading to white at margins; darker than deep purplish red (59B) at base and in throat, breaking into brushmarks at the upper margin; midveins deep purplish red (59B) in basal half, shading to brilliant yellow-green (149A). Outside of tepals deep purple (N81A) at base and beside and along basal half of midribs, darker than pale yellow-green (149D) at margins; midribs strong yellow-green (145A) in upper half. Spots absent; papillae few, only slightly raised, slightly darker than throat; nectaries greyish purple (N77A); filaments and connectives pale yellow-green; anthers dark red; pollen pale purplish pink (56); style mid-green; stigma dark greyish purple (N186B). Fls 270mm wide, strongly scented; tepals 200 × 70mm, margins slightly ruffled, tips recurved. Lvs 270 × 50mm, scattered. Stems 1.35m (under glass), with up to 6 fls; with underground stem-bulbils. Mid-season (July).

‘Labiano’ VII(a/b-c)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007**N:** Hobaho BV, 2007 **REG:** Testcentrum voor Siergewassen BV, 2009

Inside of tepals deep purplish pink (N66D) on a paler ground, becoming paler towards base; base and midveins brilliant yellow (12B). Outside of inner tepals predominantly very pale purple (73D). Spots fairly numerous to abundant, yellow-brown; papillae present; nectaries yellow-green; pollen orange-brown; stigma purple, speckled. Tepals of medium length to long, and broad, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of medium length and width, dark green. Stems tall, green with darker spots and stripes, with up to 6 fls. Mid-season to late.

‘Ladislao Sedlaro’ I(b-c/d)

‘Noris Star’ × ‘Apricot Supreme’

H: J. Dudek, pre-2007 **N:** J. Dudek, pre-2007**I:** J. Dudek, 2007 **REG:** J. Dudek, 2007

Inside of tepals orange-red; throat and outside of tepals paler; midribs orange-yellow. Spots numerous, large, purple, over half of each tepal; nectaries edged white; pollen brown-orange; stigma orange-brown. Fls 110mm wide; tepals 70 × 34mm, tips strongly recurved; pedicels 90mm, thick, green, downy; with secondary buds. Lvs 90 × 40mm, glossy, dark green. Stems 1.4m, green; infl. of average length. Early July.

‘Laimes Spēle’ I(a/b)

‘Saulkrasti’ × ‘Rotaļa’

H: Ā. Zorgevics, 1993 **G:** Ā. Zorgevics, 1999**N:** Ā. Zorgevics, 2001 **I:** University of Latvia Botanical Garden, 2003 **REG:** Ā. Zorgevics and University of Latvia Botanical Garden, 2009

Inside of tepals vivid yellow (14A); throat green. Outside of tepals brilliant yellow (13B). Spots few, quite large and conspicuous, round, vivid orange-

yellow (23A), just above throat on each tepal; nectaries clear; pollen brown; stigma greenish yellow. Fls 130mm wide, not scented; tepals 80 × 30mm, margins slightly ruffled, tips strongly recurved. Lvs 70 × 14mm, scattered, glossy. Stems 1.1m, brownish green, with up to 9 fls in a long infl. Early July. The name means “matter of chance” in Latvian.

‘Latvju Dancis’ I(a/b-c)

‘Dzintars’ × ‘Rotaļā’

H: Ā. Zorgevics, 1994 **G:** Ā. Zorgevics, 2000
N: Ā. Zorgevics, 2002 **I:** University of Latvia Botanical Garden, 2003 **REG:** Ā. Zorgevics and University of Latvia Botanical Garden, 2009
Inside of tepals vivid orange (28B); throat grey. Outside of tepals orange with a greenish patch. Spots absent; nectaries clear; pollen brown; stigma pale orange. Fls 170mm wide, not scented; tepals 90 × 30mm, margins smooth, tips strongly recurved. Lvs 110 × 10mm, scattered, glossy. Stems 1.1m, brownish green, with up to 13 fls. Early to mid-July. The name of a Latvian folk dance.

‘Lavender Ballerina’ I(a/-)

Parentage unknown

N: Cebeco Lilies, Inc., c.2008

Inside of tepals an unusual shade of reddish purple.

Stems 1.07m. Mid-season.

Published in *The Lily Nook cat.*, 2009

‘Le Baron’ VII(a/b-c)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2005 **N:** Van der Marel Lelie BV, 2005 **REG:** Van der Marel Lelie BV, 2007

Inside of tepals deep red (60A) to strong purplish red (60B), more blue with age; becoming paler towards margins; margins pale yellow-green (155A); midveins white at base. Outside of inner tepals strong purplish red (60C). Spots and papillae present; nectaries green; pollen orange-brown; stigma grey-green spotted purple. Tepals of short to medium length and of narrow to medium width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of short to medium length, and broad. Stems of average height to tall, green with darker markings, with few to average no. of fls. Late to very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

‘Lemon Supreme’ VI(b/a)

‘Lemon Delight’ × small, yellow-flowered Div. VI (Trumpet/Aurelian hybrid)

H: L.C. Bligh, 2002 **G:** L.C. Bligh, 2004

N: L.C. Bligh, 2008 **REG:** L.C. Bligh, 2008

Inside of tepals, including throat, brilliant greenish yellow (6C). Outside of tepals brilliant greenish yellow (6C) shading to pale yellow-green (4D); midribs tinged greyish red (182B). Spots absent; nectaries green; pollen brownish orange; stigma green. Fls 125mm wide, slightly scented; tepals 115 × 44mm, margins almost smooth, tips slightly recurved. Lvs 200 × 20mm, scattered. Stems 2.03m, green with maroon flecks, with up to 32 fls. Mid-January (Tasmania).

LEXUS VII(a/b-c)

Trade designation of ‘Zanlorexus’

‘Lileah’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2008

Tepals, including throat, pale yellow-green (4D); midribs vivid yellow (15B). Spots variable in size: outer large, round; inner smaller, sometimes paler and/or forming small, short dashes; grey-purple, over basal half of each tepal; papillae few, faint, grey-purple, between throat and spots; nectaries green, frosted; pollen golden orange; stigma pale green. Fls 85mm wide, not scented; tepals 65 × 26mm, margins smooth, tips slightly recurved. Lvs 110 × 6mm, scattered. Stems 0.8m, green with brown flecks, with up to 12 fls. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “fresh water”.

‘Ling Long’ VII(b/b)

‘Bergamo’ × ‘Siberia’

H: W. Dong, 2001 **G:** W. Dong, pre-2008 **N:** W. Dong, 2008 **I:** H.P. Zhang, 2006 **REG:** Wei Dong and Cheng du United Bio-technology Co. Ltd, 2008

Tepals palest flesh-pink, flushed darker soft rose-pink towards tips, between margins and midveins; margins whitish; midveins greenish yellow at base, tinged yellowish white above; throat buff. Spots and papillae few, slightly raised, varying in size, reddish orange, widely scattered over each tepal, but mostly in basal third; nectaries green; pollen orange-brown; style and stigma apple-green. Fls 250mm wide, slightly scented; tepals 140 × 60mm, margins slightly ruffled, tips slightly recurved. Lvs 130 × 36mm, scattered. Stems 1.2m, green, with up to 6 fls. June.

‘Little Pink Rascal’ VII(b/b)

Parentage unknown

S: E.T. Kirsch, 1975 **G:** E.T. Kirsch, 1975

I: Melridge Inc., c.1986

Tepals strong purplish pink (63C), shading through light purplish pink (62C) to pale purplish pink (62D) towards margins and tips; more brown with age. Papillae numerous in the centre of each tepal (c.125 on inner tepals, c.50 on outer); nectaries green; filaments 70–80mm, pale pink to white; anthers moderate reddish orange (178C), occasionally strong orange (168B) to moderate orange (168C); style pale green to white, sometimes tinged pink; stigma pale green. Buds strong purplish pink (63C) to light purplish pink (63D), with paler tips; midribs pale green, 1mm wide, pale pink on opening. Fls 200mm wide, strongly scented; tepals ovate, 30mm (inner) to 40mm (outer) wide, tips slightly recurved; pedicels to 70mm, dark green, ascending at 45° to stem. Lvs 50–100 × 20–30mm, leathery, glossy, dark green. Stems 0.5m, with up to 8 fls in a compact raceme 150mm long. Mid-season.

‘Littleton’ VII(a-b/c)

Parentage unknown

N: M.A. van den Berg en Zn, V.o.f., pre-2009

REG: M.A. van den Berg en Zn, V.o.f., 2009

Inside of tepals deep red (53A/60A), paler towards tips and shading to almost white towards margins; midveins yellow at base. Outside of inner tepals strong purplish red (64B) to deep purplish pink (64D). Spots and papillae present; nectaries green; pollen red-brown; stigma dark purple. Tepals short and of narrow to medium width, margins ruffled, tips slightly recurved. Lvs of medium length and broad to very broad. Stems 0.6m, green with darker markings, with up to 5 fls. Mid-season.

‘Louis XIV’

Unregistered seed parent of ‘**Antoni Widerski**’. No further details known.

‘Love Story’ VII(a-b/b)

Parentage unknown

H: Mak Breeding BV, pre-2003 **N:** Van der Marel Lelie BV, 2003 **REG:** Van der Marel Lelie BV, 2007

Inside of tepals dark red (59A), shading through strong purplish red (60C) to pale yellow-green (155A) towards margins; midveins yellow-green at base. Outside of tepals strong purplish red (64B). Spots and papillae present; nectaries yellow-green; pollen red-brown; stigma grey-green. Tepals of medium length to long and of medium width to broad, margins ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad to very broad. Stems of short to average height, green, with few to average no. of fls. Mid-season to late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

‘Lowana’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2007

Inside of tepals pale yellow (158B) ground tinged mauve, overlaid with light yellow (160B) in centre; throat pale yellow-green (4D). Outside of tepals pale yellow (158B); midribs marked greyish reddish orange (174B). Spots rather large, round to elliptic, mahogany-brown, well-distributed over lower two-thirds of each tepal; nectaries pale green, frosted; pollen golden brown; stigma pale green. Fls 75mm wide, not scented; tepals 65 × 28mm, margins smooth, tips slightly recurved. Lvs 105 × 5mm, scattered, dark green. Stems 1.1m, brown, flecked green, with up to 15 fls in pyramidal infl. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “woman” or “female”.

‘Luina’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2007

Inside of tepals moderate yellow (160A), shading to pale orange-yellow (159B) at tips; central band of moderate purplish pink (186D); throat moderate yellow (160A). Outside of tepals pale orange-yellow (159B) to dark reddish orange (173A). Small, V-shaped pencil-mark on each tepal and spots chocolate-brown; spots rather large, round, in moderate numbers, well-distributed over basal half of each tepal; nectaries pale green, frosted white; pollen golden brown; stigma pale green. Buds lightly downy. Fls 70mm wide, not scented; tepals 60 × 30mm, margins smooth, tips slightly recurved. Lvs 105 × 5mm, scattered, dark green. Stems 0.9m, brown, flecked green, with up to 15 fls in pyramidal infl. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for the blue wren (*Malurus cyaneus*).

‘Lullaby’ I(a/-)

Parentage unknown

N: Cebeco Lilies, Inc., c.2008

Inside of tepals soft baby-pink, central portion cream. Tepal margins very ruffled. Stems 0.9m. Early to mid-season.

Published in *The Lily Nook cat.*, 2009

‘Luz’ VII(a-b/b-c)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

N: Hobaho BV, 2007 **REG:** Testcentrum voor Siergewassen BV, 2009

Inside of tepals white (NN155B); outside of inner tepals predominantly NN155B. Spots absent; papillae present; nectaries green; pollen orange-yellow; stigma purple. Tepals of medium length and of medium width to broad, margins ruffled, tips slightly recurved. Lvs of medium length and broad, dark green. Stems 1m, green with faint darker spots and stripes, with up to 4 fls. Mid-season.

‘Macanta’ VII(a-b/c)

Div. VII (Oriental hybrid) seedling × ‘Gentile’

H: Testcentrum voor Siergewassen BV, pre-2007

N: Hobaho BV, 2007 **REG:** Testcentrum voor Siergewassen BV, 2009

Inside of tepals and outside of inner tepals greenish white (155C). Spots absent; papillae present; nectaries green; pollen reddish brown; stigma purple. Tepals of medium length and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and broad. Stems 1.25m, green with darker markings, with up to 4 fls. Mid-season to late.

‘Magic Lamp’ VI(b-c/b-c)

‘Henrieta’ × ‘Brno’

H: A. Mego, pre-2008 **N:** A. Mego, 2008 **I:** A. Mego, 2008 **REG:** A. Mego, 2008

Tepals cream-white, inside of tepals with orange rays in basal half; midribs pale green. Spots few, small, brown, over basal two-fifths of each tepal; papillae long, dark apricot; nectaries dark green, edged green; pollen chocolate-brown; stigma pale green. Fls

170mm wide; tepals 90 × 35mm, margins slightly ruffled, tips very slightly recurved; pedicels 80mm, green spotted brown; with secondary buds. Lvs 210 × 30mm, pale green. Stems 1.35m, greyish with brownish green spots, with infl. of average length. Early July.

‘Mango’ I(a/-)

Parentage unknown

N: “The Originals” U.A., Coöp. Kwekersvereniging, c.2005

Inside of tepals mango-orange. Spots dark brown, coalescing in central portion of each tepal and spreading over more than half of each tepal. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009

‘Marianne Timmer’ I(a-b/b)

Parentage unknown

H: Mak Breeding BV, pre-2006 **N:** Mak Breeding BV, 2006 **REG:** Mak Breeding Rights BV, 2008

Inside of tepals strong red (46A). Outside of inner tepals strong red (46A) to moderate red (179A). Spots and papillae present; nectaries dark red; pollen red-brown; stigma orange. Tepals short and of narrow to medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length and width. Stems 1–1.1m, green with some darker markings, with up to 11 fls. Mid-season.

Named after the Olympic gold-medal-winning Dutch speed-skater.

‘Marrawah’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2008

Tepals, including throat, pale orange-yellow (23D); midribs tinged red-brown. Spots of medium size, round to elliptic, dark brown, on basal half of each tepal; coalescing to form a thin V-shape either side of midvein beyond nectary furrow; nectaries green frosted; pollen bright orange; style and stigma soft pale orange. Fls 85mm wide, not scented; tepals 75 × 37mm, margins smooth, tips slightly recurved. Lvs 85 × 5mm, scattered. Stems 0.7m, brown, flecked green, with up to 10 fls. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “one”.

‘Marvin Joslin’ I(c/c-d)

Parentage unknown

H: M. Joslin, pre-2007 **G:** M. Joslin, pre-2007

N: K. Frey, 2007 **I:** Estate Perennials, c.2007

REG: K. Frey, 2007

Inside of tepals vivid yellow (9A); throat brilliant yellow (9C), slightly frosted white. Outside of tepals slightly paler than brilliant yellow (9C); midribs vivid yellow (9A). Heavily marked inside with greyish purple (N77) spots on basal half, extending further along margins, occasionally to tips; nectaries vivid yellow (9A); pollen strong orange (N25C); stigma pale greenish yellow (9D). Fls 102mm wide, not scented; tepals 94 × 28mm, margins smooth, tips

slightly recurved; with secondary buds. Lvs 104 × 13mm, scattered. Stems 1.125m, dark green flushed brown, with up to 20 fls. Mid-season.

Named after the former proprietor of Estate Perennials who bred the lily.

Nomenclatural standard: colour prints supplied by registrant (WSY 0112960)

‘Matignon’ VII(a/b)

Seedling RM 96-26 × seedling RM 98-18

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals deep purplish pink (N57D), shading to strong purplish red (64C) towards margins; patch above nectaries light yellow-green (2C) tinged vivid reddish orange (33B); throat yellowish white (155D). Inside of outer tepals deep purplish pink (N57D), yellowish white (155D) at base beside nectaries shading to light yellow-green (2C) above nectaries; midveins vivid reddish orange (33B) in basal half. Outside of outer tepals deep purplish pink (N66D); tips vivid purplish red (67B), base yellowish white (155D); midribs strong yellow-green (144C) at base and top, vivid purplish red (67B) between. Outside of inner tepals moderate purplish pink (65A), shading to strong purplish pink (68B) towards margins; base tinged yellowish white (155D); midribs light yellow-green (144D) at base, shading to yellowish white (155D) and strong purplish pink (68B). Spots absent; papillae vivid purplish red (N57A); nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma pale yellow-green (193D). Fls 170mm wide, scented; tepals 125 × 70mm, margins slightly ruffled, tips slightly recurved. Lvs 190 × 48mm. Stems 1.2m, green with darker markings, with up to 6 fls.

‘Matrix Orange’ I(a/b)

Syn. of ‘Orange Matrix’

‘Matrone’ VII(a/c-b)

Parentage unknown

H: Van der Marel Lelie BV, pre-2007 **N:** Van der Marel Lelie BV, 2007 **REG:** Van der Marel Lelie BV, 2009

Inside of tepals, and principal colour of outside of inner tepals, white (NN155C). Spots absent; papillae present; nectaries green; pollen orange-brown; stigma dark purple. Fls 150–200mm wide; tepals long and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and broad. Stems of average height to tall, green with some darker markings, with up to 7 fls. Mid-season.

MAURENA VII(a-b/b)

Trade designation of ‘Zanlorena’

'Maverick' VII(a-b/c-d)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006**N:** Testcentrum voor Siergewassen BV, 2006**REG:** Testcentrum voor Siergewassen BV, 2008

Inside of tepals slightly more red than strong purplish red (67A) to vivid purplish red (67B), paler towards tips and base; margins shading to nearly white; midveins deep red (53A) at base. Outside of tepals deep purplish pink (N66C/D). Spots present; papillae faint; nectaries green, fading to yellow; pollen orange-brown; stigma purple. Tepals of medium length to long and of medium width to broad, margins ruffled, tips slightly recurved to recurved. Lvs of medium length and broad. Stems 1.25m, green with darker markings, with up to 4 fls. Mid-season to late.

'Mawbanna' I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005**G:** N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2008

Inside of tepals, including throat, vivid yellowish pink (28A), shading to vivid reddish orange (33A) at tips. Outside of tepals strong yellowish pink (34D) to vivid reddish orange (44C). Small, V-shaped pencil-mark on each tepal and spots mahogany brown; spots fairly large, round, well-distributed over basal half of each tepal; nectaries vivid reddish orange (34A); pollen terracotta; stigma pale orange. Fls 70mm wide, not scented; tepals 60 × 23mm, margins smooth, tips slightly recurved. Lvs 100 × 5mm, scattered, dark green. Stems 0.9m, green, flecked brown, with up to 15 fls in pyramidal infl. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for "black".

'Maxima'**I:** Mak-Leek Inc., c.2007

No further details of this lily are known beyond that it was granted EU PBR 19991 in Aug 2007, terminated in Aug 2009.

'Mejdan' VI(b-c/b)(L. × *kewense* 'White Henry' × 'Kankán')

X unknown

H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007**I:** B. Mičulka, 2007 **REG:** B. Mičulka, 2007

Inside of tepals creamy white; throat grooved, bright yellow. Outside of tepals creamy white flushed yellow; midribs pink. Nectaries deep yellow-green; pollen ochre-brown, poorly developed; stigma green. Fls 175mm wide; tepals 120 × 42mm, margins strongly ruffled, tips recurved; pedicels 65mm, with numerous dark brown spots. Lvs 120 × 35mm, pale green, recurved. Stems 1.15m, with numerous dark brown spots; infl. short. Late June.

'Melissimo' VII(b-a/b-c)**I:** Harts Nursery, 2009

Inside of tepals palest pinkish white, sometimes pale greenish cream along and beside midveins; a narrow border of closely stippled rich rose-pink at margins, edged by a wire rim of dark rose-pink. Spots absent;

papillae few, small, greenish white; nectary furrows and style dark apple-green; filaments pale apple-green; pollen reddish orange; stigma dark brown. Outer tepals narrowly ovate; all tepal margins closely ruffled, tips slightly recurved.

Exhibited by Harts Nursery at RHS Chelsea Flower Show, 2009.

'Mella' I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005**G:** N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2008

Inside of tepals light purplish pink (73C), shading to pale orange-yellow (159C) towards and in throat.

Outside of tepals light purplish pink (73C) shading to light pink (49C). Spots of medium size, mahogany-brown, scattered fairly evenly over basal half of each tepal; nectaries green, frosted; pollen bright orange; stigma pale pink. Fls 95mm wide, not scented; tepals 65 × 23mm, margins smooth, tips slightly recurved. Lvs 100 × 4mm, scattered. Stems 0.8m, brown, flecked green, with up to 12 fls. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for "to run".

'Memphis Bell' VII(a/b-c)

Parentage unknown

H: Mak Leek, Inc. (U.S.A.), pre-2008 **N:** Ruiter Bloembollen BV, 2008 **REG:** Ruiter Bloembollen BV, 2008

Inside of tepals light purplish pink (73C); margins of outer tepals yellowish white (155D); midveins vivid purplish red (N57B); throat yellowish white (155D). Outside of outer tepals moderate purplish pink (186D); margins and beside base of midribs yellowish white (155D); midribs strong yellow-green (144C) at base, light olive (152A) at top, vivid purplish red (61C) between. Outside of inner tepals moderate purplish pink (186D), shading to dark purplish pink (186C) beside midribs; midribs light yellow-green (144D) at base shading to strong purplish red (63B). Spots deep purplish red (59B); papillae yellowish white (155D) tinged deep purplish red (59B); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma pale yellow-green (193D) with deep purplish red (71A). Fls 275mm wide, scented; tepals 155 × 60mm, margins slightly ruffled, tips recurved. Lvs 195 × 43mm. Stems 1m, green, with darker markings at base, with up to 6 fls.

'Meng Fan' VII(a/b)

'Rosy Dream' × 'Titanico'

H: W. Dong, 2001 **G:** W. Dong, pre-2008 **N:** W. Dong, 2008 **I:** H.P. Zhang, 2008 **REG:** Wei Dong and Cheng du United Bio-technology Co. Ltd, 2008

Tepals, including throat, deep rich pink, shading to narrow, whitish margins. Spots moderately large, fairly numerous, purple, in basal third to half of each tepal; nectaries green, edged yellow; pollen orange-brown; stigma purple. Fls 190mm wide, slightly scented; tepals 130 × 56mm, margins slightly ruffled, tips slightly recurved. Lvs 110 × 27mm, scattered. Stems 1.25m, green, with up to 6 fls. June.

'Mengha' I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2008

Tepals vivid yellow (9A); midribs tinged red; throat pale greenish yellow (9D). Spots of medium size, round to elliptic, mahogany-brown, scattered fairly evenly over basal half of each tepal; nectaries lemon-yellow, frosted; pollen pale orange; stigma pale lemon-yellow. Fls 85mm wide, not scented; tepals 55 × 23mm, margins smooth, tips slightly recurved. Lvs 100 × 4mm, scattered. Stems 0.75m, brown, with up to 18 fls. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for "toe".

MERIVA VII(a/b-c)

Trade designation of 'Zanlorva'

'Merluza' VIII(a/b-c) LA

Seedling BW 96-29 × seedling AW 99-10

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Tepals yellowish white (155D); basal half of midveins and throat light yellow-green (150D and 145C respectively). Outside of outer tepals flushed light yellow-green (145D); midribs 145C at base shading to strong yellow-green (144C). Outside of inner tepals light yellow-green (150D) at base; midribs 145C at base flushed moderate red (185B). Spots and papillae absent; nectaries strong yellow-green (144A); pollen brownish orange (N170A); stigma very pale purple (73D). Fls 205mm wide; tepals 130 × 66mm, margins slightly ruffled, tips slightly recurved. Lvs 165 × 25mm. Stems 1.3m, green, with up to 4 fls. The name is the Spanish word for "hake".

'Mestre' VIII(a/b) LA

Seedling BW 98-09 × seedling RS 91-09

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals dark purplish pink (186C), shading to strong purplish pink (68B) towards margins; patch above nectaries yellowish white (155D); throat brilliant yellow-green (142B). Inside of outer tepals strong purplish pink (68B), shading to deep purplish pink (68A) towards margins; patch above nectaries yellowish white (155D). Outside of tepals dark purplish pink (186C), outer tepals shading to deep purplish pink (68A) towards margins; midribs strong yellow-green (144B) at base, those of outer tepals shading to moderate olive-brown (199A) at top. Spots few, dark red (59A); papillae absent; nectaries strong yellow-green (143B); pollen dark red (187A); stigma deep purplish pink (N57D). Fls 162mm wide, slightly scented; tepals 106 × 54mm, margins slightly ruffled towards tip, tips recurved. Lvs 150 × 23mm. Stems 1.25m, green, with up to 7 fls.

Named after a town in Veneto, northern Italy.

'Metropol' I(a-b/b-c)

Parentage unknown

H: B. Mičulka, pre-2007 **N:** B. Mičulka, pre-2007

I: B. Mičulka, 2007 **REG:** B. Mičulka, 2008

Tepals pale yellow; throat paler; midribs brown. Spots absent; nectaries frosted white; pollen brown; stigma brown. Fls 170mm wide; tepals 105 × 47mm, margins of outer tepals slightly ruffled, tips slightly recurved; pedicels 55mm, with faint brown spots. Lvs 115 × 16mm, dark green. Stems 0.65m, with faint brown-red spots; infl. short. Late June.

'Millesimo' VII(a/b)

Seedling RM 97-04 × seedling RM 99-16

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Tepals, including throat, yellowish white (155D). Inside of inner tepals flushed very pale purple (73D) with a patch of light purplish pink (73C) beside nectaries; all margins deep purplish pink (73A; margins of outside of inner tepals N74C). Base of outside of outer tepals with a patch of moderate red (180A) beside midribs; midribs strong yellow-green at base and top (144B and 144B tinged 144A respectively) and light yellow-green (144D) between. Midribs of inner tepals strong yellow-green (144B at base, 144C at top). Spots and papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen dark reddish orange (173A); stigma dark purple (79B). Fls 260mm wide, scented; tepals 155 × 95mm, margins slightly ruffled, tips slightly recurved. Lvs 225 × 50mm. Stems 1.3m, green, with darker markings near leaf nodes, with up to 4 fls.

'Miramar' I(a-b/b-c)

Parentage unknown

H: Mak Breeding BV, pre-2006 **N:** Mak Breeding BV, 2006 **REG:** Mak Breeding Rights BV, 2008

Tepals moderate purplish pink (65A) to light purplish pink (65B), paler with age; midveins orange-pink at base. Spots present, coalescing along margins towards base; papillae present; nectaries green, becoming pink; pollen orange-brown; stigma dark purple. Tepals (very) short and of narrow to medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length and of narrow to medium width. Stems 1.05m, green with faint darker spots and stripes, with up to 5 fls. Mid-season to late.

'Miss Feyta' VIII(b/c) OT

Parentage unknown

H: R.J. Griesbach, pre-2005 **N:** Lily Company BV, 2005 **REG:** Lily Company BV, 2008

Inside of tepals dark red (187C), shading to deep purplish red (187D) towards margins; margins greenish white (155C); tips strong purplish red (60C); throat deep purplish red (187D). Outside of outer tepals deep purplish red (61A) to strong purplish red (61B); outside of inner tepals 61B; all margins light purplish pink (62C) to pale purplish pink (62D); all tips vivid purplish red (61C); midribs strong yellow-green (144C), 144A at top of outer tepals. Spots dark red (187A); papillae 187C; nectaries dark

greyish reddish brown (200A); pollen greyish reddish brown (200B); stigma dark red (187A). Fls 240mm wide; tepals 120 × 54mm, margins ruffled, tips recurved. Lvs 195 × 67mm, dark green. Stems 0.9m, green with some darker markings, with up to 3 fls. Named after the grand-daughter, Feya Tjittes, of P. Groot of Lily Company BV.

‘Modigliani’ VII(a/b)

Seedling RM 98-08 × seedling RM 96-24

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals light purplish pink (inner 65B, outer 68D); midveins vivid yellow (13A) above nectaries, shading to moderate purplish pink (65A); throat yellowish white (155D). Outside of outer tepals moderate purplish pink (65A) in upper half, shading through very pale purple (69C) to yellowish white (155D) at base; midribs strong yellow-green (base 144C, top 143A), light yellow-green (145D) between. Outside of inner tepals light purplish pink (65B), yellowish white (155D) at base; midribs strong yellow-green (144C), flushed light purplish pink (65B) at top, yellowish white (155D) between. Spots few, deep purplish pink (64D); papillae light greenish yellow (4B); nectaries strong yellow-green (144C); pollen dark reddish orange (175B); stigma dark purplish red (N79A). Fls 210mm wide, scented; tepals 130 × 78mm, margins ruffled, tips recurved. Lvs 225 × 55mm. Stems 1.05–1.5m, green, with up to 4 fls. Named after the Italian painter and sculptor who lived 1884–1920.

‘Monrovia’ VII(a/b)

Seedling 93-14 × seedling RW 99-21

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals white than white (NN155D); throat white (NN155D). Outside of tepals white (NN155D), outer tepals tinged strong yellow-green (144A) at tips; midribs strong yellow-green (144C) at base, shading to pale yellow-green (155A). Spots and papillae white (NN155D); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma pale yellow-green (193D). Fls 250mm wide, scented; tepals 150 × 80mm, margins slightly ruffled, tips recurved. Lvs 210 × 75mm. Stems 1.2m, green, with up to 3 fls.

‘Montebello’ I(a/b)

Parentage unknown

H: World Breeding BV, pre-2007 **N:** World Breeding BV, 2007 **REG:** World Breeding BV, 2007

Inside of tepals vivid yellowish pink (28A) to vivid yellowish orange (30C); midveins strong orange (24A) at base. Outside of tepals vivid yellowish orange (30C). Spots absent; papillae faint; nectaries orange; pollen red-brown; stigma purple. Tepals of short to medium length and of medium width, margins (very) slightly ruffled, tips slightly recurved to recurved. Lvs of medium length, and of narrow to medium width. Stems 1.45m, green with some darker

markings, with up to 7 fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

‘Montefalcone’ VIII(a/a) LA

Seedling BW 98-08 × seedling AG 98-21

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals vivid yellow (12A); throat light greenish yellow (3C). Outside of outer tepals brilliant greenish yellow (6C); margins brilliant yellow (13B); midribs strong yellow-green (base 143B, top 144C). Outside of inner tepals vivid yellow (9A); margins 14B; midribs strong yellow-green (144C) at base. Spots and papillae absent; nectaries strong yellow-green (144A); pollen strong reddish orange (169A); stigma brilliant yellow-green (154B). Fls 160mm wide, scented; tepals 140 × 64mm, margins smooth, tips slightly recurved. Lvs 155 × 35mm. Stems 1.3m, green with darker markings, with up to 3 fls.

‘Montego Bay’ VIII(a/b) OT

H: Liliande BV, pre-2008

Inside of tepals dominated by a large, rounded zone of rich reddish pink in basal half, of more intense colour towards midveins; feathering higher along midveins and towards slightly bronzed, soft apricot-yellow margins. Nectary furrows rich apple-green with a wide margin of greenish yellow at base of tepals. Inside of upper half of tepals bronzed soft apricot-yellow; towards tips overlaid peachy pink. Spots dull, dark red, quite large, evenly distributed in two sprays in rich reddish pink zone in basal half of each tepal, away from margins and midveins. Papillae absent. Fls very large, of heavy substance; tepal margins smooth, tips strongly recurved. Stems 1.5m. Late. Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

‘Monteleone’ VIII(a/b)

Seedling BW 95-07 × seedling AJ 96-27

H: Vletter en Den Haan Beheer BV, pre-2004

N: Vletter en Den Haan Beheer BV, 2004

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals brilliant yellow (20A), shading to brilliant orange (25C) towards margins; throat pale yellow (20D). Inside of outer tepals light orange-yellow (22B), shading to strong orange (25B) towards margins. Outside of outer tepals light yellow-green (145D), shading to brilliant orange (25C) towards margins; midribs light yellow-green (145B) at base shading to strong yellow-green (144B) at top. Outside of inner tepals light yellow (20B), shading to strong orange (25B) towards margins; midribs light yellow-green (145C). Spots deep red (60A); papillae absent; nectaries strong yellow-green (144B); pollen strong brown (172A); stigma pale yellow (20D) with deep red (60A) stripes. Fls 230mm wide; tepals 130 × 57mm, margins smooth, tips recurved. Lvs 170 × 37mm. Stems 1.35m, green with darker markings, with up to 9 fls.

'Morandi' VIII(a/b) LA

Seedling BG 97-14 X seedling AG 98-12

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals light greenish yellow (8B); midveins brilliant yellow (8A); throat pale greenish yellow (1D). Inside of outer tepals brilliant yellow (8A). Outside of outer tepals light greenish yellow (3D); margins 5C; midribs light yellow-green (145B) at base, strong yellow-green (143B) at top. Outside of inner tepals light greenish yellow (5C); midribs strong yellow-green (143B) at base. Spots deep red (60A); papillae absent; nectaries strong yellow-green (145A); pollen moderate brown (165A); stigma moderate red (183D). Fls 200mm wide; tepals 140 × 57mm, margins smooth, tips slightly recurved. Lvs 155 × 27mm. Stems 1.3m, dark with green markings, with up to 4 fls. Named after the Italian painter who lived 1890–1964.

'Morgan' I(a/b-c)

Un-named, yellow-flowered seedling X un-named, orange-flowered seedling

H: T. Zimmerman, 1999 **G:** T. Zimmerman, 2002**N:** T. Zimmerman, 2005 **REG:** T. Zimmerman, 2008

Inside of tepals brilliant orange-yellow (23B) in centre, shading to strong orange (25A) towards margins; throat orange. Outside of tepals orange. Spots absent; nectaries orange; filaments orange, pollen very dark brown; stigma yellowish grey. Fls 130mm wide, not scented; tepals 75 × 30mm, margins smooth, tips straight. Lvs 90 × 20mm, scattered. Stems 1m, dark green, with up to 6 fls. Early to mid-July.

'Moselle' VIII(a/c-b) LA

Seedling AW 95-07 X seedling BM 496-03

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals dark purplish pink (186C); patch above nectaries yellowish white (155D) with dark red (183A) stripe; throat yellowish white (155D). Outside of outer tepals moderate purplish pink (70D), shading to strong purplish red (71D) towards margins; midribs strong yellow-green (143B) at base and top, moderate purplish red (184C) between. Outside of inner tepals deep purplish pink (70C), shading to 72D towards margins; midribs strong yellow-green (143C) at base. Spots and papillae absent; nectaries strong yellow-green (143A); pollen brownish orange (N167B); stigma light yellow-green (145C). Fls 220mm wide, slightly scented; tepals 125 × 61mm, margins smooth, tips slightly recurved. Lvs 150 × 25mm. Stems 1.1m, green, with up to 5 fls.

'Music Art' I(a-b/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2006 **N:** W. Vink, 2006 **REG:** Mak Breeding Rights BV, 2008

Tepals slightly more orange than vivid yellow (12A); midveins yellow-green at base with purple-red spots. Spots dark red (187B) or darker, along margins and towards base inside, also on outside of tepals; papillae present; nectaries yellow-green; pollen orange-brown; stigma dark purple. Fls 300mm wide; tepals short and of

narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of medium length to long, and of narrow to medium width. Stems 1.1–1.25m, green with darker spots and stripes, with up to 9 fls. Mid-season. Published in *The Lily Nook cat.*, 2009

'Mustang' VII (a/b)

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2003**N:** Vletter en Den Haan Beheer BV, 2003**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals deep purplish pink (N57C), shading to deep purplish pink (N66C) towards margins; midveins strong purplish red (61B); throat yellowish white (155D). Outside of outer tepals deep purplish pink (67C), shading to strong purplish red (67A) towards margins, yellowish white (155D) beside midribs at base; midribs strong yellow-green (base 144C; top 144B with a small, yellowish white (155D) patch), yellowish white (155D) between. Outside of inner tepals moderate purplish red (186B), shading to deep purplish pink (67C) towards margins, yellowish white (155D) beside midribs at base; midribs strong yellow-green (144C) at base shading to light olive (152B). Spots strong purplish red (60B); papillae yellowish white (155D) with strong purplish red (61B); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma light yellow-green (145D). Fls 230mm wide, scented; tepals 135 × 80mm, margins ruffled, tips recurved. Lvs 200 × 45mm. Stems 1.3m, green, with up to 6 fls.

'Nabageena' I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005**G:** N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2008

Inside of tepals moderate purplish red (184C), basal half flushed pale yellow (158A); throat pale yellow (158A). Outside of tepals moderate purplish red (184C) shading to dark yellowish pink (181D). Spots fairly large, mostly round, mahogany-brown, scattered fairly evenly over basal two-thirds of each tepal; nectaries pink, frosted; pollen bright orange; stigma pale orange. Fls 80mm wide, not scented; tepals 65 × 22mm, margins smooth, tips recurved. Lvs 65 × 3mm, scattered. Stems 0.75m, brown, flecked green, with up to 15 fls. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for "sun".

'Naira' I(a/b)

'Holebaba' X 'Latvia'

H: M. Tlach, pre-2007 **N:** M. Tlach, pre-2007**I:** M. Tlach, 2007 **REG:** M. Tlach, 2007

Inside of tepals violet-pink, shading to white towards margins; throat greenish. Outside of tepals yellowish. Spots numerous, pink-violet; nectaries pale green, frosted white; pollen orange-brown; stigma pinkish brown. Fls 140mm wide; tepals 80 × 40mm, inner tips recurved, outer slightly recurved; pedicels 50mm, stout, green. Lvs 90 × 13mm, mid-green. Stems 0.8m, stout, green, with few, fairly large, brown bulbils; infl. short. Late June.

‘Namara’ VII(a/b)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2008

N: Testcentrum voor Siergewassen BV, pre-2008

REG: Testcentrum voor Siergewassen BV, 2008

Inside of tepals flushed strong purplish pink (73B) on a pale ground, white towards base; becoming much paler with age. Outside of inner tepals light purplish pink (62C). Spots absent; papillae present; nectaries green; pollen orange-brown; stigma grey-green.

Tepals of medium length to long and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved. Lvs of medium length and of medium width to broad. Stems of average height to tall, green with faint darker spots and stripes, with few to average no. of fls. Mid-season to late.

‘Northern Dazzle’ VIII(c/b) OT

H: R.J. Griesbach, pre-2008

Inside of tepals matt crimson tinged russet in basal two-thirds; colour extending to margins in basal third only; upper edge rounded, feathering to soft apricot-yellow in apical third and margins in upper two-thirds. Outside of tepals whitish, base dark violet overlaid with a little green; shading gradually becoming more green, speckled pale violet, beside green midribs. Spots and papillae absent; filaments and style apple-green; pollen rich reddish orange. Tepal margins wavy, tips strongly recurved. Fls large. Stems 1.5m. Late. Tetraploid.

Published in *The Lily Nook cat.*, 2009

‘Northern Delight’ VIII(c/b) OT

H: R.J. Griesbach, pre-2008

Inside of tepals soft melon-orange, shading to paler apricot-yellow at tips; base of midveins apple-green. Outside of tepals white with a stripe of rosy purple beside midribs; midribs sometimes greenish towards base. Spots few, reddish orange but faint, round or coalescing into short dashes, sprinkled over basal half of each tepal; papillae absent; filaments apple-green; pollen dark russet brown. Tepal margins smooth, tips recurved; pedicels green. Stems 1.5m, sturdy. Late. Tetraploid.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Northern Star’ VIII(c/b) OT

H: L.M. Collicutt, pre-2008

Inside of tepals lavender-pink, fading towards tips and margins in apical third; shading to darker fuchsia-pink towards base and along and beside midveins; nectary furrows strong apple-green. Outside of tepals fuchsia-pink overlaid dark violet, shading paler towards margins. Spots few, greyed violet, quite large, round to elliptic, scattered mostly on basal third of each inner tepal; papillae absent; filaments and style pale yellowish green; pollen russet-orange. Buds dark maroon. Tepal margins smooth, tips strongly recurved. Stems 1.2m. Late.

Published in *The Lily Nook cat.*, 2009

‘Nuance’ VIII(b-a/b) LO

Div. VII (Oriental hybrid) seedling × seedling LO 93003 7B T

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2008

Tepals, including throat, yellowish white (155D). Strong purplish red (58B) stripes and patches beside and along midveins. Outside of outer tepals shading to strong purplish red (58B) flushed light yellow-green (145C) beside midribs; midribs strong yellow-green (144B) at base. Outside of inner tepals flushed strong purplish red (58B) at tips; base light yellow-green (144D); midribs at base strong yellow-green (144C) tinged strong purplish red (58B). Spots absent; papillae strong purplish red (58B); nectaries strong yellow-green (144C) edged light greenish yellow (3D); pollen strong orange (25A); stigma light greenish grey (190D). Fls 215mm wide, scented; tepals 120 × 56mm, margins slightly ruffled, tips recurved. Lvs 220 × 45mm. Stems 1.1m, green, with up to 4 fls.

‘Nutcracker’ I(b-c/-)

‘Noris Star’ × ‘Tetra Nutmegger’

H: A. Evans, pre-2008

Tepals vibrant orange-red, heavily spotted. Stems 1.2m. Mid-season. A robust tetraploid.

Published in *The Lily Nook cat.*, 2009

‘Olina’ I(a-b/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** “The Originals” U.A., Coöp. Kwekersvereniging, 2003

REG: Mak Breeding Rights BV, 2008

Inside of tepals strong red (46A) to moderate red (47A), shading to dark red (187A) at base and in places along margins. Outside of tepals strong reddish orange (42B/C) spotted darker. Spots darker than dark red (187A); papillae present; nectaries purple-red; pollen red-brown; stigma purple. Fls 160mm wide; tepals short and narrow, margins (very) slightly ruffled, tips slightly recurved. Lvs of short to medium length, and of narrow to medium width. Stems 0.8–1.3m, green with darker spots and stripes, with few to average no. of fls. Mid-season.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Omaggio’ VII(b-a/b)

Seedling RW 94-09 × seedling 00-310

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Tepals, including throat, greenish white (155C). Spots absent; papillae greenish white (155C) in basal third of each tepal; nectaries strong yellow (N144B); pollen moderate reddish brown (175A); stigma greenish grey (188A). Fls 160mm wide; tepals 110 × 65mm, margins slightly ruffled, tips slightly recurved. Lvs 200mm long, mid-green. Stems 1.2m, pale green, with up to 4 fls.

‘Orange Art’ I(a/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2005 **N:** “The Originals” U.A., Coöp. Kwekersvereniging, 2005

REG: Mak Breeding Rights BV, 2008

Inside of tepals vivid yellowish pink (28A) to vivid orange (28B), dark red (187A/B) at base and in places along margins; base of midveins orange with purple-red spots. Outside of tepals vivid yellowish orange (30C) to strong orange (30D) with dark spots. Spots darker than dark red (187A/B); papillae present; nectaries orange; pollen red-brown; stigma purple. Fls 150mm wide; tepals short and narrow, margins slightly ruffled, tips slightly recurved. Lvs of medium length, and of narrow to medium width. Stems 0.8–1.35m, green with darker spots and stripes, with few to average no. of fls. Early to mid-season. Published in *The Lily Nook cat.*, 2009

‘Orange Baby’ I(a-b/b)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003

REG: World Breeding BV, 2008

Inside of tepals vivid yellowish pink (28A) to vivid orange (28B). Outside of inner tepals strong orange (30D). Spots few; papillae present; nectaries orange; pollen red-brown; stigma purple on orange ground. Tepals (very) short and narrow, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of short to medium length, and of narrow to medium width. Stems 0.35–0.45m, green with darker markings, with up to 10 fls. Very early to early.

‘Orange Cameron’ VIII(c/d)

henryi seedling hybridized by J. Hoell × (“Tetra Black Beauty” × “Tetra White Henry”)

H: M. Simpson, 2003 **G:** M. Simpson, pre-2009

N: M. Simpson, 2009 **REG:** M. Simpson, 2009

Inside of tepals dark orange, outside of tepals paler orange. Spots dark brown; papillae orange, in throat only; nectaries green; pollen brown; stigma purple-grey. Fls 160mm wide, slightly scented; tepals 201 × 35mm, margins slightly ruffled, tips recurved to strongly recurved. Lvs 201 × 38mm, scattered. Stems 1.8m, dark green, with up to 31 fls. Nomenclatural standard: colour prints supplied by registrant (WSY 0112958)

‘Orange Flash’ I(a-b/b)

Un-named, orange-flowered Div. I (Asiatic hybrid) seedling derived from (white-flowered sport Div. I × ‘Sunkissed’) × un-named, orange-flowered Div. I seedling derived from (white-flowered sport Div. I × ‘Sunkissed’)

H: J.F. McRae, pre-1988 **G:** J.F. McRae, 1988

I: Mount Hood Lilies, Inc., c.1993

Inside of tepals with a pure white central blaze up to 20mm wide at base, tapering to 5mm wide towards tips, shading to vivid reddish orange (33B) to strong yellowish pink (33C) towards margins and strong reddish orange (34C) beside nectaries at base; becoming paler with age. Spots of medium size, magenta-black, sparsely distributed over strong reddish orange base of each tepal; nectaries soft green

to pale ivory in low light, pale orange in bright light, pubescent, with short white hairs; filaments 80–100mm, very pale orange to white; pollen greyish red (178A) to moderate red (182A), bleaching to strong orange (170A) under very hot conditions; style 60–80mm, very pale orange; stigma moderate red (183C/D). Buds 90–120 × 100–140mm, moderate yellowish pink (33D) with green midribs. Fls 140–180mm wide, not scented; tepals with a satin sheen, broad, outer acute, 22.5–30mm wide, inner 30–35mm wide, tips recurved; pedicels 80–120mm, glossy, dark plum, due to dark staining over dark green ground. Lvs 10–15 × 7.5–20mm, leathery, glossy, pale to mid-green. Stems up to 1.25m, with up to 10 fls in a racemose infl. Mid-season.

‘Orange Matrix’ I(a/b)

Sport of ‘Matrix’

H: Bischoff Tulleken Lelies BV, pre-2008 **N:** “The Originals” U.A., Coöp. Kwekersvereniging, 2008

REG: “The Originals” U.A., Coöp.

Kwekersvereniging, 2009

Syn. ‘Matrix Orange’

Inside of tepals strong orange (N25A); outside of inner tepals predominantly N25A. Spots few, dark red; papillae present; nectaries orange; pollen red-brown; stigma purple-red. Fls 120mm wide; tepals short and narrow, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of short to medium length, and of narrow to medium width, dark green. Stems 0.5m, green with darker spots and stripes, with up to 10 fls. Very early to early. Floriferous. Published in *The Lily Nook cat.*, 2009, as ‘Matrix Orange’

‘Orange Miss’ VI(b/b)

‘Orange Best’ × ‘Oranža Planeta’

H: R. Widerski, pre-2008 **N:** R. Widerski, 2008

I: R. Widerski, 2008 **REG:** R. Widerski, 2008

Inside of tepals apricot-orange, paler towards tips; throat darker. Midribs greenish, tinged purple and orange. Spots absent; nectaries green; pollen orange-brown; stigma dark purple. Fls 150mm wide; inner tepals 120 × 45mm, tips recurved; with secondary buds. Lvs 80 × 15mm, dull, mid-green. Stems 1.3m, with dark purple spots, with infl. of average length. Mid-July.

‘Orange Twins’ I(a-b/b)

‘Fata Morgana’ F₂ × ‘Fata Morgana’ F₂

H: Laan Flora Facilities BV, pre-2009 **REG:** Laan Flora Facilities BV, 2009

Inside of inner tepals vivid yellowish pink (28A), shading to strong orange (N25B) along midveins. Outside of inner tepals vivid orange (28B), shading to vivid yellowish pink (28A) towards margins. Spots few to moderately numerous, dark; papillae present; nectaries orange; pollen absent; stigma orange-red. Fls double, 90mm wide; tepals (very) short and narrow, margins slightly ruffled, tips slightly recurved. Lvs of medium length, and of narrow to medium width. Stems 1–1.35m, green, with up to 8 fls. Early to mid-season.

‘Oranža Planeta’

Pollen parent of ‘Orange Miss’. No further details known.

‘Ormea’ VIII(b-a/b) OT

Parentage unknown

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals light greenish yellow (5D) in basal half; shading to pale yellow-green (155A) towards tips; midveins flushed deep yellowish pink (43C); throat yellowish white (155D). Inside of outer tepals pale greenish yellow (2D), flushed yellowish white (155D) towards tips; margins light greenish yellow (3C); midveins flushed deep yellowish pink (43C). Outside of outer tepals pale yellow-green (155A), shading to pale greenish yellow (1D); midribs strong yellow-green (143B) at base, with short, vivid reddish orange (42A) stripe on one side. Outside of inner tepals pale yellow-green (155A); margins pale greenish yellow (2D) at base; midribs strong yellow-green (143C) at base. Spots absent; papillae light greenish yellow (5D); nectaries strong yellow-green (143C); pollen greyish brown (166A); stigma greyish purplish red (N77D). Fls 190mm wide, scented; tepals 115 × 55mm, margins smooth, tips recurved. Lvs 190 × 27mm. Stems 1.2m, green, with up to 6 fls. Named after a town in the region of Piedmont, Italy.

‘Papaya’ VIII(a/b) LA

Div. I (Asiatic hybrid) seedling × Div. VIII (Other hybrids) LA seedling

H: De Jong Beheer BV, pre-2007 **N:** De Jong Lelies Holland BV, 2007 **REG:** De Jong Lelies Holland BV, 2008

Inside of tepals strong orange (25B), shading to vivid orange (28B) towards margins; midveins strong orange (25A); throat pale yellow (18D). Outside of outer tepals light yellow (20B), inner tepals light orange-yellow (23C), all shading to vivid orange (28B) towards margins; midribs strong yellow-green at base (outer 143B, inner 143C). Spots absent; papillae few, dark red (59A); nectaries moderate olive-green (146A); pollen greyish brown (166A); stigma moderate purplish red (58A). Fls 210mm wide; tepals 130 × 60mm, margins smooth, tips slightly recurved. Lvs 160 × 17mm. Stems 1.4m, green with darker markings mostly towards base, with up to 7 fls.

‘Paradero’ VII(a/c)

Parentage unknown

H: Mak ‘t Zand BV, pre-2005 **N:** Mak ‘t Zand BV, 2005 **REG:** Mak ‘t Zand BV, 2008

Inside of tepals deep red (60A) to strong purplish red (60D), paler towards tips and nearly white towards margins; more blue with age; midveins more red. Outside of inner tepals with strong purplish red (60D) stripes on a pale ground. Spots and papillae absent; nectaries yellow-green; pollen orange-brown; stigma grey-green. Tepals long and of medium width, margins ruffled to strongly ruffled, tips slightly recurved. Lvs of medium length and broad. Stems 1.2m, green with darker spots and stripes, with up to 3 fls. Mid-season.

‘Park Rudy’ VI(c-d/b)

‘Black Heart Orange’ × unknown

H: R. Widerski, pre-2007 **N:** R. Widerski, pre-2007

REG: R. Widerski, 2007 **I:** R. Widerski, 2007

Inside of tepals orange, tips yellow-orange; midribs orange; throat pale orange, with some brown dashes. Nectaries greenish; pollen bronze-brown; stigma grey-green. Fls 135mm wide; tepals 120 × 25mm, tips recurved; pedicels 140mm, stout, pale green; with secondary buds. Lvs 200 × 15mm, mid-green, glossy. Stems 2m, green; infl. long. Mid-July.

‘Parrawe’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2007

Tepals light greenish yellow (4B); midribs lightly marked with moderate reddish orange (173B); throat pale yellow-green (4D), frosted. Spots rather large, mostly round, chocolate brown, densely and evenly distributed overn basal half of each tepal; nectaries pale green, frosted; pollen golden brown; stigma pale green. Fls 65mm wide, not scented; tepals 60 × 27mm, margins smooth, tips strongly recurved. Lvs 85 × 6mm, dark green. Stems 0.7m, green, flecked brown, with up to 10 fls in pyramidal infl. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “cease” or “abstain”.

‘Pascoli’ VII(a/b)

Seedling 93-14 × seedling RM 98-25

H: Vletter en Den Haan Beheer BV, pre-2006

N: Vletter en Den Haan Beheer BV, 2006

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals moderate purplish pink (186D); inside of outer tepals deep pink (52C); tips deep purplish pink (64D); all margins, base above nectaries, and throat, yellowish white (155D). Outside of outer tepals deep purplish pink (64D), shading to pale purplish pink (62D) towards margins; margins yellowish white (155D); midribs strong yellow-green (base 144B, top 143A/B and N144D between). Outside of inner tepals yellowish white (155D) flushed pale purplish pink (65D), shading to moderate purplish pink (65A) beside midribs; midribs strong yellow-green (144B). Spots and papillae yellowish white (155D); nectaries light yellow-green (145B); pollen moderate red (180A); stigma pale yellow-green (193C). Fls 245mm wide, scented; tepals 155 × 92mm, margins ruffled, tips straight. Lvs 185 × 53mm. Stems 1.1m, green with darker markings, with up to 3 fls. ‘China Pink’ was used as a temporary name prior to registration.

‘Patricia’s Pride’ I(a-b-c)

Parentage unknown

H: J. Wademaker, pre-2006 **N:** Lily Company BV, 2006 **REG:** Lily Company BV, 2008

Syn. ‘Purple Rain’ (Wademaker)

Inside of tepals, including throat, pale greenish yellow (2D), becoming whiter with age, with a large, dark greyish purple (202A) central patch shading to dark red

(187B) towards margins. Outside of outer tepals pale greenish yellow (2D); inside dark patch showing through at margins; midribs strong yellow-green (144B) at base and top, moderate yellow-green (143D) between. Outside of inner tepals pale yellow-green (4D); midribs strong yellow-green (143B) at base shading to moderate yellow-green (143D). Spots and papillae dark red (187B); nectaries strong yellow-green (144B); pollen greyish brown (166A); stigma pale yellow (11D) with narrow, dark red (187B) margin. Fls 150mm wide; tepals 85 × 37mm, margins smooth, tips slightly recurved. Lvs 130 × 20mm. Stems up to 1.3m, dark, with up to 9 fls.

‘Pavane’ VIII(a/b) LA

Parentage unknown

H: Vletter en Den Haan Beheer BV, 2005 **N:** Vletter en Den Haan Beheer BV, 2005 **REG:** Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals strong orange (N25A); throat light orange (28D). Inside of outer tepals vivid orange (28B). Outside of outer tepals light orange-yellow (22C), shading to strong orange (N25A) towards margins; midribs moderate yellow-green (138B) at base, strong yellow-green (143B) at top, light yellow-green (150D) between. Outside of inner tepals strong orange (25A/N25A); midribs strong yellow-green (143C) at base, shading to light yellow-green (145B). Spots sparse, deep red (60A); papillae absent; nectaries and pollen greyish brown (166A); stigma moderate red (181C). Fls 190mm wide; tepals 140 × 65mm, margins smooth, tips recurved. Lvs 135 × 35mm. Stems 1.3m, green, with up to 3 fls.

Named after a slow processional dance common in Europe in the 16th and 17th centuries.

‘Peach Blush’ VI(b-c/a)

Parentage unknown

H: L.C. Bligh, 2003 **G:** L.C. Bligh, 2005

N: L.C. Bligh, 2008 **REG:** L.C. Bligh, 2008

Inside of tepals pale yellow-green (4D); throat vivid yellow (14A). Outside of tepals pale yellow-green (4D) at base and margins, shading to moderate purplish red (185C) beside midribs. Spots absent; nectaries green; pollen golden yellow; stigma brown. Fls 155mm wide, strongly scented; tepals 140 × 68mm, margins slightly ruffled, tips recurved. Lvs 165 × 11mm, scattered. Stems 1.4m, green, with up to 18 fls. Mid-January (Tasmania).

‘Peachy Keen’ I(a)

H: Van der Salm (USA), pre-2008

Inside of tepals peach, shading to dark apricot at centre. Fls large. Stems 1.07m. Mid-season.

Published in *The Lily Nook cat.*, 2009

‘Pearl Jennifer’ I(b-a/c)

Parentage unknown

H: R.J. Griesbach, pre-2008 **N:** Lily Company BV, 2008 **REG:** Lily Company BV, 2008

Inside of tepals vivid yellow (12A); throat light greenish yellow (3C). Outside of outer tepals brilliant greenish yellow (6C), shading to vivid yellow (9A) towards margins; immediate tips moderate red (180A). Outside of inner tepals vivid yellow (9A), shading to brilliant greenish yellow (6A) beside midribs, strong

yellow-green (143B) at base. Spots dark red (59A); papillae absent; nectaries strong yellow-green (143A); pollen dark reddish orange (173A); stigma dark red (183A). Fls 170mm wide; tepals 93 × 52mm, margins smooth, tips recurved. Lvs 84 × 12mm. Stems 1.2m, pale green with darker markings, with up to 4 fls.

‘Pearl Jessica’ I(b-c/b-c)

Parentage unknown

H: R.J. Griesbach, pre-2008 **N:** Lily Company BV, 2008 **REG:** Lily Company BV, 2008

Inside of inner tepals pinker than moderate orange-yellow (168D), shading to light yellowish pink (38D) towards tips; becoming more pink with age; throat light orange-yellow (19A), shading to light yellowish pink (38D) at upper margin. Inside of outer tepals light orange-yellow (19A); midveins, and towards tips, moderate yellowish pink (39C). Outside of tepals light yellowish pink (38D); margins narrow, light yellow (20B); midribs strong yellow-green (144A) at base, shading to dark violet at top of outer tepals, and dark greenish yellow (152C) on inner tepals. Spots very few, dark red (187A); papillae very few or absent; nectaries strong yellow-green (144A); pollen strong reddish orange (169A); stigma deep purplish red (71A). Fls 160–170mm wide; tepals 90 × 25–40mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs 90 × 15mm. Stems 0.7m, green, with up to 7 fls. Tetraploid.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Pearl Justien’ I(-/b-c)

H: R.J. Griesbach, pre-2008 **N:** Lily Company BV, 2008

Tepals bright orange. Fls large, of heavy substance.

Stems 1.2m, erect. Mid-season. Tetraploid.

Published in *The Lily Nook cat.*, 2009

‘Pearl Loraine’ I(b-c/b-c)

Parentage unknown

H: R.J. Griesbach, pre-2008 **N:** Lily Company BV, 2008 **REG:** Lily Company BV, 2008

Inside of inner tepals moderate reddish orange (179C), with large, moderate red (179A) patch at base; margins narrow, brilliant orange (25C); midveins moderate red (180C); throat light yellowish pink (36A), tinged strong greenish yellow (151A) beside nectaries, and strong red (51A) at upper margin. Inside of outer tepals moderate red (179A), shading to 179B to moderate reddish orange (179C) towards margins and tips; margins narrow, brilliant orange (25C). Outside of tepals dark yellowish pink (181D); margins strong red (51A); in outer tepals in basal one-third only, extending further in inner tepals; vivid reddish orange (30A) above; midribs light yellowish pink (27A), all strong yellowish green (144A) at base, and at top of outer tepals only. Spots deep reddish purple (77A) to dark greyish purple (202A); papillae few; nectaries strong yellowish green (N144A); pollen moderate red (179A); stigma deep red (60A). Fls 180mm wide; tepals 90 × 30–50mm, margins smooth to slightly ruffled, tips slightly recurved to recurved. Lvs 120 × 15mm, dark green. Stems 0.95m, green with darker spots, with up to 4 fls.

'Pearl Stacey' I(b-c/c)

Parentage unknown

H: R.J. Griesbach, pre-2008 **N:** Lily Company BV,

2008 **REG:** Lily Company BV, 2008

Inside of tepals light orange-yellow (22B); brilliant orange-yellow (23B) towards throat and shading to light yellowish pink (26D) towards tips; margins narrow, strong orange (26B); outer tepals of more intense colouring; throat pale orange-yellow (23D), strong yellow-green (144A) beside nectaries. Outside of tepals pale orange-yellow (24D); margins narrow, strong orange (24A) to strong orange-yellow (24B), and greyish purple (N77A) at top of outer tepals only; midribs strong yellow-green (144A) at base shading to pale greenish yellow (1D). Spots absent; papillae few; nectaries moderate yellow-green (146B); pollen strong reddish orange (169A); stigma deep purplish red (61A). Fls 150mm wide; tepals 85 × 30–50mm, margins smooth, tips strongly recurved. Lvs 110–150 × 10–15mm, dark green. Stems 1m, green with darker markings, with up to 12 fls.

'Peggy Wharrier' I(a-b/b-c)

White-flowered seedling derived from *lancifolium* × red-flowered Div. I (Asiatic hybrid) seedling

H: R. Hall, 2005 **G:** R. Hall, 2007 **N:** R. Hall, 2007

REG: R. Hall, 2008

Inside of tepals lime-green, flushed lilac-pink, particularly towards margins and tips; throat purple. Outside of tepals lilac-pink, mottled lime-green. Spots dark purple, of medium size, oval, sometimes elongated, well scattered over basal half of each tepal; nectaries lime-green, flushed lilac-pink; pollen yellow; style red-purple; stigma greenish white. Fls 120mm wide, not scented; tepals 80 × 50mm, margins smooth, tips slightly recurved. Lvs 80 × 20mm, scattered. Stems 1m, green, with up to 8 fls. Named after Mrs Hall's late mother.

'Penghana' I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2007

Inside of tepals strong reddish purple (70B) overlaid with pale yellowish pink (159D) towards centre; throat 159D. Outside of tepals moderate purplish pink (70D); midribs marked greyish reddish orange (174B). Small, V-shaped pencil-mark on each tepal and spots mahogany-brown; spots of medium size, round to elliptic, scattered over basal two-thirds of each tepal; nectaries pale green, frosted; pollen golden brown; stigma pale green. Fls 65mm wide, not scented; tepals 60 × 25mm, margins smooth, tips recurved. Lvs 75 × 4mm, scattered. Stems 0.65m, green, flecked brown, with up to 10 fls. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for a "river ford".

'Penthouse' VII(a/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** W. Vink,

2003 **REG:** "The Originals" U.A., Coöp.

Kwekersvereniging, 2007

Inside of tepals strong purplish pink (68B), midveins and base pale yellow-green (155A). Outside of inner tepals light purple (75B) to very light purple (75C). Spots white and pink on pink ground; papillae present; nectaries green; pollen red-brown; stigma yellow-green. Fls 180mm wide; tepals of medium length and width, margins slightly ruffled to ruffled, tips recurved. Lvs of medium length, and broad. Stems 1.1m, green with darker markings, with up to 5 fls. Late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

External images: *The Lily Nook cat.*, 2009

'Petit Brigitte' I(a/b-c)

Tepals glossy, bright canary-yellow. Spots and papillae absent; pollen orange-brown. Tepal margins smooth, tips recurved. Lvs dark green. Stems 0.45m. Early to mid-season.

'Pico' VII (a/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2005 **N:** Van der Marel Lelie BV, 2005 **REG:** Van der Marel Lelie BV, 2007

Inside of tepals deep red (60A) to strong purplish red (60B). Outside of inner tepals strong purplish red (60D). Spots and papillae present; nectaries yellow-green; pollen orange-brown; stigma purple. Tepals of short to medium length and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad to very broad. Stems of average height to tall, green with darker markings, with few to average no. of fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007).

'Picotee Lace' I(a)

H: Cebeco Lilies, Inc., pre-2008

Inside of tepals mid-soft red; margins yellow. Stems 1.07m. Mid-season.

Published in *The Lily Nook cat.*, 2009

'Pink Meadow' VII(b-a/b-c)

Div. VII (Oriental hybrid) seedling × Div. VII seedling **H:** De Jong Beheer BV, pre-2007 **N:** De Jong Lelies Holland BV, 2007 **REG:** De Jong Lelies Holland BV, 2008

Inside and outside of inner tepals deep purplish pink (N66D) in upper half; margins, basal half and throat yellowish white (155D). Inside of outer tepals deep purplish pink (N66D); margins, and patch above nectaries, yellowish white (155D). Outside of outer tepals deep purplish pink (N66C), base yellowish white (155D); all midribs strong yellow-green (outer tepals, base 143C, top 143A, 144C between; inner tepals 144C). Spots and papillae yellowish white (155D); nectaries strong yellow-green (143C); pollen

strong orange (N25A); stigma pale yellow-green (193D). Fls 235mm wide, scented; tepals 130 × 80mm, margins ruffled, tips slightly recurved. Lvs 235 × 65mm. Stems 1.1m, green, with up to 4 fls.

‘Pink Pixels’ I(a/b-c)

‘Cha Cha Cha’ F₂ × ‘Cha Cha Cha’ F₂

H: Laan Flora Facilities BV, pre-2009 REG: Laan Flora Facilities BV, 2009

Inside of tepals strong purplish pink (73B); throat yellowish white (155D). Outside of inner tepals mainly strong purplish pink (63C). Spots numerous, at base of each tepal, extending along margins, dark red (187B); papillae present; nectaries green shading to purple-red; pollen orange-red; stigma orange. Fls 110mm wide; tepals (very) short and narrow, margins slightly ruffled, tips slightly recurved. Lvs of medium length, and of narrow to medium width. Stems 1.2–1.45m, green with darker markings, with up to 7 fls. Early to mid-season.

‘Pink Secret’ VII(a/b)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

N: Marklily BV, 2007 REG: Testcentrum voor

Siergewassen BV, 2008

Inside of tepals flushed strong purplish pink (68B) on white ground, shading to white towards base. Outside of inner tepals greenish white (155C), flushed strong purplish pink (68B) towards tips. Spots absent; papillae present; nectaries green; pollen orange; stigma yellow-green. Tepals of medium length to long and of medium width to broad, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad. Stems 1.15m, green with darker spots and stripes, with up to 3 fls. Mid-season.

‘Pink Wish’ VII(a/b)

Div. VII (Oriental hybrid) seedling × Div. VII seedling

H: De Jong Beheer BV, pre-2007 N: De Jong Lelies Holland BV, 2007 REG: De Jong Lelies Holland BV, 2008

Inside of inner tepals dark purplish pink (186C), shading to deep purplish pink (64D) towards margins; margins light purplish pink (65B); patch of light yellow-green (2C) above nectaries; throat yellowish white (155D). Inside of outer tepals strong purplish pink (63C), shading to deep purplish pink (64D) towards margins; margins light purplish pink (65B); a small light yellow-green (2C) stripe above nectaries. Outside of tepals moderate purplish pink (186D), shading to deep purplish pink (N66D) towards margins and yellowish white (155D) beside midribs at base; margins of inner tepals pale purplish pink (65D); midribs of outer tepals strong yellow-green (base 143B, shading to 144C); midribs of inner tepals 144C, flushed moderate purplish pink (186D) at top. Spots few, strong purplish red (58B); papillae yellowish white (155D); nectaries strong yellow-green (143C); pollen dark orange (N163A); stigma dark purplish red (N79B). Fls 255mm wide, scented; tepals 160 × 85mm, margins slightly ruffled, tips slightly recurved. Lvs 170 × 55mm. Stems 1m, uppermost third green, lower two-thirds dark, with up to 3 fls.

‘Pirax’ VIII(a/b) LA

Seedling BW 96-47 × seedling AM 95-12

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals moderate purplish pink (186D), shading to dark purplish pink (186C); base tinged strong purplish red (58C); throat pale yellowish pink (36D). Inside of outer tepals moderate purplish pink (186D) tinged yellowish white (155D), shading to dark purplish pink (186C) towards margins; tips tinged strong purplish red (58C). Outside of outer tepals dark purplish pink (186C) tinged yellowish white (155D), shading to deep purplish pink (N57D) towards margins; midribs at base yellowish white (155D) tinged light yellow-green (145B), shading to strong yellow-green (145A) at top. Outside of inner tepals light purplish pink (63D); margins near base strong purplish pink (63C); midribs strong yellow-green (145A) at base. Fls paler with age. Spots strong purplish red (60B); papillae absent; nectaries moderate olive-green (147A); pollen vivid reddish orange (N30B); stigma light orange-yellow (25D). Fls 185mm wide, slightly scented; tepals 120 × 64mm, margins smooth, tips slightly recurved. Lvs 153 × 35mm. Stems 1.3m, green, with up to 6 fls.

‘Piston Cup’ VII(a/c)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

N: Hobaho BV, 2007 REG: Testcentrum voor

Siergewassen BV, 2009

Inside of tepals vivid purplish red (N57B) to deep purplish pink (N57C), paler towards tips and becoming paler with age, shading to yellowish white (155D) at margins; midveins deep red (60A) to strong purplish red (60B) at base. Outside of inner tepals deep purplish pink (N57D). Spots darker than tepal colour; papillae present; nectaries green; pollen orange-brown; stigma purple. Tepals of medium length to long and of medium width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length, and of medium width to broad, dark green. Stems tall, green, with up to 4 fls. Mid-season.

‘Poem’ I(a/b)

Seedling AZ 96212-1 × ‘Jonita’

H: De Jong Beheer BV, pre-2006 N: De Jong Lelies

Holland BV, 2006 REG: De Jong Lelies Holland BV, 2008

Inside of inner tepals vivid reddish orange (30A); margins and midveins N30C; throat strong reddish orange (32B). Inside of outer tepals vivid reddish orange (30A), shading to N30B towards margins. Outside of outer tepals strong orange (N25A), shading to vivid reddish orange (30A) towards margins. Outside of inner tepals vivid yellowish pink (28A). Spots numerous, dark red (59A); papillae vivid reddish orange (30A); nectaries moderate olive-green (147A); pollen strong orange (N25A); stigma vivid reddish orange (34B). Fls 200mm wide; tepals 105 × 57mm, margins smooth, tips straight. Lvs 100 × 27mm. Stems 1.2m, green, with up to 5 fls.

‘Polar’ VIII(b-a/b) LO

White-flowered Div. VII (Oriental hybrid) × seedling LO 93003 AB T

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2008

Tepals, including throat, yellowish white (155D); midveins of outer tepals tinged light greenish yellow (5D) above nectaries; midribs strong yellow-green (144C) at base, those of outer tepals tinged light yellow-green (144D) at top. Spots absent; papillae few, yellowish white (155D); nectaries strong yellow-green (144B); pollen strong orange (N163B); stigma light greenish grey (190D). Fls 210mm wide, scented; tepals 140 × 70mm, margins ruffled, tips recurved. Lvs 185 × 35mm. Stems 1.1m, green, with up to 4 fls.

‘Polymer’ VI(b-c/a)

(*regale* tetraploid form × ‘Tetra White Henry’) × (‘Tetra White Henry’ × (‘Tetra White Henry’ × unknown))

H: B. Mičulka, pre-2008 **N:** B. Mičulka, 2008

I: B. Mičulka, 2008 **REG:** B. Mičulka, 2008

Tepals white, inside of tepals with dark apricot rays in basal two-thirds which show through to outside of tepals; midribs speckled pale brown. Spots absent; nectaries dark green; pollen apricot-brown; stigma pale yellow-green. Fls 170mm wide; tepals 140 × 50mm, margins (very) slightly ruffled, tips recurved; pedicels 110mm, slightly spotted dark brown. Lvs 155 × 15mm, dull, dark green. Stems 1.3m, spotted brownish black, with infl. of average length. Mid-July.

‘Portage Gold’ I(a/b-c)

H: W.G. Ronald, pre-2008

Inside of tepals concolorous clear golden yellow. Spots chocolate-brown, evenly sprinkled towards base of each tepal; outer spots larger and more elongated than inner ones; nectaries white frosted; filaments slightly darker, duller golden yellow than inside of tepals, becoming sulphur-white; anthers slender; pollen russet-orange; stigma and style pinkish bronze. Tepals of medium length and width, margins smooth to very slightly ruffled, tips becoming strongly recurved. Stems 1.07m. Mid-season. Tetraploid. Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

‘Portage Orange’ I(a/b-c)

H: W.G. Ronald, pre-2008

Inside of tepals shimmering dark apricot-orange. Spots few, dark brown, small, evenly sprinkled towards base of each tepal, outer spots more elongated; filaments paler apricot-orange than inside of tepals; pollen dark chocolate-brown; stigma orange. Tepals of medium length and width, margins smooth to very slightly ruffled, tips strongly recurved. Stems 1.07m. Mid-season. Tetraploid. Published in *The Lily Nook cat.*, 2009
External images: *The Lily Nook cat.*, 2009

‘Prajská Hvězda’ VI(b-c/d)

‘Henrietta’ × ‘Bright Star’

H: J. Dudek, pre-2007 **N:** J. Dudek, pre-2007

I: J. Dudek, 2007 **REG:** J. Dudek, 2007

Inside of tepals white with yellow rays. Outside of tepals white; midribs greenish yellow, creamy at top. Brushmarks fine, purple, over one-third of each tepal; nectaries green; pollen orange; stigma pale green. Fls 65mm wide; inner tepals 68 × 27mm, margins slightly ruffled, tips strongly recurved; pedicels 65mm, stout, dark green; with secondary buds. Lvs 88 × 36mm, mid-green, glossy. Stems 1.25m, green; infl. of average length. Late July.

‘Pratolo’ VIII(a/c-b) LA

Seedling BG 96-15 × seedling AG 96-26

H: Vletter en Den Haan Beheer BV, pre-2006

N: Vletter en Den Haan Beheer BV, 2006

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals vivid yellow (9A); throat pale greenish yellow (2D). Outside of outer tepals light greenish yellow (8B); margins vivid yellow (12A); midribs strong yellow-green (base 143C, top 143A). Outside of inner tepals brilliant yellow (8A), shading to vivid yellow (12A) towards margins; midribs pale greenish yellow (2D) at base. Spots and papillae absent; nectaries strong yellow-green (143B); pollen dark red (183A); stigma light yellow (15D) with dark red (59A) stripes. Fls 240mm wide; tepals 130 × 75mm, margins smooth, tips slightly recurved. Lvs 145 × 25mm. Stems 1.05m, green, with up to 3 fls.

‘Premio’ VII (a/b)

Div. VII (Oriental hybrid) seedling × Div. VII seedling

H: De Jong Beheer BV, pre-2007 **N:** De Jong Lelies Holland BV, 2007 **REG:** De Jong Lelies Holland BV, 2008

Inside of tepals strong purplish red (60D); margins tinged moderate purplish red (70A); tips of inner tepals tinged yellowish white (155D); basal half and top of midveins of outer tepals yellowish white (155D); throat yellowish white (155D). Outside of outer tepals strong purplish red (60C), yellowish white (155D) beside midribs at base; margins tinged deep purplish red (71A); midribs strong yellow-green (base 144A shading to 143A). Outside of inner tepals strong purplish red (60D), yellowish white (155D) beside midribs at base; margins tinged moderate purplish red (70A); midribs strong yellow-green (base 144C shading to 144A towards top). Spots deep purplish red (59B); papillae yellowish white (155D) tinged 59B; nectaries strong yellow-green (144C) edged pale greenish yellow (2D); pollen moderate reddish orange (171A/N172A); stigma light greenish grey (190C). Fls 225mm wide, scented; tepals 130 × 80mm, margins smooth, tips straight. Lvs 200 × 45mm. Stems 0.9m, green, with up to 5 fls.

‘Pretty Maid’ VIII(a-b/a) OT

Pink-flowered (a/a) unregistered seedling × (‘Starburst Sensation’ × (*L. × kewense* ‘White Henryi’ × ‘Dominique’))

H: R.H. Bazett, 1997 **G:** R.H. Bazett, 2000

N: R.H. Bazett, 2007 **REG:** R.H. Bazett, 2007

Inside of each tepal purple-red in central two-thirds, shading to white towards margins and tips; midveins dark purple-red; throat pale yellow. Outside of outer tepals greyed purple-red, paler towards margins; midribs purple-red tinged grey-green towards base. Outside of inner tepals greyed pink, shading to white towards margins; midribs purple-red tinged grey-green towards base. Spots and papillae absent; nectaries dark green; pollen orange-brown; stigma grey-green. Fls 135mm wide, strongly scented; tepals 145 × 50mm, margins slightly ruffled, tips recurved. Lvs 205 × 20mm, dark green. Stems 2m, grey-green, with up to 18 fls in racemose infl. July. Triploid.

‘Prima Donna’ I(a/c)

Un-named, soft-yellow-flowered Div. I (Asiatic hybrid) (a/-) seedling × un-named, pink-tipped, white-flowered Div. I seedling

H: R. Mak-Leek, pre-1989 **G:** R. Mak-Leek, 1989

I: Mak-Leek, Inc., c.1998

Inside of tepals yellowish white (158C) to light yellowish pink (159A), shading to darker pink towards margins; tips light purplish pink (68D); base brilliant yellow (13B) to vivid yellow (14B). Spots small, greyish red (178A), in basal one-third of each tepal; filaments 45–60mm, pale green to very pale pink; pollen brownish orange (171B); stigma moderate purplish red (185C). Buds 60–80mm long; light yellowish pink (159A); dark red (183B) at tip; midribs close to strong yellow-green (144B). Fls 145–165mm wide, with a satin sheen, not scented; outer tepals 25–35mm wide, inner 40–50mm wide, tips slightly recurved; pedicels 50–80mm, strong yellow-green (144A) to moderate olive-green (146A). Lvs 100–140 × 15–25mm, leathery, glossy, deep yellowish green (141A) above, paler beneath. Stems up to 0.75m, with up to 14 fls in a racemose infl. Mid-season.

‘Pup Art’ I(a/-)

N: W. Vink, 2003

Inside of tepals bright red. Spots burgundy, coalescing to form a ring towards the centre of the flower. Stems 1.07m. Mid-season.

Published in *The Lily Nook cat.*, 2009

‘Purple Rain’ (Wadecomper) I(a/b-c)

Syn. of ‘Patricia’s Pride’

‘Rebound’ VII (a/b-c)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

N: Testcentrum voor Siergewassen BV, 2007

REG: Testcentrum voor Siergewassen BV, 2008

Inside of tepals light purplish pink (73C) to very pale purple (73D), paler towards tips and base; light greenish yellow (8C) to pale yellow (8D) beside midveins at base and along midveins. Outside of inner tepals light purplish pink (73C) to very pale purple (73D). Spots yellow; papillae present; nectaries green; pollen orange-brown; stigma grey-green. Tepals of medium length and width, margins ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad to very broad. Stems 1.2m, green, with up to 8 fls. Mid-season to late.

‘Red Empire’ VII(a/c)

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV, 2003

REG: World Breeding BV, 2008

Inside of tepals deep red (53A/60A), becoming more blue with age; paler towards margins and tips; margins very narrow, white (NN155C). Outside of tepals strong purplish red (60C). Spots and papillae present; nectaries green shading to yellow; pollen red-brown; stigma grey-green, flecked purple. Fls 240mm wide; tepals of medium length and width, margins strongly ruffled, tips slightly recurved to recurved. Lvs of medium length, and of medium width to broad. Stems 0.95–1.1m, green with some darker markings, with up to 6 fls. Mid-season.

Published in *The Lily Nook cat.*, 2009

‘Red Eyes’ VII(a/-)

Parentage unknown

N: H.P. Imanse BV, 2007 **REG:** H.P. Imanse BV, 2008

Inside of tepals strong purplish red (60B), shading to 60C towards margins; margins and tips paling to almost white; midveins deep red (60A). Outside of tepals deep purplish pink (N57D). Spots numerous, darker than tepals; papillae present; nectaries green, flushed red; pollen red-brown; stigma dark purple. Tepals of medium length to long and of medium width, margins strongly ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad. Stems of average height, green, with few to average no. of fls. Late.

RED IMPRESSION VII(a/b)

Trade designation of ‘Zanlopres’

‘Red Latin’ I(a/b)

Parentage unknown

H: Mak Breeding BV, pre-2008 **N:** Mak Breeding BV, 2008 **REG:** Mak Breeding Rights BV, 2008

Tepals vivid reddish orange (42A) to vivid red (45B), inside of tepals paler towards tips. Spots and papillae present; nectaries red; pollen reddish brown; stigma dark purple. Fls of average diameter; tepals short and of narrow to medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length, and of narrow to medium width. Stems 1.45m, green, with up to 11 fls.

‘Redpa’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2007

Tepals, including throat, strong red (47B); inside of tepals flushed light orange (28C) at centre. Spots of medium size, round, mahogany-brown, scattered over basal half to two-thirds of each tepal; nectaries red; pollen red-brown; stigma pale orange. Fls 70mm wide, not scented; tepals 55 × 20mm, margins smooth, tips recurved. Lvs 83 × 3mm, scattered. Stems 0.9m, brown, with up to 18 fls. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for ‘mosquito’.

'Red Reflex' VII(a/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** W. Vink, 2003 **REG:** "The Originals" U.A., Coöp. Kwekersvereniging, 2007

Inside of tepals deep red (60A), paler towards tips; more blue with age. Outside of inner tepals strong purplish red (60C). Spots and papillae present; nectaries yellow-green; pollen red-brown; stigma grey-green. Fls 160mm wide; tepals of medium length and width, margins ruffled to strongly ruffled, tips recurved. Lvs of medium length, and broad. Stems 1–1.25m, green, with up to 5 fls. Mid-season. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

'Red Strap' I(c/d)

H: C.F. Patterson, pre-2008

Tepals dark cherry-red with a lustrous sheen. Spots black, raised; pollen orange-red. Stems 0.8m. Mid-season.

Published in *The Lily Nook cat.*, 2009

'Red Twin' I(a/b)

Div. I (Asiatic hybrid) seedling × Div. I seedling

H: De Jong Beheer BV, pre-2007 **N:** De Jong Lelies Holland BV, 2007 **REG:** De Jong Lelies Holland BV, 2008

Inside of tepals moderate reddish orange (35A) to vivid reddish orange (40B), with speckled patch of dark red (59A) above nectaries; throat vivid reddish orange (30A). Outside of outer tepals light olive (152B) flushed deep red (60A); margins vivid reddish orange (30A) towards tips and deep red (60A) closely spotted vivid reddish orange (N30A) towards base. Outside of inner tepals strong reddish orange (31B); margins vivid reddish orange (N30A) towards tips and strong red (53C) towards base; all midribs light olive (152B); those of outer tepals greyish olive-green (N137A) at top. Papillae absent; nectaries moderate olive-green (137B); filaments petaloid, broad at base, linear in upper half; stigma dark purple (79A). Fls 180mm wide, double; tepals 105 × 42mm, margins smooth to very slightly ruffled, tips straight. Lvs 135 × 25mm. Stems 1.1m, dark at base, shading to green, with up to 7 fls.

'Relucida' VIII(a-b/b) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

N: Testcentrum voor Siergewassen BV, 2007

REG: Testcentrum voor Siergewassen BV, 2008

Inside of tepals strong purplish red (60B/C), becoming more blue with age. Outside of inner tepals purple. Spots and papillae present; nectaries purple-red, greenish at base; pollen orange-yellow; stigma purple. Tepals short and of narrow to medium width, margins slightly ruffled, tips (very) slightly recurved. Lvs of medium length and width. Stems 1.35m, green with faint darker spots and stripes, with up to 10 fls. Mid-season to late.

'Remine' I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2003 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2007

Tepals brilliant greenish yellow (6A); midribs dark reddish orange (173A); throat light yellow-green (154D). Small, V-shaped pencil-mark on each tepal and spots mahogany-brown; spots round, of medium size, evenly scattered over basal half of each tepal; nectaries yellow-green; pollen golden orange; stigma yellow. Fls 70mm wide, not scented; tepals 65 × 25mm, margins smooth, tips slightly recurved. Lvs 90 × 6mm, scattered, dark green. Stems 0.8m, green, flecked brown, with up to 10 fls in pyramidal infl. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for *Blandfordia punicea* (Christmas bells).

'Reunion' VII(b/c-d)

speciosum var. *gloriosoides* × 'Siberia'

H: Q.F. Cheng, 2004 **G:** Q.F. Cheng, 2008

N: Q.F. Cheng, 2008 **REG:** Yongkang Zhejiang Jiangnan Lily Breeding Co. Ltd, 2008

Inside of tepals white, with pencil line of carmine-red at margins, in outer tepals in basal quarter to one-third; in inner tepals in basal half; midveins green at base, becoming paler towards top; throat whitish green. Outside of tepals white; midribs green. Spots and papillae large, oval, numerous, increasing in size, and sometimes merging, towards margins, carmine-red, over basal two-thirds of each tepal; nectaries green; pollen dark blackish brown; stigma purple. Fls 230mm wide, slightly scented; tepals 115 × 50mm, margins strongly ruffled, tips strongly recurved. Lvs 182 × 55mm, scattered. Stems 1.13m (under glass), green. External web links: www.lanlilium.com
Nomenclatural standard: colour print supplied by registrant (WSY 0112957)

'Rexona' VIII(a/b) OT

Parentage unknown

H: World Breeding BV, pre-2003 **N:** Liliande BV,

2003 **REG:** World Breeding BV, 2008

Inside of tepals and outside of inner tepals pale yellow (8D); young fls greenish white (155C). Spots and papillae absent; nectaries green; pollen red-brown; stigma purple, speckled. Fls 220mm wide; tepals of medium length to long and of medium width, margins slightly ruffled, tips recurved. Lvs of medium length to long, and of medium width to broad. Stems 1–1.3m, green with some darker markings, with up to 5 fls. Late to very late.

'Richmond' VIII(a/b) LA

Seedling BW 94-14 × seedling AW 94-09

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Tepals yellowish white (155D); throat pale yellow-green (149D). Midribs tinged strong yellow-green (144C) at base, with a dark pink (182C) spot or stripe above; midribs of outer tepals tinged strong

yellow-green (143C) at top. Spots and papillae absent; nectaries strong yellow-green (143C); pollen dark reddish orange (175B); stigma pale greenish yellow (2D). Fls 200mm wide, slightly scented; tepals 115 × 66mm, margins smooth, tips recurved. Lvs 170 × 30mm. Stems 1.2–1.3m, green with darker markings, with up to 6 fls.

‘Riley’ I(a/c)

Un-named, white-flowered, pollen-free seedling × un-named ‘Latvia’ seedling

H: T. Zimmerman, 2004 **G:** T. Zimmerman, 2006
N: T. Zimmerman, 2008 **REG:** T. Zimmerman, 2008
Inside of tepals white; throat purplish white. Outside of tepals greenish purple. Spots small, numerous, brownish burgundy-purple, stippling up to three-quarters of each margin, coalescing in basal half of each tepal to form patch speckled with white; nectaries white; pollen brown; stigma purple. Fls 120mm wide, not scented; tepals 70 × 35mm, margins slightly ruffled, tips slightly recurved. Lvs 100 × 15mm, scattered. Stems 1m, mid-green, with up to 7 fls. Early July.

‘Rinadeena’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005
G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2007

Tepals light greenish yellow (4C) to pale yellow-green (4D); midribs of inner tepals with small, moderate reddish orange (173B) patch; throat pale yellow-green (4D). Small, V-shaped pencil-mark on each tepal and spots mahogany-brown; spots of medium size, round, scattered over basal half of each tepal; nectaries pale green, frosted; pollen pale golden brown; stigma pale green. Fls 65mm wide, not scented; tepals 60 × 24mm, margins smooth, tips strongly recurved. Lvs 90 × 5mm, scattered. Stems 0.8m, green, flecked brown, with up to 10 fls. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “raindrop”.

‘Ring of Fire’ I(a/b-c)

‘Campfire’ × un-named seedling

H: T. Zimmerman, 1995 **G:** T. Zimmerman, 1997
N: P. Olson, 1998 **REG:** T. Zimmerman, 2008
Inside of tepals, including throat, vivid reddish orange (42A); semi-circular patch brilliant yellow (7A), with feathered upper margin, occupying basal third of each tepal. Outside of tepals reddish yellow; midribs green. Spots absent; nectaries yellow; pollen orange-brown; stigma yellowish beige. Fls 120mm wide, not scented; tepals 75 × 30mm, margins smooth, tips slightly recurved. Lvs 100 × 15mm, scattered. Stems 1m, dark green, with up to 14 fls. Early July.

‘River Rouge’ I(a/b-c)

Parentage unknown

H: Van der Salm (USA), pre-2008
Tepals concolorous: clear, lustrous, raspberry-red; a patch of blackish red in the centre of each tepal at the base. Spots few, small, blackish brown, sprinkled evenly over the basal third of each tepal; filaments reddish pink; pollen dark, rich orange-brown; style and stigma

purplish brown. Buds deep red. Tepal margins smooth, tips recurved. Stems 0.9m. Mid-season. Tetraploid. Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Rosy Dream’

Unregistered seed parent of ‘Meng Fan’ and pollen parent of ‘Zhonghua Huangguan’. No further details known.

‘Rouge’ VII(a/b)

Syn. of ‘Color Cosmetics’

External web links: www.lanlilium.com as ‘Rouge’ (image)

‘Royal’ VII(a/b)

‘Miami’ × ‘Sorbonne’

H: Q.F. Cheng, 2004 **G:** Q.F. Cheng, 2008

N: Q.F. Cheng, 2008 **REG:** Yongkang Zhejiang Jiangnan Lily Breeding Co. Ltd, 2008

Inside of tepals rich pink; margins and tips whitish; midveins of outer tepals darker; throat pale yellow. Outside of tepals pink; midribs pale green. Spots and papillae of moderate size, sparse, reddish pink, scattered over basal quarter of each tepal; nectaries green; pollen dark reddish brown; stigma green suffused with red spots. Fls 200mm wide, strongly scented; tepals 100 × 55mm, margins slightly ruffled, tips slightly recurved. Lvs 125 × 35mm, scattered. Stems 1.05m (under glass), yellow-green.

RUBATO VII(a/b-c)

Trade designation of ‘Zanlorbato’

‘Ruyi’ VII(b/b)

‘Con Amore’ × ‘Acapulco’

H: S. Zheng, 2002 **G:** S. Zheng, Mrs J. Huang, H. Zhang, 2006 **N:** S. Zheng, Mrs J. Huang, H. Zhang, 2007 **REG:** S. Zheng, Mrs J. Huang, H. Zhang, 2009

Tepals strong pink; throat yellow and green. Spots pinkish red, in basal quarter of each tepal; papillae white, at base of each tepal; nectaries green; pollen orange-brown; stigma milky white. Fls 225mm wide, strongly scented; tepal margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs 173 × 52mm, whorled. Stems 1.15m (under glass), green with purple spots, with up to 5 fls. May. Triploid. The epithet is a transcription of Mandarin Chinese meaning “good luck and happiness”.

‘Sabatini’ VIII(a/b-c) LA

Div. I (Asiatic hybrid) seedling × seedling LA 93008-6T

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2008

Inside of tepals moderate purplish red (184C); margins closely spotted dark purplish red (N186D); inner tepals with patch of yellowish white (155D) above nectaries; base of outer tepals light yellow-green (145C) beside nectaries; throat pale yellow-green (155A). Outside of outer tepals deep pink (185D) with strong yellow-green (144C) stripes, shading to moderate purplish red (186A) towards margins;

midribs strong yellow-green (base 144C, top 144B). Outside of inner tepals moderate purplish red (184B), shading to 186A towards margins; basal third pale greenish yellow (2D); midribs strong yellow-green (144C) at base shading to moderate purplish red (186A) towards top. Spots absent; papillae yellowish white (155D) and moderate purplish red (184C); nectaries strong yellow-green (143C); pollen greyish brown (166A); stigma purple-red. Fls 215mm wide; tepals 120 × 50mm, margins smooth, tips recurved. Lvs 170 × 25mm. Stems 1.1m, green, with up to 5 fls.

‘Sabella’ VII(a/b)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

N: Hobaho BV, 2007 **REG:** Testcentrum voor Siergewassen BV, 2009

Inside of tepals white (NN155C); outside of inner tepals predominantly NN155C. Spots absent; papillae conspicuous; nectaries pale green; pollen orange-brown; stigma purple. Tepals of medium length to long and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved. Lvs of medium length and of medium width to broad, dark green. Stems tall, green with darker spots and stripes, with up to 5 fls. Mid-season.

‘Sadie’ I(a/b-c)

Parentage unknown

H: Mak-Leek, Inc., pre-2009

Inside of tepals dog-rose pink in upper half, basal half creamy yellow, darkest towards centre of each tepal, feathering to milk-white towards margins, base and upper half. Spots and papillae absent. Filaments and style deep rose-pink; pollen old-gold yellow; stigma milk-white. Buds pinkish red, tip apple-green. Tepals slender, margins smooth, slightly incurving, tips slightly recurved. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Saldais Rīts’ II(c/d)

Un-named seedling × unknown

H: D. Hercbergs, pre-2003 **G:** D. Hercbergs, 2003

N: D. Hercbergs, 2004 **I:** D. Hercbergs, 2007

REG: D. Hercbergs, 2008

Tepals, including throat, vivid yellow (17B); midribs moderate purplish red (186A) at base. Spots strong purplish red (58B) over two-thirds of each tepal; nectaries vivid yellow (17B); pollen and stigma moderate reddish orange (171A). Fls 45mm wide, slightly scented; tepals 25 × 12mm, margins smooth, tips strongly recurved. Lvs 120 × 30mm, whorled. Stems 1.1m, green, with up to 15 fls. Early.

The name means “sweet morning” in Latvian.

‘Salmon Star’ VII(a-b/b-c)

Inside of tepals milk-white, flushed pale salmon-orange beside midveins in central third; midveins soft salmon-orange, becoming a more apricot shade towards top; basal third a conspicuous narrow triangle of apple-green with a narrow margin of pale yellowish green. Midribs bright, pale yellowish green. Spots salmon-pink, relatively large, round to elliptic, evenly and quite densely distributed over basal two-thirds of each tepal; papillae

sometimes present, pale creamy yellow; filaments slender, greenish white; pollen soft orange; style dull apple-green; stigma greenish white. Tepal margins more or less smooth, inside tepals much broader and less strongly recurved than outer tepals. Stems 0.9m. Late. Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

It is not possible to register this cultivar epithet as it is too similar to, and likely to cause confusion with, ‘Salmonstar’, which was registered for another lily by B. Mićulka in 1990 (ICNCP, 2004: Art. 19.25).

‘Samarinda’ VIII(a/b) LA

Seedling BW 98-09 × seedling RS 01-18

H: Vletter en Den Haan Beheer BV, pre-2003

N: Vletter en Den Haan Beheer BV, 2003

REG: Vletter en Den Haan Beheer BV, 2008

Tepals yellowish white (155D); throat light yellow-green (142D). Outside of outer tepals with strong red (53B) stripe at base; midribs strong yellow-green (base 144C, top 144B). Outside of inner tepals strong yellow-green (144C) at base; midribs tinged strong red (53B). Spots and papillae absent; nectaries strong yellow-green (144A); pollen moderate reddish brown (175A); stigma light yellow-green (150D). Fls 190mm wide, scented; tepals 115 × 62mm, margins smooth, tips slightly recurved. Lvs 150 × 20mm. Stems 1.3m, green with darker markings, with up to 6 fls. Named after a city on Borneo.

‘Santa Rosa’ VIII(b/a) LO

longiflorum × Div. VIII (Other hybrids) LO hybrid

H: Testcentrum voor Siergewassen BV, pre-2008

REG: Testcentrum voor Siergewassen BV, 2008

Inside of tepals flushed strong purplish pink (73B) to light purplish pink (73C) on a white ground, paling to white towards tips; midveins green at base. Outside of inner tepals greenish white (155C) flushed pale pink. Spots and papillae absent; nectaries green; pollen pale brown; stigma pale green. Tepals long to very long, and broad, margins slightly ruffled, tips recurved. Lvs of medium length to long, and of medium width to broad. Stems 1.35m, green with darker spots and stripes, with up to 3 fls. Mid-season to late.

‘Sarabanda’ VIII(a/b) LA

Seedling BW 97-61 × seedling AW 96-04

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Tepals yellowish white (155D); midribs of outer tepals moderate yellow-green (143D) at base, strong yellow-green (144C) at top; midribs of inner tepals light yellow-green (150D); throat light yellow-green (145C). Spots and papillae absent; nectaries strong yellow-green (143B); pollen brownish orange (N170A); stigma brilliant yellow-green (150B). Fls 195mm wide, slightly scented; tepals 112 × 50mm, margins smooth, tips recurved. Lvs 113 × 26mm. Stems 1.2m, green, with up to 11 fls.

‘Schneesturm’ I(a/c)

Seedling NN W 7.7 × ‘Berliner Morgen’

H: G. Steinbrück, 1992 **G:** G. Steinbrück, 1995

N: G. Steinbrück, 2009 **I:** G. Steinbrück, 2009

REG: G. Steinbrück, 2009

Tepals pale yellow-green (155A); throat vivid white flushed pale yellow-orange. Spots numerous, small, orange, over basal two-thirds of each tepal; nectaries creamy white; pollen red-brown; stigma beige. Fls 100mm wide, not scented; tepals 50 × 22mm, margins smooth, tips strongly recurved. Lvs 55 × 12mm, scattered. Stems 0.7m, pale green, with up to 10 fls. Late June.

‘Schweriner Glocke’ I(c/b)

Seedling Fe 4 × seedling C 7.44

H: G. Steinbrück, 1992 **G:** G. Steinbrück, 1995

N: G. Steinbrück, 2009 **I:** G. Steinbrück, 2009

REG: G. Steinbrück, 2009

Tepals light yellowish pink (159A); midribs pale brownish beige, darker towards base; throat pale yellowish white. Spots small, relatively few, dark red-brown, evenly scattered over basal quarter of each tepal; nectaries pale green; pollen blackish brown; stigma very long, blackish violet. Fls 100mm wide, not scented; tepals 62 × 32mm, margins smooth, tips recurved to strongly recurved. Lvs 70 × 11mm, scattered. Stems 1m, brown, with up to 8 fls. Late June. The name means Schwerin Bell in German, and commemorates the flower show held in 2009 in Schwerin, the capital of the state Mecklenburg-Vorpommern.

‘Schweriner Morgen’ I(a/b-c)

Seedling LV 15 × ‘Beppo’

H: G. Steinbrück, 1992 **G:** G. Steinbrück, 1995

N: G. Steinbrück, 2009 **I:** G. Steinbrück, 2009

REG: G. Steinbrück, 2009

Tepals, including throat, strong red (53D). Spots few, inconspicuous, dark violet, over three-quarters of each tepal; nectaries bright red; pollen dark brown; stigma violet. Fls widely expanded, star-shaped, 140mm wide, not scented; tepals 50 × 8mm, margins smooth, tips slightly recurved. Lvs 65 × 12mm, scattered. Stems 0.7m, mid-green, with up to 8 fls. Late June. The name means Schwerin Morn in German, and commemorates the flower show held in 2009 in Schwerin, the capital of the state Mecklenburg-Vorpommern.

‘Schweriner Schloß’ I(a/c)

‘Skywatcher’ × ‘Vater Zille’

H: G. Steinbrück, 1992 **G:** G. Steinbrück, 1995

N: G. Steinbrück, 2009 **I:** G. Steinbrück, 2009

REG: G. Steinbrück, 2009

Inside of tepals vivid yellow (9A), shading to yellowish white (158D) at tips; midveins slightly raised, darker yellow. Outside of tepals yellow. Spots small, russet-brown, evenly scattered over basal third to half of each tepal; nectaries pale green and cream; pollen golden yellow; stigma pale brown. Fls 110mm wide, not scented; tepals 60 × 30mm, margins smooth, tips slightly recurved. Lvs 70 × 15mm, scattered. Stems 0.7m, mid-green, with up to 8 fls. Mid-July. The name means Schwerin Castle in German, the seat of the state parliament of Mecklenburg-Vorpommern, and commemorates the flower show held in Schwerin in 2009.

‘Seguret’ VIII(a/b) LA

Seedling BW 96-03 × seedling W 93-04

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Tepals, including throat, yellowish white (155D); outside of outer tepals flushed strong red (51A) on one side of midrib; midribs of outer tepals light yellow-green (144D); midribs of inner tepals strong yellow-green (144C) at base shading to strong greenish yellow (151A). Spots and papillae absent; nectaries strong yellow-green (143B); pollen moderate orange (167A); stigma light yellow-green (154D). Fls 195mm wide; tepals 120 × 55mm, margins smooth, tips slightly recurved. Lvs 140 × 18mm. Stems 1.1m, green, with up to 7 fls.

Named after one of the official “most beautiful villages in France”, in Provence.

‘Sensual’ VII (a/b)

Div. VII (Oriental hybrid) seedling × Div. VII seedling

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies

Holland BV, 2006 **REG:** De Jong Lelies Holland BV,

2008

Inside of tepals strong purplish pink (63C), with large patch of yellowish white (155D) above nectaries; margins and throat yellowish white (155D). Outside of outer tepals strong purplish pink (63C), yellowish white (155D) beside midribs at base; midribs strong yellow-green (144C) at base and top, strong purplish red (63B) between. Outside of inner tepals strong purplish pink (63C) mixed with yellowish white (155D), becoming just 155D beside midribs at base; midribs light yellow-green (144D) at base shading to strong purplish red (63B). Spots and papillae yellowish white (155D); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma deep purple (79C). Fls 230mm wide, scented; tepals 135 × 75mm, margins smooth, tips recurved. Lvs 190 × 47mm. Stems 1.17m, green, with up to 3 fls.

‘Shanti’ VII (a/b)

Div. VII (Oriental hybrid) seedling × Div. VII seedling

H: De Jong Beheer BV, pre-2005 **N:** De Jong Lelies

Holland BV, 2005 **REG:** De Jong Lelies Holland BV,

2008

Inside of tepals vivid purplish red (61C); midveins vivid red (52A) tinged yellowish white (155D) at top; throat yellowish white (155D). Outside of outer tepals vivid purplish red (61C), shading to strong purplish red (64C) towards margins and tinged yellowish white (155D) beside midribs at base; midribs strong yellow-green (143C) at base shading to strong yellow (153D). Outside of inner tepals deep purplish pink (64D), shading to very pale purple (69C) beside midribs at base; midribs strong yellow-green (143C) at base shading to strong yellow (153D) towards top. Spots and papillae strong purplish red (60B); nectaries strong yellow (N144B) edged brilliant greenish yellow (5A); pollen vivid yellowish pink (28A); stigma pale yellow-green (193C). Fls 250mm wide, scented; tepals 155 × 75mm, margins smooth, tips slightly recurved. Lvs 190 × 45mm. Stems 0.95m, green with darker markings, especially at base, with up to 3 fls.

SIGNUM VII(a/b)

Trade designation of 'Zanlorsig'

'Siracl' I(a/b)

'Holebaba' × 'Latvia'

H: M. Tlach, pre-2007 **N:** M. Tlach, pre-2007

I: M. Tlach, 2007 **REG:** M. Tlach, 2007

Tepals sulphur-yellow, margins closely spotted violet; throat green-yellow. Midribs brown-green. Nectaries pale green, frosted white; pollen orange-brown; stigma green-yellow. Fls 130mm wide; tepals 80 × 30mm, tips slightly recurved to recurved; pedicels 75mm, stout, brownish. Lvs 130 × 20mm, glossy, pale green. Stems 0.85m, stout and stiff, green; infl. long. Late June.

'Slovačka' VI(b-c/a-b)

Syn. of 'Bonviván'

'Smoky Lemon' VI(b-c/a)

Herald Angel Group × 'Damsón'

H: B. Strohman, 1999 **G:** L.C. Bligh, 2003

N: L.C. Bligh, 2007 **REG:** L.C. Bligh, 2008

Inside of tepals light greenish yellow (1C), tinged deep purplish pink (70C), margins white in places; throat brilliant yellow (7A). Outside of outer tepals heavily mottled deep purplish pink (70C) over a light greenish yellow ground, shading to pale yellow-green (155A) towards margins. Outside of inner tepals light greenish yellow (1C) tinged deep purplish pink (70C); midribs deep purplish pink (70C). Spots absent, nectaries lemon-yellow; pollen golden orange; stigma red-purple. Buds deep purplish pink. Fls 150mm wide, strongly scented; tepals 145 × 55mm, margins smooth, tips slightly recurved. Lvs 180 × 9mm, scattered. Stems 1.4m, maroon, flecked green, with up to 17 fls in pyramidal infl. Late December (Tasmania).

'Sniegbaltitis' II(c/d)

Un-named seedling × unknown

H: D. Hercbergs, pre-2003 **G:** D. Hercbergs, 2003

N: D. Hercbergs, 2005 **I:** D. Hercbergs, 2007

REG: D. Hercbergs, 2008

Tepals pinkish white (N155B); midveins and midribs green at base. Spots absent; nectaries pale green; pollen vivid orange-yellow (21A); stigma pale green. Fls 50mm wide, slightly scented; tepals 40 × 16mm, margins smooth, tips strongly recurved. Lvs 130 × 50mm, whorled. Stems 1.1m, green, with up to 18 fls. Late. The name means "snow-white boy" in Latvian.

'Snow Prince' VII(a/b)

Parentage unknown

H: E.T. Kirsch, pre-1986 **G:** E.T. Kirsch, pre-1986

I: Melridge Inc., c.1988

Tepals white, sometimes tinged pink. Spots absent; papillae numerous, white, in central 120mm of fl.; nectaries green; filaments 70–80mm, white to pale green; anthers 28mm, dark reddish orange (172B); style pale green to white, sometimes tinged pink; stigma pale purple-grey. Buds 120–150mm long, white flushed pale green or cream, midribs flushed pale pink in basal half, pale green in upper; peduncles 10–20mm, dark green. Fls 200–220mm wide, strongly scented; tepals ovate, inner 60mm wide, outer 30mm wide,

tips slightly recurved; pedicels 60–80mm, dark green. Lvs 50–100 × 20mm, leathery, glossy, dark green to plum-coloured. Stems to 1m, plum-coloured, with many fls in a compact raceme. Mid-season.

'Solomon' VII(a/-)

Parentage unknown

H: World Breeding BV, 2003 **N:** Liliande BV, 2003

REG: World Breeding BV, 2008

Tepals greenish white (155C); inside of tepals brilliant yellow (8A) to pale yellow (8D) at base. Spots yellow; papillae present; nectaries yellow-green; pollen red-brown; stigma purple. Tepals long and of medium width to broad, margins ruffled, tips recurved. Lvs of medium length to long, and broad. Stems 1.4m, green, with 1 or 2 fls. Mid-season to late.

'Specto' VII(b-a/c)

Div. VII (Oriental hybrid) seedling × Div. VII seedling

H: De Jong Beheer BV, pre-2008 **N:** De Jong Lelies

Holland BV, 2008 **REG:** De Jong Lelies Holland BV,

2008

Inside of tepals whiter than white (NN155D), brilliant greenish yellow (6B) above and beside nectaries; throat white (NN155D). Outside of outer tepals white (NN155D), tinged strong purplish red (60B) beside midribs at base; midribs strong yellow-green (base 144B flushed strong purplish red (60B); top 144A, 144C between). Outside of inner tepals white (NN155D); midribs dark red (59A) at base shading to light greenish yellow (3D). Spots white (NN155D); papillae brilliant greenish yellow (6B); nectaries strong yellow-green (N144D); pollen strong reddish orange (169A); stigma light purplish grey (N187D). Fls 270mm wide, scented; tepals 130 × 78mm, margins smooth, tips slightly recurved. Lvs 200 × 47mm, dark green. Stems 1.2m, green, with up to 4 fls.

'Starburst' (b/b-c)

I: H.W. Hyde and Son, 2009

Inside of tepals white with a conspicuous flare along and beside midveins in basal two-thirds; strong lemon-yellow in basal third; bright raspberry-red in central third; tips white. Spots carmine-pink, evenly distributed over basal three-fifths of each tepal, from near margins to beside, rarely on, flare; spots towards margins small and round, becoming progressively larger towards flare, sometimes coalescing to form short dashes. Papillae few, greenish white; nectary furrows rich apple-green; filaments and style paler apple-green; pollen reddish orange; stigma dark purplish brown. Tepal margins ruffled, tips recurved. Exhibited by H.W. Hyde and Son at RHS Chelsea Flower Show, 2009. It is not possible to register this cultivar epithet as it is in use already for another lily (ICNCP, 2004: Art. 27.1).

'Starburst' (Tango) I(a-b/b-c)

Tepals bright rich red in upper half. Outside of tepals stained dark dull brown in basal half. Spots coalescing to form a solid patch of dark purple-black in basal half of each tepal, becoming progressively less dense towards margins and beyond upper edge of patch, producing a picotee or stippling effect; anthers bright

red; filaments and style pale coral-red. Buds orange-red, tips green. Fls large, long lasting; tepals margins smooth, tapering quite abruptly to slightly recurved tips. Stems 0.8–1.2m. Early summer to early autumn. It is not possible to register this cultivar epithet as it is in use for another lily which was exhibited by H.W. Hyde and Son at RHS Chelsea Flower Show, 2009 (ICNCP, 2004: Art. 27.1).

‘Starlight Express’ VII (a-b/b-c)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** W. Vink, 2003 **REG:** “The Originals” U.A., Coöp.

Kwekersvereniging, 2007

Inside of tepals deep red (60A) to strong purplish red (60B), shading to pale yellow-green (155A) towards margins and tips. Outside of tepals strong purplish red (60C/D). Spots and papillae present; nectaries yellow-green; pollen red-brown; stigma dark purple. Fls 200mm wide; tepals of short to medium length and of medium width, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad to very broad. Stems 0.5m, green with some darker markings, with up to 5 fls. Mid-season to late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

‘Starshell’ VII (a-b/b)

Parentage unknown

H: E.T. Kirsch, pre-1986 **G:** E.T. Kirsch, 1986

I: Melridge Inc., c.1988

Inside of tepals moderate purplish red (58A) to strong purplish red (58B) (occasionally vivid purplish red (57B)) at centre, shading to vivid purplish red (57D) towards margins; margins white, 5–6mm wide on inner tepals, 2–4mm on outer. Papillae conspicuous, moderate purplish red (58A) to strong purplish red (58B), over central 120mm of each tepal; nectaries green, with yellow margin 2–4mm wide; filaments 70–80mm, pale green; anthers strong reddish orange (169A) to strong orange (169B); style pale green; stigma pale to mid- purple-grey. Buds 100–120mm long, pale pink, tips moderate purplish red (59C) to strong purplish red (59D); peduncles 20–30mm, dark green. Fls 200–220mm wide, strongly scented; tepals ovate, inner 50mm wide, outer 30mm wide, margins (very) slightly ruffled, tips slightly recurved; pedicels 70–80mm, dark green. Lvs 50–100 × 20mm, leathery, glossy, mid-green. Stems 0.65–1m, with up to 8 fls; infl. a raceme. Mid-season.

‘Stowport Buccaneer’ VII(b/b)

‘Casa Blanca’ × *L. × parkmanii* ‘Red Sun’ hybrid

G: L.C. Bligh, 2000 **S:** L.C. Bligh, c.2006

N: L.C. Bligh, 2006 **REG:** L.C. Bligh, 2009

Tepals and throat white. Spots of medium size, round to elliptic, strong purplish red (72A), scattered fairly densely over basal half to two-thirds of each tepal; nectary furrows pale green; pollen golden brown; stigma brown. Fls 175mm wide, slightly scented; tepals 118 × 54mm, margins slightly ruffled, tips slightly recurved. Lvs 157 × 46mm, scattered. Stems 1.5m, green, with up to 8 fls. Early February (Tasmania).

‘Stowport Optimum’ VII (b/b)

‘Stowport Buccaneer’ × ‘Stowport Lolita’

H: L.C. Bligh, 2002 **G:** L.C. Bligh, 2006

N: L.C. Bligh, 2008 **REG:** L.C. Bligh, 2008

Inside of tepals, including throat, strong purplish pink (68B). Outside of tepals pale purplish pink (65D). Spots numerous, round to elliptic, of medium size to small, strong purplish red (60B), densely and fairly evenly scattered over each tepal; papillae few, strong purplish red (60B), near nectaries; nectaries green; pollen chocolate-brown; stigma dull green. Fls 160mm wide, slightly scented; tepals 100 × 59mm, margins smooth, tips slightly recurved, those of outer tepals twisted. Lvs 200 × 45mm, scattered. Stems 1.65m, green, with up to 30 fls. Mid-February (Tasmania).

‘Stowport Sunrise’ VI(b-c/b)

Herald Angel Group × ‘Damsen’

H: B. Strohmman, 1999 **G:** L.C. Bligh, 2003

N: L.C. Bligh, 2007 **REG:** L.C. Bligh, 2008

Inside of tepals vivid yellow (9A), shading to paler 9B, some moderate red (183C) shading showing through towards margins from outer face; throat brilliant yellow (9C). Outside of tepals brilliant yellow (9C), shading to moderate red (183C) especially beside and along midribs. Spots and papillae absent; nectaries bright green; pollen golden yellow; stigma red-purple. Fls 150mm wide, slightly scented; tepals 115 × 62mm, margins slightly ruffled, tips very slightly recurved. Lvs 115 × 12mm, scattered. Stems 1.25m, green, flecked maroon, with up to 8 fls. January (Tasmania).

‘Strawberry and Cream’ I(a-b/b-c)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2005 **N:** “The Originals” U.A., Coöp. Kwekersvereniging, 2005

REG: Mak Breeding Rights BV, 2008

Inside of tepals, including throat, pale purplish pink (65C) to greenish white (155C); basal half overlaid with a broad patch of deep purplish red (59B) to moderate purplish red (59C), upper edge disintegrating into numerous spots. Outside of tepals pale purplish pink (65C) to greenish white (155C), with dark spots. Spots darker than deep purplish red (59B); papillae present; nectaries green; pollen orange-brown; stigma purple-red. Fls 150mm wide; tepals short and narrow, margins slightly ruffled, tips slightly recurved. Lvs of medium length, and of narrow to medium width. Stems 0.8–1.35m, green, with up to 10 fls. Late. Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Sulpice’ VIII(a-b-c) LA

Seedling AM 96-01 × seedling BM 496-18

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals deep red (60A) to strong purplish red (60B), shading to deep red (60A) towards margins and along midveins; throat light yellow-green (150D). Inside of outer tepals strong purplish red (60B/C). Outside of tepals strong purplish red (60B), shading to deep red (60A) towards margins; midribs of outer tepals strong yellow-green at base (143C) and top

(144B), deep red (60A) between; midribs of inner tepals strong yellow-green (144C) at base shading to deep red (60A). Spots and papillae absent; nectaries strong yellow-green (143A); pollen greyish brown (166A); stigma light yellow-green (154D). Fls 210mm wide; tepals 130 × 60mm, margins smooth, tips recurved. Lvs 150 × 30mm. Stems 1.3m, green, with up to 5 fls.

‘Summer’s Beauty’ VIII(b/a) OT

Un-named, pink-flowered (a/a) seedling × (‘Starburst Sensation’ × (*L. × kewense* ‘White Henryi’ × ‘Dominique’))

H: R.H. Bazett, 1997 **G:** R.H. Bazett, 2000

N: R.H. Bazett, 2007 **REG:** R.H. Bazett, 2007

Inside of inner tepals white, overlaid in lower half above nectary furrows by arc of purple-red, a more solid and darker colour extending further along midveins; throat pale yellow. Outside of tepals white, pale purple-red showing through from inner face; midribs greyed green. Spots and papillae absent; nectary furrows deep, moss green, conspicuously edged golden yellow; pollen dark brown; stigma pale grey-green. Fls 155mm wide, strongly scented; tepals 116 × 49mm, margins smooth, tips strongly recurved. Lvs 160 × 24mm, scattered. Stems 1.9m, dark green, with up to 25 fls; infl. a raceme. July. Triploid.

‘Sunny Bonaire’ VII(a/b)

Parentage unknown

H: Mak Breeding BV, pre-2007 **N:** Mak Breeding BV, 2007 **REG:** Mak Breeding Rights BV, 2008

Inside of tepals light purplish pink (65B), paler towards base; midveins purplish pink at base. Outside of tepals pale purplish pink (65C). Spots darker than tepal colour; papillae present; nectaries green; pollen orange-brown; stigma purple. Tepals of short to medium length and of narrow to medium width, margins strongly ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad. Stems 0.55m, green, with up to 7 fls. Late.

‘Sunny Borneo’ VII(a-b/b-c)

Parentage unknown

H: Mak Breeding BV, pre-2007 **N:** Mak Breeding BV, 2007 **REG:** Mak Breeding Rights BV, 2009

Inside of tepals strong purplish pink (73B) on a paler ground, shading to white (NN155C) in basal portion. Outside of inner tepals predominantly very pale purple (73D). Spots absent; papillae white; nectaries green; pollen orange-brown; stigma purple-red. Fls 150–200mm wide; tepals short and of medium width, margins slightly ruffled, tips slightly recurved. Lvs of medium length to long, and (very) broad, dark green. Stems 0.45m, green with some darker spots and stripes, with up to 8 fls. Mid-season to late.

‘Sunny Star’ VIII(a/b) LA

Derived from ‘High Class’, *longiflorum* ‘Avita’ and ‘Shiraz’

H: Testcentrum voor Siergewassen BV, pre-2007

N: Testcentrum voor Siergewassen BV, 2007

REG: Testcentrum voor Siergewassen BV, 2008

Inside of tepals strong purplish red (60B/C), more blue with age; midveins deep red (60A) at base.

Outside of inner tepals strong purplish red (60C).

Spots absent; papillae inconspicuous; nectaries green; pollen yellow-brown; stigma orange-yellow. Tepals short and of narrow to medium width, margins (very) slightly ruffled, tips (very) slightly recurved. Lvs of medium length, and of narrow to medium width.

Stems 1.4m, green with darker spots and stripes, with up to 7 fls. Mid-season to late.

‘Sunny Sulawesi’ VII(a-b/b-c)

Parentage unknown

H: Mak Breeding BV, pre-2007 **N:** Mak Breeding BV, 2007 **REG:** Mak Breeding Rights BV, 2009

Inside of tepals white (NN155C); outside of inner tepals predominantly NN155C. Spots absent; papillae present; nectaries green; pollen orange-brown; stigma dark purple. Tepals short and of medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of short to medium length, and broad, dark green. Stems 0.4–0.45m, green, with up to 5 fls. Mid-season.

‘Sunshine’ (Taeon) I(a/b)

‘Holebibi’ × ‘Shocking Pink’

H: Taeon Lily Experiment Station, 1998

G: Taeon Lily Experiment Station, 2000

N: Taeon Lily Experiment Station, pre-2006

I: Taeon Lily Experiment Station, c.2006

Inside of tepals light yellow-orange; outside of tepals darker, vivid orange. Spots absent; pollen and stigma reddish brown. Fls 133mm wide; tepal margins smooth, tips only very slightly recurved. Lvs 110 × 25mm. Stems 0.925m, light brown, with c.3 fls in a racemose infl. Early June (under glass).

Epithet published in Abstract S05-P-31 from 27th International Horticultural Congress and Exhibition, Korea, 2006.

It is not possible to register this cultivar epithet as it is in use for another lily, bred by J. de Graaff c.1961 (ICNCP, 2004: Art. 27.1).

‘Suntrip’ VIII(a-b/b) LA

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2007

N: Testcentrum voor Siergewassen BV, 2007

REG: Testcentrum voor Siergewassen BV, 2008

Tepals strong orange, inside 25A; outside of inner tepals N25B/C; midveins N25A at base. Spots present, especially along margins of each tepal towards base; papillae present; nectaries orange; pollen red-brown; stigma orange-red. Tepals of short to medium length and of medium width, margins (very) slightly ruffled, tips (very) slightly recurved. Lvs of medium length to long, and of narrow to medium width. Stems 1.25m, green with faint darker spots and stripes, with up to 5 fls. Early to mid-season.

‘Suspension’ VII(a/b)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006

N: Testcentrum voor Siergewassen BV, 2006

REG: Testcentrum voor Siergewassen BV, 2007

Inside of tepals strong purplish red (61B), shading to greenish white (155C) towards margins and tips; midveins deep red (60A) at base. Outside of inner

tepals deep purplish pink (N66D). Spots fairly numerous to numerous; papillae present; nectaries yellow-green; pollen brown; stigma green, with small purple spots. Tepals of medium length and width, margins slightly ruffled to ruffled, tips recurved. Lvs of medium length to long, and broad. Stems of average height to tall, green with faint darker spots and stripes, with average no. of fls. Mid-season to late. Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

‘Sutherland Belle’ I(b-c/d)

‘Sutherland Flamethrower’ × Tiger Babies Group
H: D.J. Hay, 2000 **G:** D.J. Hay, 2005 **N:** D.J. Hay, 2008
REG: D.J. Hay, 2008

Tepals pink; inside of tepals gold and cream at base; throat dark pink; midribs pale green. Spots dark purple, numerous; nectaries pink; pollen dark rust-orange; stigma pink. Fls 125mm wide, not scented; tepals 85 × 25mm, margins slightly ruffled, tips strongly recurved. Lvs 85 × 25mm, scattered. Stems 0.76m, robust, greenish brown, with up to 8 fls. July.

‘Sutherland Citrus Melody’ I(b/c)

‘Catherine Laburi’ × ‘Sutherland Flamethrower’
H: D.J. Hay, 2001 **G:** D.J. Hay, 2004 **N:** D.J. Hay, 2007
REG: D.J. Hay, 2008

Inside of tepals soft lemon-yellow, basal half flushed vivid orange. Spots of medium size, dark red, evenly distributed, mostly over orange flush, avoiding margins; papillae absent; nectary furrows white; pollen rust-orange; stigma yellow. Fls 155mm wide, not scented; tepals 72 × 36mm, margins smooth, tips slightly recurved. Lvs 48 × 18mm, scattered. Stems 0.69m, brown, with up to 6 fls. July.

‘Sviesta Sāpnis’ II(c/d)

Un-named seedling × unknown
H: D. Hercbergs, pre-2003 **G:** D. Hercbergs, 2003
N: D. Hercbergs, 2004 **I:** D. Hercbergs, 2007
REG: D. Hercbergs, 2008

Inside of tepals brilliant greenish yellow (5B), shading to pale yellow (11D) towards tips. Outside of tepals light greenish yellow (5C); midribs pale yellow (11D). Spots and papillae absent; pollen vivid orange-yellow (21A); stigma green. Fls 55mm wide, slightly scented; tepals 30 × 14mm, margins smooth, tips strongly recurved. Lvs 130 × 28mm, whorled. Stems 1.2m, green, with up to 25 fls. Mid-season. The name means “butter’s dream” in Latvian.

‘Tabledance’ VIII(a/b-c) OT

H: Mak Breeding BV, pre-2008
 Buds and tepals lilac-pink, inside of tepals with a small, circular patch of white at base; nectary furrows pale green. Spots and papillae absent. Tepal margins smooth, tips strongly recurved. Stems 1.2m. Late. Published in *The Lily Nook cat.*, 2009

‘Take Five’ VII (a/b)

Parentage unknown
H: Vletter en Den Haan Beheer BV, pre-2003
N: Vletter en Den Haan Beheer BV, 2003
REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals dark red (187C), inside of outer tepals deep red (60A), all with patch of light greenish yellow (8C) above nectaries; midveins strong purplish red (60B); throat yellowish white (155D). Outside of outer tepals deep purplish red (187D), tinged yellowish white (155D) beside midribs at base; midribs strong yellow-green (144B) at base, moderate olive-green (147A) at top, moderate red (47A) between. Outside of inner tepals deep purplish red (61A), shading to strong purplish red (60D) beside midribs, tinged yellowish white (155D) at base; midribs strong yellow-green (144C) at base shading to strong purplish red (61B). Spots and papillae dark red (59A); nectaries strong yellow-green (144B); pollen moderate reddish orange (171A); stigma moderate purplish red (70A). Fls 220mm wide; tepals 140 × 70mm, margins smooth, tips slightly recurved. Lvs 180 × 45mm. Stems 1m, green with darker markings, with up to 3 fls.

‘Tambor’ VI(b-c/b-d)

(‘Honeydew’ × ‘Gold Eagle’) × (((*henryi* × ‘Gold Eagle’) × ‘Toni’) × *L. × kewense* ‘White Henryi’)
H: B. Mićulka, pre-2007 **N:** B. Mićulka, pre-2007
I: B. Mićulka, 2007 **REG:** B. Mićulka, 2007
 Inside of tepals lemon-yellow; tips paler; throat dusted with linear trails of cinnamon-brown. Midribs pale green. Nectaries green; pollen apricot-brown; stigma pale green. Fls 135mm wide; tepals 120 × 47mm, margins slightly ruffled, tips recurved; pedicels 75mm, with numerous dark brown spots. Lvs 125 × 11mm, dull yellowish green, tips strongly recurved. Stems 1.15m, stiff, green, with faint reddish mottling; infl. short. Mid-June.

‘Tamburo’ VIII(a-b/b) LA

Seedling AM 97-14 × seedling BM 497-22
H: Vletter en Den Haan Beheer BV, pre-2005
N: Vletter en Den Haan Beheer BV, 2005
REG: Vletter en Den Haan Beheer BV, 2008
 Inside of inner tepals strong red (53D) to strong purplish red (54A), with patch of pale purplish pink (56D) above nectaries; margins deep red (53A); throat strong yellow-green (144C). Inside of outer tepals strong purplish red (54A), with patch of pale purplish pink (56D) above nectaries; margins strong red (53B). Outside of tepals strong purplish red (59D), tinged yellowish white (155D) beside midribs at base; margins strong red (46A); midribs of outer tepals light yellow-green (base 145B, top 145C); midribs of inner tepals tinged strong yellow-green (N144C) at base. Spots strong red (53C); papillae absent; nectaries strong yellow-green (144A); pollen moderate reddish brown (175A) to dark reddish orange (175B); stigma light yellow-green (145C). Buds a little pubescent. Fls 195mm wide; tepals 120 × 70mm, margins smooth, tips straight. Lvs 155 × 24mm. Stems 1.3–1.4m, dark with green markings, with up to 5 fls. Published in *The Lily Nook cat.*, 2009
 External images: *The Lily Nook cat.*, 2009
 The Italian word for “drum”.

‘Tanita’ VIII(b-c/a) LO

Tetraploid *longiflorum* ‘White Fox’ × tetraploid (*longiflorum* ‘Lorina’ × ‘Touch’)

H: Testcentrum voor Siergewassen BV, pre-2004
N: Marklily BV, 2004 **REG:** Testcentrum voor Siergewassen BV, 2008

Inside of tepals and outside of inner tepals greenish white (155C); midveins yellow-green at base. Spots and papillae absent; nectaries green; pollen orange-brown; stigma purple. Tepals long to very long and of medium width to broad, margins slightly ruffled, tips recurved. Lvs long and of medium width to broad. Stems 1.55m, green, with up to 3 fls. Late.

‘Tartuga’ VIII(a/b-a) LA

Seedling BG 96-15 × seedling AJ 98-20

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals brilliant yellow (7B); margins and midveins 7A; throat pale greenish yellow (2D). Outside of outer tepals pale yellow-green (4D); margins vivid yellow (9A); midribs strong yellow-green (base 143C, top 144C). Outside of inner tepals light greenish yellow (8C), shading to pale yellow-green (4D) beside midribs at base; margins vivid yellow (9B); midribs strong yellow-green (143C) at base shading to pale yellow-green (4D). Spots and papillae absent; nectaries strong yellow-green (143A); pollen moderate reddish brown (175A); stigma yellowish grey (156B) tinged deep purplish red (59B). Fls 170mm wide; tepals 135 × 70mm, margins smooth, tips slightly recurved. Lvs 160 × 28mm. Stems 1.3m, green with some darker markings, with up to 5 fls.

‘Taschkent’ VIII(b/b) LA

Parentage unknown

H: P. Wistuba, pre-2007 **N:** P. Wistuba, pre-2007

I: P. Wistuba, 2007 **REG:** P. Wistuba, 2007

Inside of tepals pale bronze-orange, margins dark brown at paler throat. Outside of tepals pale orange; midribs pink and yellow-green. Spots fairly large, numerous, dark crimson-brown, over one-quarter of each tepal; papillae absent; nectaries brown-green; pollen orange-brown; stigma pale orange. Fls 145mm wide; tepals 115 × 45mm, tips slightly recurved; pedicels 90mm, very stout, with numerous dark brown spots. Lvs 135 × 21mm, mid-green. Stems 0.85m, very stout, dark brown, with average no. of fls. Late June.

Tas Townships Group I(c/d)

(*davidii* × ‘Edith Cecilia’ hybrid) × yellow-flowered Div. I (b/-) (Asiatic hybrid) seedling bred by Henry Foster) × ‘Edith Cecilia’ hybrid

H: N. Jordan, 2003 **G:** N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2007

Inside of tepals from ivory-white (‘Lileah’, ‘Lowana’ and ‘Rinadeena’) to pale creamy yellow (‘Parrawe’); clear citron-yellow (‘Mengha’ and ‘Remine’) and citron-yellow flushed rose-pink (‘Luina’); soft apricot (‘Marrawah’) and richer apricot-yellow with peach-pink tips (‘Teepookana’), to richer orchid-pink at margins and tips (‘Nabageena’ and ‘Penghana’); through soft coppery pink (‘Mella’) and dark, rich

pink (‘Togari’) to carmine-red (‘Tullah’) with orange (‘Trowutta’), coral-red (‘Temma’) and fiery red (‘Mawbanna’ and ‘Redpa’). Small, mahogany-brown, “V”-shaped brushmarks sometimes present: most conspicuous in ‘Penghana’, ‘Rinadeena’ and ‘Trowutta’; spots numerous, particularly widespread in ‘Lowana’ and ‘Parrawe’, mahogany- or chocolate-brown. Tepal tips recurved (in ‘Parrawe’ and ‘Teepookana’ strongly so), margins smooth. Flowers single, not scented. Leaves numerous, scattered, 65–105 × 4–6mm, dark green. Stems 0.6–1.1m, green with brown flecks, or brown with green flecks; occasionally brown throughout (‘Mengha’ and ‘Redpa’); with 10–20 flowers in a pyramidal inflorescence. Mid-season: December (Tasmania). All cultivars in this Group bear names of western Tasmanian townships that are derived from Tasmanian Aboriginal words.

‘Teepookana’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005 **G:** N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007

REG: N. Jordan, 2007

Tepals, including throat, light orange-yellow (25D), shading to moderate pink (51D) in uppermost third. Spots of medium size to small, round to elliptic, mahogany-brown, fairly densely scattered over basal two-thirds of each tepal; nectaries rose-pink, frosted; pollen rich golden brown; stigma orange-brown. Fls 65mm wide, not scented; tepals 60 × 27mm, margins smooth, tips strongly recurved. Lvs 95 × 5mm, scattered. Stems 1m, brown, flecked green, with up to 18 fls. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal name for azure kingfisher (*Alcedo azurea*).

‘Temma’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007

I: Cam Lilies, 2007 **REG:** N. Jordan, 2007

Inside of tepals strong red (53B), especially towards tip, shading to vivid purplish red (57B) towards margins, flushed light orange (28D) especially towards base; throat deep pink (51B). Outside of tepals deep pink (51B) shading to light orange (28C). Small, V-shaped pencil-mark on each tepal and spots mahogany-brown; spots of medium size to small, round to elliptic, scattered over basal third to half of each tepal; nectaries pink, frosted; pollen pale golden brown; stigma pale orange. Fls 70mm wide, not scented; tepals 60 × 28mm, margins smooth, tips slightly recurved. Lvs 90 × 4mm, scattered. Stems 0.9m, green, flecked brown, with up to 18 fls in pyramidal infl. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “hut”.

‘Termoli’ VIII(b-a/b) LA

Seedling BW 96-47 × seedling AM 95-12

H: Vletter en Den Haan Beheer BV, pre-2005**N:** Vletter en Den Haan Beheer BV, 2005**REG:** Vletter en Den Haan Beheer BV, 2008

Inside of tepals light purplish pink (63D); midveins of inner tepals, and margins of outer tepals, strong purplish pink (63C); throat pale purplish pink (65D). Outside of outer tepals light purplish pink (65B); margins strong purplish pink (63C); midribs tinged moderate yellow-green (138C) at base. Outside of inner tepals moderate purplish pink (65A); midribs tinged strong yellow-green (143C) at base. Spots dark red (59A); papillae absent; nectaries moderate olive-green (137A); pollen vivid reddish orange (32A); stigma light orange-yellow (22D). Fls 210mm wide; tepals 120 × 58mm, margins smooth, tips slightly recurved. Lvs 160 × 30mm. Stems 1.25–1.3m, green, with up to 8 fls. A town on the Adriatic coast of Italy.

‘Tetra Black Beauty’ VIII(b-c/d)

‘Black Beauty’, registered by the I.C.R.A. in 1992, was originally classified in Div. VII (Oriental hybrids), but latterly is considered to be an Oriental/Trumpet (Orientalpet or O.T.) hybrid, and has, accordingly, been reclassified in Div. VIII (Other hybrids). Consequently, the tetraploid form of ‘Black Beauty’, ‘Tetra Black Beauty’, has been reclassified in Div. VIII also.

‘The Favourite’ II(c/d)

Un-named seedling × unknown

H: D. Herbergs, pre-2003 **G:** D. Herbergs, 2003**N:** D. Herbergs, 2004 **I:** D. Herbergs, 2007**REG:** D. Herbergs, 2008

Tepals greenish white (155C); throat green. Spots small, red-purple, on two-thirds of each tepal; nectaries green; pollen strong orange-yellow (17A); stigma pale greenish yellow (1D). Fls 60mm wide, slightly scented; tepals 40 × 20mm, margins smooth, tips strongly recurved. Lvs 130 × 40mm, whorled. Stems 1.4m, green, with up to 30 fls. Mid-season.

‘Tibo’ VIII(a/b) LA

Parentage unknown

H: J. Berbée, pre-2008 **N:** J. Berbée, 2008**REG:** J. Berbée, 2008

Tepals yellowish white (NN155A), shading to light yellow-green (154D) inside above nectaries; midribs of outer tepals strong yellow-green (143C at base and top, 144C between); midribs of inner tepals brilliant yellow-green (142B) at base shading to light yellow-green (145D) towards top; throat light yellow-green (145C). Spots and papillae absent; nectaries strong yellow-green (143A); pollen vivid orange-yellow (23A); stigma light yellow-green (145C). Fls 200mm wide; tepals 113 × 60mm, margins smooth to slightly ruffled, tips recurved. Lvs 150 × 20mm. Stems 1.1m, green with darker markings, with up to 7 fls.

‘Tiffany’ I(b/b-c)

Parentage unknown

H: Mak-Leek, Inc., pre-2009

Inside of tepals soft lilac-pink, golden yellow along and beside midveins in basal third; throat deep rose-pink.

Spots and papillae absent; filaments and pollen orange-red; stigma and style dark brown. Buds greenish flushed pink. Tepal margins very slightly ruffled, tips slightly recurved. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009

This cultivar epithet is not acceptable for registration as it was registered by Van der Salm Bulb Farm, Inc. in 1990 for another lily (ICNCP, 2004: Art. 27.1).

‘Tigerplay’ I(a-b/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003**N:** Q.J. Vink en Zn, 2003 **REG:** Mak Breeding

Rights BV, 2008

Inside of tepals vivid yellow (9A); margins and base dark red (187A/B). Outside of tepals brilliant yellow (8A) to light greenish yellow (8B), with spots darker than dark red (187A/B). Papillae present; nectaries yellow-green; pollen pale brown; stigma dark purple. Fls 180mm wide; tepals (very) short and of narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of short to medium length, and of narrow to medium width. Stems 0.8m, green, with average no. of fls. Early to mid-season.

Published in *The Lily Nook cat.*, 2009External images: *The Lily Nook cat.*, 2009**‘Timezone’ VIII(a/b) OT**

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2006**N:** Testcentrum voor Siergewassen BV, 2006**REG:** Testcentrum voor Siergewassen BV, 2008

Inside of tepals strong purplish red (64B/C), paler with age, shading to nearly white towards base; midveins yellow-green at base shading to white. Outside of inner tepals moderate purplish red (64A) to strong purplish red (64B). Spots and papillae absent; nectaries dark green; pollen reddish brown; stigma purple, with small spots. Fls 100–120mm wide; tepals of medium length to long and of narrow to medium width, margins (very) slightly ruffled, tips recurved. Lvs of medium length and breadth. Stems 1.4m, green with darker spots and stripes, with few fls. Late.

‘Tiny’ VII(b/b)

Parentage unknown

S: E.T. Kirsch, c.1979 **G:** E.T. Kirsch, 1979**I:** Melridge Inc., c.1988

Tepals pale purplish pink (62D), with 10mm-wide, moderate purplish pink (62B) band along midveins; fading browner with age. Papillae strong purplish red (63B); nectaries green, with 2–4mm-wide yellow margin; filaments 70–80mm, pale green; anthers vivid yellow (16A) to light orange-yellow (16B); style pale green to white, sometimes tinged pink; stigma pale purple-grey. Buds 100–120mm long, strong purplish pink (63C) to light purplish pink (63D), paler at tips; peduncles 20–30mm, dark green. Fls 180–200mm wide, scented; tepals ovate, inner 40mm wide, outer 25mm, margins of outer tepals slightly ruffled, tips of inner tepals slightly recurved; pedicels 50–100mm. Lvs 40–80 × 15–20mm, leathery, glossy, dark green. Stems 0.3–0.5m, with up to 8 fls in a compact raceme. Mid-season.

‘Tiny Padhye’ I(a-b/b-c)

Parentage unknown

H: Mak ‘t Zand BV, pre-2007 **N:** Mak ‘t Zand BV,

2007 **REG:** Mak ‘t Zand BV, 2008

Inside of tepals yellowish white (155D), shading to dark red (187A/B) along margins in upper half of tepal. Outside of inner tepals greenish white (155C). Spots few, darker than dark red (187A/B); papillae present; nectaries green; pollen orange-yellow; stigma pale yellow. Tepals (very) short and of narrow to medium width, margins slightly ruffled, tips slightly recurved. Lvs of short to medium length, and of narrow to medium width. Stems 0.45m, green, with up to 9 fls. Early to mid-season.

‘Togari’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 **S:** N. Jordan, c.2005

G: N. Jordan, 2005 **N:** N. Jordan, 2007 **I:** Cam Lilies, 2007 **REG:** N. Jordan, 2008

Inside of tepals moderate purplish red (185C) shading to pale orange-yellow (159B) in basal third; throat pale orange-yellow (159B). Outside of tepals moderate purplish red (185C) shading to moderate purplish pink (186D). Spots of medium size, round, chocolate-brown, scattered evenly over basal half of each tepal; papillae few, pale grey-purple, between throat and spots; nectaries pink, frosted; pollen bright orange; stigma pale orange. Fls 70mm wide, not scented; tepals 60 × 23mm, margins smooth, tips slightly recurved. Lvs 75 × 3mm, scattered. Stems 0.85m, green, flecked brown, with up to 12 fls. December (Tasmania).

Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “crown”.

‘Tome’

Unregistered seed parent of ‘Yujiao’. No further details known.

‘Topstar’ VII(a/b-c)

Parentage unknown

H: Mak ‘t Zand BV, pre-2005 **N:** Mak ‘t Zand BV,

2005 **REG:** Mak ‘t Zand BV, 2008

Inside of inner tepals deep purplish pink (N57C/D), paler and bluer with age; midveins yellow-green at base. Outside of inner tepals deep purplish pink (N57D). Spots and papillae absent; nectaries green, fading to yellow; pollen reddish brown; stigma dark purple. Tepals of medium length to long and of medium width, margins ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad. Stems 1.15m, green with faint darker spots and stripes, with up to 4 fls. Mid-season.

‘Torpedo’ I(a/b)

Mixture of Div. I (Asiatic hybrid) tetraploids × ‘Lenora’ un-named seedling

H: De Jong Beheer BV, pre-2006 **N:** De Jong Lelies Holland BV, 2006 **REG:** De Jong Lelies Holland BV, 2008

Inside of tepals vivid yellow (12A); throat light yellow-green (150D). Outside of tepals vivid yellow (9A), strong yellow-green (143C) at base; tips of outer

tepals vivid yellow (13A), tips of inner tepals strong yellow-green (143C) tinged vivid yellow (13A). Spots dark red (59A); papillae vivid yellow (12A); nectaries strong yellow-green (143A); pollen strong orange (N163B); stigma brilliant greenish yellow (1B). Fls 193mm wide; tepals 115 × 60mm, margins smooth, tips slightly recurved. Lvs 145 × 20mm. Stems 0.9m, green, with up to 6 fls.

‘Touch of Pink’ VIII(b/a) OT

Un-named Div. VII (Oriental hybrid) seedling × (‘Starburst Sensation’ × (*L. × kewense* ‘White Henryi’ × ‘Dominique’))

H: R.H. Bazett, 1988 **G:** R.H. Bazett, 1991 **N:** R.H. Bazett, 2007 **REG:** R.H. Bazett, 2007

Inside of tepals white, flushed pink over two-thirds of tepal but especially beside midveins; midveins yellow-green at base, shading to pink; throat pale yellow-green. Outside of tepals creamy white; midribs pale green. Spots and papillae absent; nectaries dark green; pollen dark brown; stigma pale grey-green. Fls 183mm wide, strongly scented; tepals 145 × 55mm, margins slightly ruffled, tips slightly recurved. Lvs 225 × 23mm, scattered. Stems 1.74m, dark green, with up to 24 fls; infl. racemose, with secondary buds. July. Triploid.

‘Trebisonda’ VIII(b-a/b) OT

Seedling PG 96-62 × seedling RH 97-09

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals brilliant yellow (inner 7B; outer 7A, shading to 8A towards margins); midveins of inner tepals 7A; throat light greenish yellow (6D). Outside of outer tepals pale greenish yellow (2D), shading to light greenish yellow (5C) towards margins and deep yellowish pink (39B) beside midribs; midribs light olive-brown (199B) at base, strong yellow-green (144C) at top, deep yellowish pink (39B) between. Outside of inner tepals light greenish yellow (8B), shading to pale yellow-green (4D) beside midribs; midribs at base light yellow-green (144D) with deep yellowish pink (39B) stripe, shading to light yellow-green (150D) towards top. Spots absent; papillae few, brilliant yellow (7A); nectaries strong yellow-green (144B); pollen moderate reddish brown (175A); stigma dark purplish red (N79A). Fls 230mm wide, scented; tepals 140 × 75mm, margins slightly ruffled, tips recurved. Lvs 235 × 35mm. Stems 1.2m, green with darker markings, with up to 4 fls.

‘Tropical Breeze’ I(a/b-c)

N: Mak-Leek, Inc., pre-2008

Inside of tepals greyish pale lilac-pink; tips yellowish green. Spots burgundy-red, variable in size, in basal third of each tepal coalescing as dashes beside midveins and more solid patches of colour towards margins, becoming progressively less densely distributed along margins towards tips; papillae absent; nectaries pale green; filaments whitish; anthers dull russet-brown; style pale yellowish green; style russet-brown. Outer tepals rather narrow elliptical, inner tepals larger, ovate, margins smooth,

tips becoming strongly recurved with age. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Tropical Dream’ I(a/-)

H: Van der Salm, pre-2008

Inside of tepals coral-pink; basal portion of each tepal warm flesh-pink/apricot. Fls borne on long pedicels. Stems 0.9m. Mid-season. Tetraploid.

Published in *The Lily Nook cat.*, 2009

‘Tropical Garden’ VI(b-c/b-c)

‘Marilyn’ × (‘Mimosa Star’ × ‘Caramba’)

H: A. Mego, pre-2008 G: A. Mego, 2008 N: A. Mego, 2008 I: A. Mego, 2008 REG: A. Mego, 2008

Tepals orange; outside of tepals paler; midribs pale green. Spots and dashes varying in size, dark red, raised, widely distributed over each tepal; nectaries pale green; filaments, style and stigma pale green; pollen brown. Fls 140mm wide; tepals 90 × 38mm, margins slightly ruffled, tips strongly recurved; pedicels 90mm, green slightly spotted brown; with secondary buds. Lvs 190 × 38mm, slightly glossy, dark green. Stems 1.35m, stout, with dark brown spots, with infl. of average length. Mid-July.

‘Tropic Delight’ I(a/-)

H: Van der Salm, pre-2008

Inside of tepals papaya-pink; a blend of dark pinkish red with a slight orange flush. Stems 0.9m. Mid-season. Tetraploid.

Published in *The Lily Nook cat.*, 2009

‘Trowutta’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 S: N. Jordan, c.2005

G: N. Jordan, 2005 N: N. Jordan, 2007 I: Cam Lilies, 2007 REG: N. Jordan, 2008

Inside of tepals moderate purplish red (185C) at tips, base and throat, flushed strong orange-yellow (163B) in central portion of each tepal. Outside of tepals moderate purplish red (185C), shading to dark pink (182D). Small, V-shaped pencil-mark on each tepal and spots mahogany-brown; spots of medium size, round to elliptic, scattered over basal half of each tepal; nectaries grey-purple, forming a conspicuous, six-pointed star; pollen orange-red; stigma pale orange. Fls 75mm wide, not scented; tepals 60 × 23mm, margins smooth, tips recurved. Lvs 70 × 4mm, scattered. Stems 0.8m, green, flecked brown, with up to 15 fls; infl. pyramidal. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “stone tool”.

‘Tullah’ I(c/d)

Selection from Tas Townships Group

H: N. Jordan, 2003 S: N. Jordan, c.2005

G: N. Jordan, 2005 N: N. Jordan, 2007 I: Cam Lilies, 2007 REG: N. Jordan, 2007

Inside of tepals dark red (59A), shading to pale greenish yellow (160C) in basal half; throat 160C. Outside of tepals dark red (59A) to deep purplish pink (70C). Spots and ill-defined, V-shaped pencil-mark

on each tepal, mahogany-brown; spots fairly large, round to elliptic, scattered fairly evenly over basal half of each tepal; nectaries pink, frosted; pollen golden brown; stigma pale green. Fls 75mm wide, not scented; tepals 60 × 28mm, margins smooth, tips slightly recurved. Lvs 95 × 4mm, scattered, dark green. Stems 1.05m, green, flecked brown, with up to 18 fls; infl. pyramidal. December (Tasmania). Named after a township in western Tasmania, derived from the Tasmanian Aboriginal word for “seashell”.

‘Tumalo’ VII(b-a/b)

Parentage unknown

S: E.T. Kirsch, c.1985 G: E.T. Kirsch, pre-1985

I: Melridge Inc., c.1988

Tepals light purplish pink (62C), shading to strong yellowish pink (38A) in 10–15mm-wide band along midveins/midribs. Papillae numerous, strong purplish red (63B); nectaries green, with 2–4mm, yellow margin; filaments 70–80mm, pale green; anthers strong brown (172A) to dark reddish orange (172B); style pale green; stigma pale purple-grey. Buds 100mm long; white to very pale pink; midribs tinged green at base, pale yellow-cream at tips; peduncles <10mm. Fls c.200mm wide, strongly scented; inner tepals broadly ovate, tips acuminate, 50mm wide; outer tepals 30mm wide; margins slightly ruffled, tips slightly recurved; pedicels 30–50mm, dark green, ascending. Lvs 50–100 × 20mm, leathery, glossy, dark green. Stems 0.4–0.5m, with up to 8 fls in a compact raceme. Mid-season (late July).

‘Tyrolia’ I(a/c)

Un-named, small-flowered, white-flowered, pollen-free sport of ‘Sunkissed’ O.P. × un-named, tall, exceptionally vigorous, large-flowered, orange-flowered seedling of ‘Chinook’ O.P.

H: H. Mantel, pre-1989 G: H. Mantel, 1989

I: Mount Hood Lilies, Inc., c.1995

Syn. ‘White Bouquet’

Tepals open creamy white, becoming pure white in upper part at least; inside occasionally tinged light greenish yellow (4C) to pale yellow-green (4D) above nectaries. Spots occasional, 2–4, deep magenta, at base of inner tepals; nectaries pale green, shading to ivory white, inconspicuous, with thin down of short white hairs; filaments 20–30mm long, cream-white; anthers usually absent, but occasionally 1–4mm, indehiscent; pollen absent; style 40–90mm long, cream-white to pale yellowish pink (27D) or paler; stigma moderate purplish red (185C) to deep pink (185D). Buds 70–100mm long; cream to white, green at tips; midribs green; peduncles 50–80mm, green flushed deep plum, appearing almost black. Fls 120–160mm wide, starry, glossy, not scented; tepals 25–30mm wide, tips very slightly recurved; pedicels 70–100mm, ascending, dark green flushed deep plum, appearing almost black. Lvs 70–120 × 20–40mm, leathery, glossy, very dark green. Stems 1.25–1.75m, with up to 12 fls; infl. racemose. Mid-season.

‘Uguntiņa’ I(a/b)

‘Rotaļa’ × ‘Dīana’ (Kireeva)

H: Ā. Zorgevics, 1990 **G:** Ā. Zorgevics, 1993

N: Ā. Zorgevics, 1994 **I:** University of Latvia Botanical Garden, 2001 **REG:** Ā. Zorgevics and University of Latvia Botanical Garden, 2008

Tepals inside, including throat, vivid reddish orange (32A); outside paler. Spots brown, scattered, in basal part of tepal; nectaries clear; pollen brown; stigma orange. Fls 120mm wide, not scented; tepals 70 × 20–40mm, margins smooth, tips slightly recurved. Lvs 80 × 15mm, scattered, glossy. Stems 0.9m, greenish, tinged grey, with up to 9 fls. Mid-season (early July).

Published in *Lilium Balticum. Latvijas liliju selekcionāru izaudzētās lilijas*, No. 1, 1995

Name means “small light” in Latvian.

‘Vainode’ VI(b/a)

‘Atblāzma’ × ‘Laimrota’

H: Ā. Zorgevics, 1990 **G:** Ā. Zorgevics, 1993

N: Ā. Zorgevics, 1994 **I:** University of Latvia Botanical Garden, 2000 **REG:** Ā. Zorgevics and University of Latvia Botanical Garden, 2008

Tepals yellowish white (155D); inside with light yellow (15D) patch towards base; outside centre flushed with greenish violet stippling; throat pale green. Nectaries clear; pollen orange-yellow; stigma brownish. Fls 130mm wide, strongly scented; tepals 90 × 20–40mm, margins slightly ruffled, tips strongly recurved. Lvs 110 × 10mm, whorled. Stems 1.2m, green, with up to 9 fls. July.

Published in *Lilium Balticum. Latvijas liliju selekcionāru izaudzētās lilijas*, No. 1, 1995

Named after a town in Latvia.

‘Veļetoč’ VI(b-c/b)

Syn. of ‘Divoženka’

‘Velocita’ I(a-b/b)

Parentage unknown

H: Mak Breeding BV, pre-2007 **N:** Mak Breeding BV, 2007 **REG:** Mak Breeding Rights BV, 2008

Tepals vivid reddish orange (N30C); midveins N30B at base. Spots few; papillae present; nectaries orange; pollen orange-red; stigma purple-red. Tepals short and of narrow to medium width, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length and breadth. Stems of average height to tall, green with some darker markings, with few fls. Early.

‘Vivendi’ VIII(a-b-c) LA

Seedling BW 95-03 × seedling W 91-22

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of inner tepals, including throat, greenish white (157D). Inside of outer tepals pale yellow-green (4D). Outside of outer tepals 4D; midribs strong yellow-green (144C), tinged 144A at top. Outside of inner tepals greenish white (157D); midribs strong yellow-green (144C). Spots and papillae absent; nectaries strong yellow-green (145A); pollen dark orange (N163A); stigma light yellow-green (145C).

Fls 235mm wide; tepals 135 × 75mm, margins smooth, tips recurved. Lvs 183 × 20mm, dark green. Stems 1.3m, green, with up to 5 fls.

‘Vivendum’ VIII(a/b) OT

Seedling PH 96-06 × seedling RM 96-49

H: Vletter en Den Haan Beheer BV, pre-2005

N: Vletter en Den Haan Beheer BV, 2005

REG: Vletter en Den Haan Beheer BV, 2008

Inside of tepals, including throat, yellowish white (155D); light greenish yellow (5D) patch next to and above nectaries; base of young fls tinged red-brown. Outside of tepals pale yellow-green (155A, flushed 4D); midribs light yellow-green (145B). Spots absent; papillae few, light greenish yellow (5D); nectaries strong yellow-green (144C); pollen greyish red (176A); stigma yellowish grey (156B). Fls 240mm wide, scented; tepals 155 × 73mm, margins ruffled, tips recurved. Lvs 185 × 37mm. Stems 1m, green, with up to 3 fls.

‘Wawel’ VIII(a-b/b-c) LA

‘Viking’ × Longistar Group

H: J. Kapias, pre-2009 **N:** J. Kapias, pre-2009

I: A. Berner, 2009 **REG:** A. Berner, 2009

Tepals pale greenish yellow; midribs green. Spots small, dark violet-brown, over basal half, fairly numerous on inner tepals, sparse on outer tepals; nectaries green, with frosted white edge; pollen brown; stigma pale yellow-green. Fls 145mm wide; tepals 90 × 40mm, tips slightly recurved; pedicels 50mm, stout, sparsely spotted dark brown; with secondary buds. Lvs 120 × 18mm, matt, mid-green. Stems 0.8m, very stout, with a few dark brown spots, with short infl. Late June.

‘White Bouquet’ I(a/c)

Syn. of ‘Tyrolia’

‘White Dancer’ I(a/d)

(‘Eng’ × ‘Last Dance’) × ‘Stardom’

H: S.D. Paulson, 2004 **G:** S.D. Paulson, 2006

N: S.D. Paulson, 2009 **REG:** S.D. Paulson, 2009

Inside of tepals light yellow-green (154D) on opening, maturing to white; throat, and outside of tepals, light yellow-green (154D). Spots fairly large, elliptic, dark purplish pink (186C), widely and fairly evenly scattered over basal two-thirds of each tepal; nectaries light yellow-green (154D); pollen dark purplish pink (186C); stigma light greenish yellow (4B). Fls 80mm wide, not scented; tepals 80 × 25mm, margins smooth, tips strongly recurved. Lvs 120 × 15mm, scattered, slightly hairy. Stems 1.2m, deep yellowish green (141B), with up to 8 fls. Dec.-Jan. (New Zealand).

‘White Grace’ V(b-c/a)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2003

N: Hobaho BV, 2003 **REG:** Testcentrum voor

Siergewassen BV, 2007

Inside of tepals, and outside of inner tepals, greenish white (155C); midveins green at base. Spots and papillae absent; nectaries green; pollen orange-yellow;

stigma cream. Tepals long and of medium width to broad, margins (very) slightly ruffled, tips slightly recurved. Lvs of medium length, and of medium width to broad. Stems 1.45m, green, with up to 8 fls. Very late.

Published in *Registraties* week 44-2006 t/m 45-2007, KAVB (2007)

‘White Hero’ VII(a-b/c)

Parentage unknown

H: Testcentrum voor Siergewassen BV, pre-2009

REG: Testcentrum voor Siergewassen BV, 2009
Inside of tepals white (NN155C); outside of inner tepals predominantly NN155C. Spots and papillae absent; nectaries green; pollen orange-brown; stigma dark purple. Tepals of medium length to long and of medium width to broad, margins slightly ruffled to ruffled, tips slightly recurved to recurved. Lvs of medium length, and broad, dark green. Stems (very) tall, green with faint darker spots and stripes, with few to average no. of fls. Mid-season.

‘Wildcat’ I(a/-)

H: Cebeco (USA), pre-2008

Inside of tepals fawn-gold shading to ivory-white in upper half; midveins pale gold. Spots dark brown-black, elliptic, very conspicuous, of medium size, evenly and quite densely distributed on basal half of each tepal, sometimes coalescing along basal half of midveins to form lines and dashes; papillae absent; filaments white-gold; pollen dark brown-black; style russet brown shading to golden brown towards stigma. Tepal margins more or less smooth, sometimes a little wavy, tips slightly recurved. Stems 0.9m. Mid-season.

Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Wizard’ I(a-b/b)

Parentage unknown

H: Bischoff Tulleken Lelies BV, pre-2003 **N:** Liliande BV, 2003 **REG:** Mak ‘t Zand BV, 2008

Inside of tepals vivid yellow (12A), overlaid dark red (187B/C) in basal portion and along margins in places. Outside of tepals brilliant yellow (12B) to light yellow (12C), with dark spots. Spots numerous, dark red (187C); papillae present; nectaries green; pollen orange-yellow; stigma yellow, with small, purple-pink spots. Tepals (very) short and narrow, margins (very) slightly ruffled, tips (very) slightly recurved. Lvs of medium length, and of narrow to medium width. Stems of average height to tall, green with some darker markings, with average no. to many fls. Early to mid-season.

Published in *The Lily Nook cat.*, 2009

‘X Factor’ VII(a/-)

Parentage unknown

REG: M.A. van den Bergen Zn, V.o.f., 2008

Inside of tepals strong purplish pink (73B) to light purplish pink (73C), paler with age; midveins yellow at base. Outside of inner tepals light purplish pink (73C). Spots numerous, white and pink; papillae present; nectaries green; pollen red-brown; stigma

grey-green or white. Tepals of medium length to long and of medium width to broad, margins ruffled, tips recurved. Lvs of medium length, and broad. Stems 1.1m, green with darker spots and stripes, with up to 3 fls. Mid-season to late.

‘Yellow Cocotte’ I(a/-)

N: D. Schoon, 2007

Inside of tepals bright, clear yellow; inner tepals with narrow, dark red margins. Stems 1.07m. Mid-season. Published in *The Lily Nook cat.*, 2009

External images: *The Lily Nook cat.*, 2009

‘Yellow Gadget’ I(a-b/b)

Parentage unknown

H: Mak ‘t Zand BV, pre-2008 **REG:** Mak ‘t Zand BV, 2008

Tepals vivid yellow (9A). Spots and papillae present; nectaries yellow-green; pollen orange-brown; stigma purple. Tepals (very) short and of narrow to medium width, margins slightly ruffled, tips slightly recurved to recurved. Lvs of short to medium length, and of narrow to medium width. Stems of short to average height, green with darker spots and stripes, with few to average no. of fls. Early.

‘Yellow Sun’ (Taeon) I(a/b-c)

‘Madras’ × ‘Galant’

H: Taeon Lily Experiment Station, 1998

G: Taeon Lily Experiment Station, pre-2004

N: Taeon Lily Experiment Station, pre-2006

I: Taeon Lily Experiment Station, c.2006

Tepals brilliant greenish yellow, central portion orange-yellow. Spots few, reddish brown; filaments fairly short, upright, yellowish; pollen and stigma reddish brown. Fls 170mm wide; tepal margins ruffled at mid-point, tips becoming strongly recurved. Lvs 110 × 15mm. Stems c.1.21m, reddish brown, with 4 or 5 fls in a racemose infl. Early June (under glass). Epithet published in Abstract S05-P-30 from 27th International Horticultural Congress and Exhibition, Korea, 2006.

It is not possible to register this cultivar epithet as it was registered for another lily by De Vries Veredeling BV in 2003 (ICNCP, 2004: Art. 27.1).

‘Yujiao’ VII(b/b)

‘Tome’ × ‘Siberia’

H: W. Dong, 2001 **G:** W. Dong, 2007 **N:** W. Dong, 2007 **I:** H.P. Zhang, 2006 **REG:** Wei Dong & Cheng du United Bio-technology Co. Ltd, 2007

Tepals white; yellow patch beside midveins at base inside; midveins yellow at base shading to white at top; throat yellow. Spots yellow, sparse but widely distributed over basal three-quarters of tepal; papillae yellow, conspicuous; nectaries green; pollen red; stigma green. Fls 250mm wide, strongly scented; tepals 130 × 65mm, margins strongly ruffled, tips strongly recurved. Lvs 85 × 22mm, scattered. Stems 1.15m, green, with up to 6 fls. Mid-season.

The epithet is a transcription of Mandarin Chinese meaning “pretty girl”.

‘Yun Xia’ VII(a/c)

Seedling D32 × seedling D6

H: X.W. Wu, 2001 **G:** X.W. Wu, 2003 **N:** X.W. Wu, 2004 **REG:** Institute of Flower Research, Yunnan Academy of Agricultural Sciences, 2008

Tepals deep purplish pink (N66D); throat pale orange-yellow (159C). Brushmark pale orange-yellow (159C), above throat; spots and papillae deep red (60A); nectaries light olive (152B); pollen strong brown (172A); stigma yellowish white (N155D). Buds orange-yellow, flushed green and pink. Fls 230mm wide, strongly scented; tepals 120 × 45mm, margins slightly ruffled, tips slightly recurved. Lvs 100 × 30mm, scattered. Stems 0.85m, purple, with up to 8 fls.

The epithet is a transcription of Mandarin Chinese meaning “rosy clouds”.

‘Yun Ying’ VII(a/c)

Seedling D36 × seedling D6

H: X.W. Wu, 2003 **G:** X.W. Wu, 2007 **N:** X.W. Wu, G. Cui, 2007 **REG:** Institute of Flower Research, Yunnan Academy of Agricultural Sciences, 2008

Tepals deep purplish pink (N74C), outside shading to moderate red (47A) towards base of tepals; narrow, white margins; throat strong yellow (153D). Spots moderate red (47A); nectaries and stigma bluish white (N155A); pollen strong orange (169C). Fls 210mm wide, strongly scented; tepals 105 × 30mm, margins slightly ruffled, tips slightly recurved. Lvs 80 × 26mm, scattered. Stems 0.88m, green, with up to 8 fls.

The epithet is a transcription of Mandarin Chinese meaning “clouds of Oriental cherry blossom”.

‘Zanloberge’ VIII(b/b) LO

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2007 **N:** Van Zanten Flowerbulbs BV, 2007 **REG:** Van Zanten Flowerbulbs BV, 2008

Trade designation AUBERGE

Inside of tepals strong purplish pink (73B) to light purplish pink (73C), shading to nearly white towards tips and base; midveins green at base. Outside of inner tepals very pale purple (73D). Spots and papillae absent; nectaries green; pollen orange-brown; stigma grey-green. Fls 200mm wide; tepals long and broad, margins slightly ruffled, tips slightly recurved. Lvs of medium length to long, and of medium width to broad. Stems 1.2–1.55m, green with faint darker spots and stripes, with up to 5 fls. Late.

‘Zanlopres’ VII(a/b)

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2009 **N:** Van Zanten Flowerbulbs BV, pre-2009

Trade designation RED IMPRESSION

Inside of tepals deep red (60A), shading to moderate purplish red (64A) towards margins; midveins deep red (53A); throat yellowish white (155D). Outside of outer tepals deep red (60A), closely spotted yellowish white (155D) at tips; midribs strong yellow-green (N144C) at base shading to moderate yellow-green (146B) towards top with moderate yellowish green (138A) tip. Outside of inner tepals strong purplish red (60C), shading to moderate purplish red (70A)

towards margins and yellowish white (155D) beside midribs at base; midribs strong yellow-green (N144D) at base. Spots dark red (59A); papillae yellowish white (155D) and dark red (59A); nectaries strong yellow-green (N144C); pollen dark orange (N163A); stigma pale yellow-green (195C). Fls 250mm wide, scented; tepals 135 × 80mm, margins slightly ruffled, tips slightly recurved. Lvs 165 × 60mm. Stems 0.95–1.15m, green, with up to 6 fls.

An application for registration was turned down in 2009, as the cultivar epithet is too similar to, and likely to cause confusion with, ‘Zanlapres’, registered by Van Zanten Flowerbulbs BV in 2006 (ICNCP, 2004: Art. 19.25). However, as this lily was granted European Plant Breeders’ Rights under the name ‘Zanlopres’ in 2009, it remains the accepted epithet (*ibid.*: Art. 24.4).

‘Zanlorbato’ VII(a/b-c)

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2007 **N:** Van Zanten Flowerbulbs BV, 2007 **REG:** Van Zanten Flowerbulbs BV, 2008

Trade designation RUBATO

Tepals deep purplish pink (68A) to strong purplish pink (68B), bluer with age; inside paler towards base; midveins yellow-green at base. Spots and papillae present; nectaries yellow-green; pollen reddish brown; stigma grey-green. Fls 250mm wide; tepals of medium length and breadth, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and (very) broad. Stems 1.6–1.85m, green with darker spots and stripes, with up to 6 fls. Mid-season.

‘Zanlorena’ VII(a-b/b)

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2007 **N:** Van Zanten Flowerbulbs BV, 2007 **REG:** Van Zanten Flowerbulbs BV, 2008

Trade designation MAURENA

Inside of tepals deep purplish pink (N57C/D), paler towards base; paler with age. Outside of inner tepals deep purplish pink (N57D). Spots and papillae present; nectaries green shading to white; pollen orange-brown; stigma purple, with small spots. Tepals of medium length and breadth, margins ruffled to strongly ruffled, tips slightly recurved to recurved. Lvs of medium length to long, and of medium width to broad. Stems 1.35m, green with some darker markings, with up to 5 fls. Early to mid-season.

‘Zanlorexus’ VII(a/b-c)

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2008 **N:** Van Zanten Flowerbulbs BV, 2008 **REG:** Van Zanten Flowerbulbs BV, 2008

Trade designation LEXUS

Inside of tepals dark purplish pink (186C); tips strong purplish red (64C); base above nectaries, and throat, yellowish white (155D). Outside of tepals dark purplish pink (186C), yellowish white (155D) beside midribs at base; midribs of outer tepals strong

yellow-green (144C) at base and top, yellowish white (155D) between; midribs of inner tepals light yellow-green (144D) at base, dark purplish pink (186C) at top, yellowish white (155D) between. Spots vivid purplish red (N57B); papillae deep purplish pink (64D); nectaries strong yellow-green (144C); pollen strong reddish orange (169A); stigma light yellow-green (145B). Fls 225mm wide, scented; tepals 123 × 56mm, margins smooth to slightly ruffled, tips slightly recurved. Lvs 190 × 45mm. Stems 1–1.2m, green, with up to 7 fls.

‘Zanlorsig’ VII(a/b)

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2009 **N:** Van Zanten Flowerbulbs BV, 2009 **REG:** Van Zanten Flowerbulbs BV, 2009

Trade designation **SIGNUM**

Inside of tepals whiter than white (NN155D), with a small pale greenish yellow (2D) patch above nectary; throat white (NN155D). Outside of tepals white (NN155D); midribs of outer tepals strong yellow-green at base (144C) and top (144B), pale greenish yellow (2D) between; midribs of inner tepals strong yellow-green (144C) at base shading to pale greenish yellow (2D) towards top. Spots and papillae few, white (NN155D); nectaries strong yellow-green (144B); pollen strong reddish orange (169A); stigma dark purplish red (N79A). Fls 255mm wide, scented; tepals 150 × 90mm, margins slightly ruffled, tips recurved. Lvs 205 × 58mm. Stems 1.23m, green, with up to 5 fls.

‘Zanlorva’ VII(a/b-c)

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2009 **N:** Van Zanten Flowerbulbs BV, pre-2009 **REG:** Van Zanten Flowerbulbs BV, 2009

Trade designation **MERIVA**

Inside of tepals whiter than white (NN155D), with a small pale greenish yellow (2D) patch above nectary; throat white (NN155D). Outside of tepals white (NN155D); midribs strong yellow-green (143C) at base, shading to light yellow-green (150D), top of outer tepals moderate yellow-green (143D). Spots and papillae white (NN155D); nectaries strong yellow-green (144C); pollen dark orange (N163A); stigma dark purplish red (N79A). Fls 290mm wide, scented; tepals 160 × 86mm, margins slightly ruffled, tips recurved. Lvs 160 × 58mm. Stems 1.2m, green, with up to 4 fls.

‘Zanotfora’ VIII(a/b) OT

Parentage unknown

H: Van Zanten Flowerbulbs BV, pre-2009 **N:** Van Zanten Flowerbulbs BV, pre-2009 **REG:** Van Zanten Flowerbulbs BV, 2009

Trade designation **AMFORA**

Inside of tepals brilliant yellow (8A), inner tepals shading to light greenish yellow (7D) at tips; outer tepals shading to 7D towards margins; throat light greenish yellow (7D). Outside of outer tepals yellowish white (NN155A), with vivid reddish orange (40A) stripe beside midribs; midribs strong

yellow-green (143C) at base shading to strong yellowish green (N144A) at top. Outside of inner tepals pale yellow (11D), shading to light greenish yellow (7D) towards margins; midribs strong yellow-green (144B) at base shading to light greenish yellow (8B) towards top. Spots brilliant yellow (8A); papillae absent; nectaries strong yellow-green (144C); pollen greyish brown (166A); stigma greyish yellow-green (191B). Fls 245mm wide, scented; tepals 155 × 75mm, margins smooth, tips slightly recurved. Lvs 230 × 45mm, dark green. Stems 1.2m, green, with up to 3 fls.

‘Zhonghua Huangguan’ VII(b/b)

‘Devotion’ × ‘Rosy Dream’

H: W. Dong, 2001 **G:** W. Dong, 2007 **N:** W. Dong, 2007 **I:** H.P. Zhang, 2006 **REG:** Wei Dong & Cheng du United Bio-technology Co. Ltd, 2007

Tepals, including throat, yellow; midribs pinkish at base. Spots absent; papillae few; nectaries green; pollen yellow; stigma purple. Fls 160mm wide, slightly scented; tepals 115 × 53mm, margins slightly ruffled, tips slightly recurved. Lvs 80 × 20mm, scattered. Stems 1.1m, green, with up to 5 fls.

The epithet is a transcription of Mandarin Chinese meaning “Chinese crown”.

‘Zooty Beauty’ VI(b-c/b)

Beautifully scented yellow-flowered Div. VI

(Trumpet/Aurelian hybrid) seedling × unknown

H: D. Hercbergs, pre-2008 **G:** D. Hercbergs, pre-2008 **N:** D. Hercbergs, c.2008 **I:** D. Hercbergs, c. 2008 **REG:** D. Hercbergs, 2008

Inside of tepals pinkish white (N155B); midveins brilliant greenish yellow (6B); throat and base of tepals moderate yellowish green (139B). Outside of outer tepals light purplish pink (65B). Spots absent; pollen scarce, strong orange (24A), some fls with none; stigma dark. Fls 120mm wide, slightly scented; tepals 110 × 45mm, margins smooth, tips recurved. Lvs 120 × 25mm, scattered. Stems 1.2m, green with darker markings, with up to 9 fls. Mid-season to late.

‘Zotina’ VII(a-b/b)

A temporary name for ‘**Frederique’s Choice**’ prior to registration.

Hybridizers, registrants and others


This list includes selectors, namers and introducers in addition to hybridizers and registrants.

(† denotes deceased)

- BAZETT, R.H.
Kelowna, British Columbia, Canada
- BERBÉE, J.
Hillegom, the Netherlands
- BERG, M.A. van den, en Zn V.o.f.
't Zand, the Netherlands
- BERNER, A.
Dolistowo, Jaświly, Poland
- BISCHOFF TULLEKEN LELIES BV
Wieringerwerf, the Netherlands
- BLIGH, L.C.
Burnie, Tasmania, Australia
- CAM LILIES
Camdale, Tasmania, Australia
- CEBECO LILIES, Inc. (USA)
Aurora, Oregon, USA
- CHENG, Qian Fa
Yongkang City, Zhejiang, People's Republic of China
- COLLICUTT, L.M.
Agriculture Canada, Morden, Manitoba, Canada
- COLUMBIA-PLATTE LILIES, Inc.
The Lily Garden, Woodland, Washington, USA
- CUI, Guangfen
Institute of Flower Research, Yunnan Academy of Agricultural Sciences, Kunming, People's Republic of China
- DONG, Wei
Chengdu, Sichuan, People's Republic of China
- DUDEK, J.
Hlučín, Czech Republic
- EGGER, D.L.
Wilsonville, Oregon, USA
- ESTATE PERENNIALS
Stony Plain, Alberta, Canada
- EVANS, A.
[address details unknown]
- FREEMAN, Mrs J.F. [formerly McRae]
Boring, Oregon, USA
- FREY, K.
Estate Perennials, Stony Plain, Alberta, Canada
- GRIESBACH, R.J.
Delavan, Wisconsin, USA
- HALL, R.
Chapel House, nr Newcastle upon Tyne, England, UK
- HART'S NURSERY
Oakhanger, Cheshire, England, UK
- HAY, D.J.
Saskatoon, Saskatchewan, Canada
- HERCBERGS, D.
Vecumnieki, Bauskas rajons, Latvia
- HOBACHO BV
Lisse, the Netherlands
- HOOGLAND, P.D.
the Netherlands
- HUANG, Jing
Institute of Horticultural Research, Guizhou Academy of Agricultural Sciences, Guiyang, People's Republic of China
- HYDE and SON, H.W.
Ruscombe, Berkshire, England, UK
- IMANSE BV, H.P.
Voorhout, the Netherlands
- INSTITUTE of FLOWER RESEARCH,
YUNNAN ACADEMY of AGRICULTURAL SCIENCES
Kunming, People's Republic of China
- JONG Beheer BV, De
Andijk, the Netherlands
- JONG LELIES HOLLAND BV, De
Andijk, the Netherlands
- JORDAN, N.
Camdale, Tasmania, Australia
- † JOSLIN, M.
Spruce Grove, Alberta, Canada
- KAPIAS, J.
Radostowice, Poland
- KIRSCH, E.T.
Myrtle Point, Oregon, USA
- KRUMINŠ, A.
Latvia
- LAAN, Gebr. N. & B., en Zn
Bovenkarspel, the Netherlands
- LAAN FLORA FACILITIES BV
Zwaagdijk, the Netherlands
- LIEROP, W. van, en Zn BV
Anna Paulowna, the Netherlands
- LILIANDE BV
't Zand, the Netherlands
- LILY COMPANY BV
Andijk, the Netherlands
- LILY NOOK, The
Neepawa, Manitoba, Canada
- LIN, Qingdan
Institute of Horticultural Research, Guizhou Academy of Agricultural Sciences, Guiyang, People's Republic of China
- LUO, Fuxian
Institute of Horticultural Research, Guizhou Academy of Agricultural Sciences, Guiyang, People's Republic of China
- MAK BREEDING BV
Wieringerwerf, the Netherlands
- MAK BREEDING RIGHTS BV
Wieringerwerf, the Netherlands
- MAK 't ZAND BV
't Zand, the Netherlands
- MAK-LEEK, Inc. (USA)
Scio, Oregon, USA
- MAK-LEEK, R.
Independence, Oregon, USA
- MANTÉL, H.
Mount Hood Lilies, Sandy, Oregon, USA

- MAREL LELIE BV, Van der
's-Gravenzande, the Netherlands
- MARKLILY BV
Lisse, the Netherlands
- MAVERIDGE INTERNATIONAL BV
Groenveldsdijk, Sint Maarten, the Netherlands
- MEGO, A.
Bratislava, Slovakia
- MELRIDGE Inc.
Gresham, Oregon, USA
- MICULKA, B.
Velehrad, Czech Republic
- MOUNT HOOD LILIES, Inc.
Sandy, Oregon, USA
- PATTERSON, C.F.
c/o University of Saskatchewan Dept of
Horticultural Science
- PAULSON, S.D.
Timaru, New Zealand
- ROBINSON, E.
Gaybird Nursery, Wawanesa, Manitoba, Canada
- RONALD, W.G.
La Prairie, Manitoba, Canada
- RUITER BLOEMBOLLEN BV
Andijk, the Netherlands
- SALM, R.W.J. van der
't Zand, the Netherlands
- SALM (USA), Van der
Woodland, Washington, USA
- SALM, Fa Van der, & De GOEDE, V.o.f.
't Zand, the Netherlands
- SCHOON, D.
Callantsoog, the Netherlands
- SHAW, J.M.H.
Selston, Nottinghamshire, England, UK
- SHI, Lejuan
Institute of Horticultural Research, Guizhou
Academy of Agricultural Sciences, Guiyang,
People's Republic of China
- SIMPSON, M.
Ulverstone, Tasmania, Australia
- † SKINNER, F.L.
Dropmore, Manitoba, Canada
- STEINBRÜCK, G.
Berlin, Germany
- STROHMAN, B.
The Lily Nook, Neepawa, Manitoba, Canada
- TAEAN LILY EXPERIMENT STATION
Chungcheongnamdo Agricultural Research and
Extension Services, Tae'an-Gun, Korea
- TESTCENTRUM voor SIERGEWASSEN BV
Hillegom, the Netherlands
- "THE ORIGINALS" U.A., Coöp. Kwekersvereniging
Breezand, the Netherlands
- TLACH, M.
Šumice, Czech Republic
- UNIVERSITY of LATVIA BOTANICAL
GARDEN
Rīga, Latvia
- VINK, Fa Q.J., en Zn BV
Sint Maartensvlotbrug, the Netherlands
- VINK, W.
the Netherlands
- VLETTER en Den HAAN Beheer BV
Rijnsburg, the Netherlands
- VLETTER, J.A., en Gebr.
Rijnsburg, the Netherlands
- VLETTER, J.A., en Gebr. & FLORIS
Rijnsburg, the Netherlands
- WADEKAMPER, J.
Parkdale, Oregon, USA
- WEI DONG & CHENG DU UNITED
BIO-TECHNOLOGY Co. Ltd
Cheng du, Sichuan, People's Republic of China
- WHITELEY, T.
Evenley, Northamptonshire, England, UK
- WIDERSKI, R.
Rybnik, Poland
- WISTUBA, P.
Orzepowicka, Rybnik, Poland
- WORLD BREEDING BV
Breezand, the Netherlands
- WU, Xue Wei
Institute of Flower Research, Yunnan
Academy of Agricultural Sciences, Kunming,
People's Republic of China
- WYNN-JONES, B. and S.
Crŷg Farm Plants, Caernarfon, Gwynedd,
Wales, UK
- YONGKANG ZHEJIANG JIANGNAN LILY
BREEDING Co Ltd,
Yongkang City, Zhejiang, People's Republic
of China
- ZANTEN FLOWERBULBS BV, Van
Hillegom, the Netherlands
- ZHANG, Hailong
Institute of Horticultural Research, Guizhou
Academy of Agricultural Sciences, Guiyang,
People's Republic of China
- ZHANG, Hong Ping
Cheng du, Sichuan, People's Republic of China
- ZHENG, Sixiang
Institute of Horticultural Research, Guizhou
Academy of Agricultural Sciences, Guiyang,
Guizhou Province, People's Republic of China
- ZHOU, Qi
Institute of Horticultural Research, Guizhou
Academy of Agricultural Sciences, Guiyang,
People's Republic of China
- ZIMMERMAN, T.
Minnetrista, Minnesota, USA
- ZORGEVICS, A.
University of Latvia Botanical Garden, Rīga, Latvia

ISBN 978-1-907057-07-6



9 781907 057076 >