



The International Rhododendron Register & Checklist (2004) *Seventh Supplement*



© 2012 The Royal Horticultural Society
80 Vincent Square, London SW1P 2PE, United Kingdom
www.rhs.org.uk

International Rhododendron Registrar:
Dr A C Leslie

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781907057311

Printed and bound in the UK by Page Bros, Norwich

Cover: Rhododendron 'Mrs Charles E. Pearson', by Alfred John Wise
Image courtesy of RHS Lindley Library

The International Rhododendron Register and Checklist (2004)

Seventh Supplement

Names registered, January–December 2011	page 2
Names and addresses	page 30
Corrections	page 32

Introduction

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first six Supplements included names registered from 1 January 2003 to 31 December 2010. This seventh Supplement includes all those registered in 2011.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7–10 of the Introduction. It may be helpful here to repeat that the following abbreviations are employed:

(a)	azalea
(az)	azaleodendron
cv.	cultivar
G	grown to first flower by
Gp	Group
H	hybridised by
I	introduced by
N	named by
(r)	rhododendron
R	raised by
REG	registered by
S	selected by
(s)	seed parent
(v)	vireya

Where possible colour references are given to the *RHS Colour Chart* (1966, 1986, 1995 & 2001) with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS website (www.rhs.org.uk/plants/plant-science/plant-registration-forms/rhodoform) or from one of the national or regional registrars as follows:

North America	Mrs J.W. Murray, 21 Squire Terrace, Colts Neck, New Jersey 07722-1021, USA
Australia	Mr K. Gillanders, Woodbank, 2040 Huon Road, Longley, Tasmania 7150, Australia
New Zealand	Mr B.J. Coker, 42 Massey Crescent, St Albans, Christchurch 8014, New Zealand
Japan	Dr K. Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken, Japan 768-0021

Dr A.C. Leslie
International Rhododendron Registrar
109 York Street
Cambridge CB1 2PZ
United Kingdom

(email: alanleslie@rhs.org.uk)

Names registered January–December 2011

(r) 'Adrien Franchet'

cv. Elepidote rhododendron: 'Kernéostic' × 'Jalisco Elect' (s) × 'Fred Wynniatt' × 'Ring of Fire'
H: M. Colombel (1997), G: M. Colombel (2006),
N: M. Colombel (2011), REG: M. Colombel (2011)
Fls 8–12/domed truss, broadly funnel-campanulate, 65–70 × 100–115mm, with 7 wavy-edged lobes, buds moderate red (47A), opening inside deep pink (51C) at the margins and light orange-yellow (16B) in the centre, outside similarly coloured; inconspicuous brownish spots on the three dorsal lobes; ovary green and hairy, but not glandular; style exserted. Calyx 2–8mm long, vivid reddish orange (33B), lobes ciliate. Pedicels deep red (60A). Lvs elliptic, 110–135 × 60–70mm, base mostly cordate (a few rounded), apex obtuse, margins flat, matt green above, hairless; petioles mostly red shading towards the apex into the greenish yellow midvein. Shrub 1.3 × 2.0m in 12 years. Mid- to late May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the famous French botanist who described many rhododendron species

(r) 'Aeschbacher's Rubin'

cv. Elepidote rhododendron: 'Oratorium' (s) × 'Feuerschein'
H: H. Hachmann (1979), G: H. Hachmann (1984),
N: Holger Hachmann (2007), I: Holger Hachmann (2007), REG: M. Oppenrieder (2011)
Fls 10–16/dense, rounded truss, funnel-shaped, 60–85mm wide, with 5 slightly wavy-edged lobes, vivid red (53A) inside and out, with a blotch of dark red (almost black or red-brown) spots on the dorsal lobe; filaments and style bright red; anthers vanilla yellow. Lvs elliptic, up to 160 × 61mm, margins wavy and decurved, glossy green above. Shrub 1.2 × 1.4m in 10 years. Mid-May to early June.

Syn: 'Schwitter's Rubin'
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009). Etymology: named for a well-known Swiss showmaster (MC or Master of Ceremonies) Kurt Aeschbacher

(r) 'Aivars Lasis'

cv. Elepidote rhododendron: open-pollinated seedling from ('Dace' × 'Purple Splendour')
H: R. Kondratovičs (1985), G: R. Kondratovičs (1991),
N: R. Kondratovičs (2008), REG: Rhododendron Nursery Babite, University of Latvia (2011)
Fls 17/strongly domed truss, open funnel-shaped, 80 × 80mm, with 5 very wavy-edged lobes, buds moderate purplish red (59C), opening inside light purple (75A), outside strong reddish purple (72C), with a strong blotch of greyish purple (N77A) on the dorsal lobes; filaments white, anthers white; style white, stigma cherry red. Calyx 2mm long, light green. Lvs elliptic, 130 × 60mm, rounded at base, pointed at apex, with

wavy-edged margins, glossy green above, hairless. Shrub 2.5 × 2.3m in 20 years. Late May to early June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March 2009, p.37. Etymology: named for the Latvian dendrologist, Aivars Lasis

(r) ANDREA

Trade designation for 'Hachmann's Andrea'

(r) 'Andris'

cv. Elepidote rhododendron: 'Catawbiense Grandiflorum' (s) × *yakushimanum* 'Koichiro Wada'
H: R. Kondratovičs (1974), G: R. Kondratovičs (1980),
N: R. Kondratovičs (2009), REG: Rhododendron Nursery Babite, University of Latvia (2011)
Fls 15/slightly domed truss, open funnel-shaped, 60 × 60mm, with 5 wavy-edged lobes, buds light purple (75A), opening inside white (155D) in the centre and very pale purple (69C) at the margins, outside very light purple (75C), with inconspicuous, discrete, strong greenish yellow (153C) spots on the dorsal lobes; filaments white, anthers light brown; style white, stigma purple-red. Calyx 1mm long, light green. Lvs elliptic, 115 × 40mm, rounded at base, pointed at apex, margins flat, matt green above, hairs white when young, aging to brown. Shrub 1.9 × 2.5m in 20 years. Late May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2010, p. 36. Etymology: Andris is a Latvian male christian name

(r) 'Angelina Dee'

cv. Elepidote rhododendron: 'Edith Bosley' (s) × 'Edwin O. Weber'
H: J. Doppel (2005), G: J. Doppel (2009),
N: J. Doppel (2010), REG: J. Doppel (2011)
Fls 13/ball truss, broadly funnel-shaped, c.50 × 65mm, with 5 very wavy-edged lobes, deep purplish red (71A) in bud, opening inside light reddish purple (74C), with a large, prominent, dorsal blotch of moderate yellow-green (146C) spots on a paler (nearly white) ground; outside light reddish purple (74C). Lvs elliptic, 75 × 30mm, broadly acute at apex, rounded at base, margins flat, hairless. Shrub with a dense habit, 0.6 × 0.8m in 10 years. Mid-May.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the granddaughter of the registrant

(r) 'Aotearoa'

cv. Elepidote rhododendron: parentage unknown
H: J.B. Brueckner (pre 1994), G: J.B. Brueckner,
N: C. Woodward (2010), REG: C. Woodward (2011)
Fls 10/lax truss, open funnel-shaped, 55–60 × 60mm, with 6 wavy-edged lobes, deep pink (51B–C) in bud, opening inside pale yellow-green (4D), with a strong yellow-green (145A) spotted blotch; outside pale yellow-green (4D). Calyx 2mm long, strong yellow-green (144C). Lvs elliptic, 150 × 55mm, broadly acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.8 × 2.2m in 25 years. Early

May to early June. Both very cold hardy and tolerant of prolonged periods of high temperatures. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: Dr Brueckner lived for many years in New Zealand, where his interest in rhododendrons originated; this epithet is the most widely known and accepted Maori name for that country – as translated it means “the land of the long white cloud”

(a) **‘Arismajesty’**

cv. Evergreen azalea: ‘Ostalett’ (s) × un-named seedling (YB-1556)
H: W.R. Bergman (1999), G: W.R. Bergman (2001), S: W.R. Bergman (2001), N: Aris Horticulture, Inc. (c. 2009), REG: Aris Horticulture, Inc. (2011)
Fls semi-double, 1–3/shoot apex, ‘rotate and rose-like’, 63mm wide, with a 5-lobed outer whorl and 4–6 petaloids, lobes undulate and ruffled, close to strong purplish red (72A) at first, when fully open more intense than vivid reddish purple (N74A) and shading to deep purplish pink (N74C) to very light purple (75C) towards the base; spots and speckles close to deep red (60A); no normal stamens or pistil present. Calyx 5-lobed, 7mm long, strong yellow green (144A). Lvs mostly elliptic, c.58 × 27mm, mucronate at apex, attenuate at base, darker than moderate olive green (147A) above, hairy, retained well during the forcing process; petioles c.12mm long, green. Shrub with an outward-spreading, mounding habit, 23 × 37cm in one year. Spring (when not forced).
Trade designation: MAJESTY

(a) **‘Arouca’**

cv. Deciduous azalea: ‘Stagecoach Frills’ (s) × ‘Twilight Sky’
H: K.-H. Hübbers (1989), G: K.-H. Hübbers (1994), N: Mrs L. Hübbers (1994), REG: J. Westhoff (2011)
Fls 12/lax, domed truss, tubular funnel-shaped (opening out widely at the mouth of the tube), 110mm wide, with 5 wavy-edged lobes (some flowers slightly doubled by virtue of some petaloid stamens), buds greenish (strong yellow-green, 145A) with red stripes and yellow patches, opening inside creamy yellow (light orange-yellow/pale yellow, 19B/C), outside creamy yellow (pale orange-yellow, 19D) with red (strong red, 39A) stripes. Calyx 3mm long, greenish. Lvs oblong-elliptic, 90 × 35mm, margins flat to slightly upcurved, cuneate at base, broadly acute at apex, glossy green above, slightly bronzed when young and with good autumn colour, hairy (with tiny white hairs along the margin and the midribs). Shrub 1.0 × 0.8m in 10 years. Early to mid-May.
Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: named for a town in Trinidad

(r) **‘Atis’**

cv. Elepidote rhododendron: open-pollinated seedling from ‘Catawbiense Grandiflorum’
H: R. Kondratovičs (1957), G: R. Kondratovičs (1963), N: R. Kondratovičs (2007), REG: Rhododendron Nursery Babīte, University of Latvia (2011)
Fls 15/strongly domed truss, funnel-shaped, 70 × 70mm,

with 5 wavy-edged lobes, buds moderate purplish red (59C), opening inside very pale purple (75D) in centre, with deep purplish pink (73A) margins, outside deep purplish pink (73A), with discrete, relatively inconspicuous strong yellow-green (144A) spots on the dorsal lobes; filaments white, anthers light grey; style light pink, stigma cherry red. Calyx 1mm long, light green. Lvs elliptic, 150 × 60mm, rounded at base, pointed at apex, margins flat, glossy green above, with hairs white when young, aging to light brown. Shrub 2.5 × 3.0m in 20 years. Late May to early June.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March 2009, p. 37. Etymology: Atis is a Latvian male christian name

(r) **ATTRAKTION**

Trade designation for ‘Hachmann’s Attraktion’

(r) **‘Austin’s Firecracker’**

cv. Elepidote rhododendron: ‘Red River’ (s) × ‘John Paul II’
H: J. Doppel (2001), G: J. Doppel (2005), N: J. Doppel (2011), REG: J. Doppel (2011)
Fls 10/ball truss, broadly funnel-shaped, c.75 × 70mm, with 5 wavy-edged lobes, red in bud, opening inside close to deep pink (52B) at margins, changing to white in the centre of each lobe, with brilliant orange-yellow (21B) spotting on the dorsal lobe; outside red. Lvs elliptic, 115 × 45mm, broadly acute at apex, rounded at base, margins downcurved, dull and olive green above, hairless. Shrub 0.8 × 0.9m in 10 years. Late May.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: Austin is the name of the registrant’s grandson

(r) **‘Ayomi’**

cv. Elepidote rhododendron: ‘Janet Blair’ (s) × *yakushimanum*
H: J.B. Brueckner (1969), G: J.B. Brueckner, N: A. Brueckner, A. Woodward & A. Keefe (2010), REG: C. Woodward (2011)
Fls 17/conical truss, broadly funnel-shaped, 45 × 75mm, with 5 very wavy-edged lobes, strong reddish purple (70B) and strong purplish pink (63C) in bud, opening inside very pale purple (76D), edged light purple (84C), with a rather faint dorsal blotch of brilliant yellow (11A) spots; outside very pale purple (76D), with a light purple (84B) streak on the centre of each lobe. Calyx 2mm long, light yellow-green (150D). Lvs elliptic, 100–135 × 40–50mm, broadly acute at apex, oblique at base, the margins flat, dull and moderate olive green (147A) to dark yellowish green (139A) above, hairless. Shrub 1.2 × 1.5m in 41 years. May to mid-June. Not only very cold hardy but responded well to extended heat waves.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the epithet is an amalgam of the names (Aya and Omi) that Marta Brueckner’s grandchildren have always called her. She herself had a significant influence and played a vital support role in the work of her husband (J.B. Brueckner) in his breeding of new rhododendron cultivars

(r) **'Babītes Sarkanaīs'**

cv. Elepidote rhododendron: 'Catawbiense Grandiflorum' (s) × 'Scandinavia'
H: R. Kondratovičs (1976), G: R. Kondratovičs (1982), N: R. Kondratovičs (2004), I: Rhododendron Nursery Babīte, University of Latvia (2004), REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 16/strongly domed truss, open funnel-shaped, 80–90 × 80–90mm, with 5 wavy-edged lobes, buds deep red (60A), opening deep red (53A) throughout, with discrete, dark greyish purple (202A) spots on the dorsal lobes; filaments red, anthers light violet; style red, stigma black. Calyx 2mm long, dark red. Lvs elliptic, 130 × 50mm, rounded at base, pointed at apex, margins wavy, matt green above, hairless. Shrub 1.6 × 2.0m in 10 years. Early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2011, p. 29. Etymology: "Sarkanaīs" means red in Latvian and refers to the flower colour, "Babītes" refers to the nursery which introduced the plant

(r) **'Biruta'**

cv. Elepidote rhododendron: 'Nova Zembla' (s) × 'Panenka'

H: R. Kondratovičs (1985), G: R. Kondratovičs (1991), N: R. Kondratovičs (2005), I: Rhododendron Nursery Babīte, University of Latvia (2005), REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 16/strongly domed truss, open funnel-shaped, 100 × 100mm, with 5–6 flat-edged lobes, buds deep red (60A), opening strong purplish pink (68B) throughout, with relatively inconspicuous, discrete, deep red (60A) spots on the dorsal lobes; filaments pink, anthers yellow; style pink, stigma dark violet. Calyx 2mm long, light green. Lvs elliptic, 145 × 55mm, pointed at base and apex, margins flat, matt green above, hairless. Shrub 3.0 × 4.0m in 20 years. Late May to early June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2011, p. 29. Etymology: Biruta is a Latvian female christian name

(r) **'Blauschimmer'**

cv. Elepidote rhododendron: selection from *campanulatum* subsp. *aeruginosum*

H: H. Hachmann, G: H. Hachmann, N: H. Hachmann, I: H. Hachmann (1992), REG: M. Oppenrieder (2011)

Fls 5–8/dense, rounded truss, campanulate, 38–42mm wide, purplish violet. Lvs elliptic-oblong, margins flat, glossy above and light bluish for several weeks at first, later becoming green, with a dense russet brown indumentum below. Dense spherical shrub, 1.0 × 1.4m in 10 years, not flowering until c.15 years old. Mid- to late May.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 1993/94 (1993)

(a) **BLOOM-A-THON LAVENDER AZALEA**

Trade designation for 'RLH1-4P19

(a) **BLOOM-A-THON PINK DOUBLE AZALEA**

Trade designation for 'RLH1-2P8'

(a) **BLOOM-A-THON RED AZALEA**

Trade designation for 'RLH1-1P2'

(a) **BLOOM-A-THON WHITE AZALEA**

Trade designation for 'RLH1-3P3'

(r) **'Blue Tango'**

cv. Elepidote rhododendron: 'Purple Amethyst' (s) × 'Plum Passion'

H: J. Barlup (2002), G: J. Barlup (2006), N: J. Barlup (2010), REG: J. Barlup (2011)

Fls 18/ball truss, broadly funnel-shaped, 45 × 70mm, with 5 wavy-edged lobes, deep purplish red (71A) in bud, opening inside light purple (77D), edged strong purple (77B), with moderate red (183C) spotting on the dorsal lobe extending c.30mm from the base; outside light purple (77D), edged strong purple (77B), with midveins strong purple (77B); filaments and anthers white. Lvs elliptic, 125 × 50mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 8 years. Late May. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY)

(a) **'Bonny'**

cv. Deciduous azalea: *viscosum* (s) × 'Sarina'

H: H. Hachmann (1985), G: H. Hachmann (1989), N: H. Hachmann (1991), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls single, 13–14/dense, flat rounded truss, tubular funnel-shaped, 50–55 × 50–60mm with 5 wavy-edged lobes, inside strong purplish red (63B), outside purplish rose pink, with a strong orange (25B) blotch on the dorsal lobe extending from the base to the middle; filaments and style deep red (60A); style very long, exerted, stigma green; intensely sweet-scented; flowers star-shaped, the lobes moderately tapered and only slightly recurved. Lvs elliptic, with flat margins, orange-red to red-brown in autumn. Shrub 1.0 × 1.2m in 10 years. Late May to mid-June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **'Burgundy Moon'**

cv. Elepidote rhododendron: 'Black Sport' (s) × 'Purple Lace'

H: R. Shellenberger (1988), G: R. Shellenberger (1996), N: R. Shellenberger (2005), REG: R. Shellenberger (2011)

Fls 16/domed truss, broadly funnel-shaped, c.50 × 65mm, with 5 wavy-edged lobes, dark red (187A) in bud, opening inside dark red (59A), with a dark red (187A), spotted blotch in the throat; outside dark red (59A). Lvs elliptic, 150 × 50mm, broadly acute at apex, cuneate at base, the margins flat, dull and moderate olive green (137A) above, hairless. Shrub 1.5 × 1.5m in 10 years. Mid-May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY)

(r) **'Calypso'**
Syn. of 'Hachcher'

(a) **'Cape Cod'**

cv. Evergreen azalea: 'Wakaebisu' (s) × 'Gumpo White'
H: S.F. McDonald (1974–1975), G: S.F. McDonald (1977), N: S.F. McDonald (c. 1980), I: Middle Atlantic Chapter, American Rhododendron Society, REG: S.F. McDonald (2011)

Fls solitary, hose-in-hose, broadly funnel-shaped, 35 × 60mm, with 5+5 very wavy-edged lobes, opening white inside and out, tinged pale green near the base; unmarked; very floriferous Calyx 35mm long, white, with pale green near the base. Spring leaves elliptic, c.40 × c.15mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (137A) above, with hairs above and below (very pale tan at first, aging to pale tan); summer leaves obovate and smaller. Shrub 1.1 × 1.8m in 35 years. Late May. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the plant was first displayed at an American Rhododendron Society conference in Cape Cod

(r) **'Carl Peter'**

cv. Elepidote rhododendron: 'September Song' (s) × 'White Dimples'
H: T. Ahern (1997), G: J. Doppel (2000), N: J. Doppel (2011), REG: J. Doppel (2011)
Fls 10/ball truss, broadly funnel-shaped, c.45 × 75mm, with 5 wavy-edged lobes, moderate purplish pink (54D) in bud, opening inside white, with a light greenish yellow (8C) throat and a dorsal blotch of discrete brilliant greenish yellow (151D) spots; outside white. Lvs elliptic, 115 × 50mm, obtuse at apex, rounded at base, margins upcurved, semi-glossy and mid-green above, hairless. Shrub 1.5 × 1.7m in 17 years. Mid-May. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a son of the registrant

(r) **'Carolyn Scholz'**

cv. Elepidote rhododendron: 'Aeschbacher's Rubin' (s) × 'Tarantella'
G: H. Hachmann (2000), H: H. Hachmann (1995), N: Holger Hachmann (2007), I: Holger Hachmann (2007), REG: M. Oppenrieder (2011)
Fls 19–26/dense, rounded truss, funnel-shaped, 72–80mm wide, with 5 wavy-edged lobes, red inside and out, with dark red marks on the central, basal part of the dorsal lobe; filaments bright red, with white hairs at the base, anthers creamy yellow; style bright red. Lvs elliptic, glossy green above, hairless. Shrub 1.2 × 1.4m in 10 years. May. Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009). Etymology: named to honour a young woman who died a violent death

(a) **'Charlotte Megan'**

cv. Deciduous azalea: hybrid derived from 'Homebush'
H: W.G.T. Hyde (1970s), G: W.G.T. Hyde (1970s), N: Mrs R. Legrand (2011), I: Millais Nurseries (2011), REG: Mrs R. Legrand (2011)

Fls double, 10–12/tight ball truss, trumpet-shaped, 55 × 21mm, with 9–10 very wavy-edged lobes, strong purplish red (54A) in bud, opening deep purplish pink (55A) both inside and out, with a blotch of brilliant orange-yellow (23B) on the two dorsal lobes; sweetly scented. Calyx irregular, 3–8mm long, strong yellow-green (144B). Lvs lanceolate, 95 × 30mm, attenuate at base, narrowly acute at apex, matt green above, margins recurved, hairless. Shrub 2.1 × 1.6m in at least 30 years. May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a granddaughter of the registrant

(r) **'Cherry Kiss'**

Syn. of 'Hachcher'

(r) **'Christian Gehler'**

cv. Elepidote rhododendron: 'Soubrette' (s) × 'Sarasate'
H: H. Hachmann (1993), G: H. Hachmann (1998), N: Holger Hachmann (2004), I: Holger Hachmann (2004), REG: M. Oppenrieder (2011)

Fls 11–15/ dense, rounded truss, open funnel-shaped, up to 90mm wide, with 5 wavy-edged lobes, the dorsal lobe slightly reflexed, bright pink both inside and out, with the centre of the lobes paler and with a large cherry-red blotch on the dorsal lobe and parts of the adjacent lobes; filaments short, white; style longer, white, exerted, stigma red. Lvs elliptic, up to 88 × 45mm, glossy green, margins mostly decurved, some wavy and upcurved. Shrub 0.9 × 1.0m in 10 years. Late May to mid-June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009). Etymology: named for the editor-in-chief of the German gardening journal *Gartenzeitung*

(a) **'Cinzia'**

cv. Deciduous azalea: 'Humboldt Picotee' (s) × 'Schneegold'
H: Holger Hachmann (1993), G: Holger Hachmann (1997), N: Holger Hachmann (1999), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls single, 7–13/dense, rounded truss, tubular funnel-shaped, 70–80mm wide, with 5 wavy-edged lobes, white inside, with a large golden yellow flare on the dorsal lobe and parts of the adjacent lobes, outside intense yellowish cream in bud, with central veins and base of the tube reddish, opening creamy white, with traces of pink; filaments and style exerted, pinkish white; spicily scented. Lvs elliptic-obovate, margins flat, reddish green at first, becoming dark green, and with autumn colours of brown-red and yellow-orange. Shrub 1.0 × 1.2m in 10 years. Mid-May to mid-June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(a) **'Cita'**

Syn. of 'Hachkissen'

(r) **'Copper Dust'**

cv. Elepidote rhododendron: 'Cimarron Sun' (s) × 'Big Deal'

H: J. Barlup (2001), G: J. Barlup (2006), N: J. Barlup (2010), REG: J. Barlup (2011)
Fls 25/conical truss, broadly funnel-shaped, 50 × 75mm, with 7 wavy-edged lobes, deep yellowish pink (39B) in bud, opening inside pale yellow (20D), with dorsal lobes light yellow (20B) fading to pale yellow (20D) at edges, with a dorsal flare of deep red (185A) spots extending upwards c.40mm from the base; outside similar to inside but without the markings. Calyx irregular, up to c.45mm long, light yellow (20B), edged pale yellow (20D), with moderate purplish red (186A) flares. Lvs elliptic, 115 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.9m in 10 years. Early May.

(r) **'Cranberry Ice'**

cv. Elepidote rhododendron: parentage unknown – grown from seed collected at Muckross Gardens, Killarney, Ireland, in June 1990
G: R.E. George (2011), S: R.E. George, N: R.E. George (2011), REG: R.E. George (2011)
Fls 9 or more/ball truss, funnel-campanulate, 50 × 75mm, with 5–7 wavy-edged lobes, cranberry red in bud, opening inside deep red in the throat; outside red at base, paling more gradually to the pale pink lobes; unmarked. Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, semi-glossy and deep green above, hairless. Shrub 3.0–3.7 × 1.5m in 21 years (growing in deep shade until recently). Mid-May. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the name suggested by the flower colour

(a) **'Dalina'**
Syn. of 'Hachneg'

(r) **'Dan Carter'**

cv. Elepidote rhododendron: 'Lem's Monarch' (s) × 'Alice Hennessy'
H: T.A. Garbutt (1990), G: A. Sim (1995), N: T.A. Garbutt (2011), REG: T.A. Garbutt (2011)
Fls 20/ open ball truss, open campanulate, with 7–8 strongly wavy-edged lobes, red in bud, opening inside pink, with a deeper pink margin and a small darker pink blotch at the base of the dorsal throat, outside mainly deep rose pink. Lvs lanceolate, 110 × 35mm, acute at apex, oblique at base, margins flat, matt green above, hairless. Shrub 1.0–3.0 × 2.0m in 10 years. Mid-October (New Zealand).
Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the nephew of the registrant's son

(r) **'Daphne's Pearl'**

cv. Elepidote rhododendron: 'Lemon Lodge' (s) × 'Alice Hennessy'
H: T.A. Garbutt (1992), G: J. Leggett (2000), N: T.A. Garbutt (2011), REG: T.A. Garbutt (2011)
Fls 15/conical truss, open campanulate, 65 × 95mm, with 7 wavy-edged lobes, buds white, opening pure white, with inconspicuous greenish spots towards the base of the dorsal throat. Calyx 2mm long, green. Lvs ovate, 145 × 55mm, acute at apex, obtuse at base,

margins flat, matt green above, hairless. Shrub 2.0 × 1.5m in 18 years. Late November (New Zealand).
Nomenclatural standard: colour print provided by the registrant (WSY)

(r) **'Dave's Delight'**

cv. Elepidote rhododendron: 'Newburyport Belle' (s) × 'Olin O. Dobbs'
H: J. Doppel (2004), G: J. Doppel (2007), N: J. Doppel (2011), REG: J. Doppel (2011)
Fls 10/ball truss, broadly funnel-shaped, c.45 × 70mm, with 5 flat-edged lobes, vivid reddish purple (72B) in bud, opening inside very light purple, edged with a broad band of strong reddish purple (70B), with a dense, dark red (187A) basal blotch and numerous spots of the same colour extending upwards on the dorsal and two adjacent lobes; outside moderate purplish pink (74C). Lvs elliptic, 115 × 40mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and mid-green above, hairless. Shrub 0.8 × 0.7m in 7 years. Mid-May. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the son-in-law of the registrant

(r) **DENISE**

Trade designation for 'Hachmann's Denise'

(az) **'Dentelle'**

cv. Azaleodendron: *periclymenoides* (s) × 'Karen Triplett'
H: L. & A. Le Duigou (2005), G: L. & A. Le Duigou (2010), N: L. & A. Le Duigou (2011), REG: L. & A. Le Duigou (2011)
Fls 8/ truss (flat early on but domed when fully open), tubular funnel-shaped, 50 × 50mm, with 5 wavy-edged lobes, buds pale yellow (20C), touched with strong yellowish pink (32C), opening white inside and out, without markings; stamens 10; style 50mm long, pale yellow-green (149D), stigma moderate orange (171C), exserted; ovary hairy; slightly scented. Calyx 35mm long, very deeply divided into linear, white (155A), strongly reflexed lobes. Lvs elliptic, 100 × 30mm, cuneate at base, acute at apex, margins slightly decurved, matt green above, with no evident indumentum; semi-deciduous. Shrub 47 × 30cm in 5 years. June.
Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY). Etymology: derived from the finely divided calyx, "dentelle" being the French for lace

(a) **'Dita Krenberga'**

cv. Deciduous azalea: a Rustica azalea (s) × 'Uguns'
H: R. Kondratovičs (1978), G: R. Kondratovičs (1984), N: R. Kondratovičs (2008), REG: Rhododendron Nursery Babīte, University of Latvia (2011)
Fls fully double, 13/slightly domed truss, tubular funnel-shaped, 70 × 70mm, with 15 wavy-edged lobes, buds vivid reddish orange (34B), opening inside light purplish pink (55C), outside strong purplish pink (62A), with pale orange-yellow (159C) ribs; a brilliant yellow (21C) blotch on the dorsal lobes; normal stamens absent; style red, stigma green. Calyx 3mm

long, green. Lvs elliptic, 130 × 40mm, pointed at base and apex, wavy-edged, glossy green above, hairless. Shrub 3.0 × 3.0m in 20 years. Late May to early June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2010, p.36. Etymology: named for the Latvian musician Dita Krenberga

(a) **'Doctor Pratt's Late Red'**

cv. Deciduous azalea: parentage unknown
R: M.C. Pratt, N: The High Beeches Gardens Conservation Trust (2011), REG: The High Beeches Gardens Conservation Trust (2011)
Fls single, 5–6/loose hemispherical truss, tubular funnel-shaped, with 5 slightly wavy-edged lobes, inside deep yellowish pink (43C), with the dorsal lobe strong orange (30C), outside with the lobes deep to moderate yellowish pink (39B–C), with more yellowish midribs, the dorsal lobe light orange/light yellowish pink (29B–C), the tube moderate reddish orange (42D) and with stalked glands and eglandular hairs; unspotted; style and filaments vivid reddish orange (43A–B), long exserted; anthers orange, tipped red; stigma whitish, darkening with age; not scented. Calyx c.5mm long, pale green, with dense, long, white marginal hairs. Lvs broadly elliptic, to oblanceolate or obovate, up to 60 × 27mm, acute and apiculate at apex, narrowly cuneate at base, margins very slightly decurved, with pale hairs on the margins, sparsely hairy above (these hairs lost with age), hairy on the main veins below. Shrub 3 × 3m in c.40 years. Late June to early July.
Nomenclatural standard: herbarium specimen provided by the registrant (WSY)

(r) **'Eda-Jenny'**

cv. Elepidote rhododendron: 'Admiral Vanessa' (s) × 'Welse'
H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2007), N: K.-H. Hübbers (2008), REG: J. Westhoff (2011)
Fls up to 10/rounded truss, broadly funnel-shaped, 45 × 80mm, with up to 7 wavy-edged lobes, buds light purplish pink (63D) at apex, strong purplish red (64B) at base, opening inside pale purplish pink (62D) at the margins and pale yellowish pink (27D) along the middle of each lobe, outside pale purplish pink (62D) with a darker line along the midrib of each lobe; strong purplish red (59D) spots forming a fan over the three dorsal lobes. Calyx 2mm long, greenish red. Lvs oblong to obovate, 140 × 55mm, margins flat, rounded at base, broadly acute and mucronate at apex, hairless. Shrub 0.5 × 0.4m in 5 years. Late May to early June. Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: named for two sisters, artists from Albania, Jenny Joosten and Eda Zari. Shrub broader than 'Holi' and not so well-branched

(r) **'Egons'**

cv. Elepidote rhododendron: open-pollinated seedling from 'Catawbiense Grandiflorum'
H: R. Kondratovičs (1969), G: R. Kondratovičs (1975), N: R. Kondratovičs (2003), I: Rhododendron Nursery Babīte, University of Latvia (2003),

REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 14/strongly domed truss, open funnel-shaped, 90 × 80mm, with 5 wavy-edged lobes, buds deep reddish purple (77A), opening strong reddish purple (N78B) throughout, with conspicuous, dark greyish purple (N186B) markings on the dorsal lobes; filaments purple, anthers violet; style red, stigma purple. Calyx 1mm long, purple violet. Lvs elliptic, 150 × 55mm, pointed at base and apex, margins flat, glossy green above, hairless. Shrub 1.3 × 1.4m in 8 years. Late May to early June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2010, p.35. Etymology: Egons is a Latvian male christian name

(r) **'Eisblume'**

cv. Elepidote rhododendron: *yakushimanum* 'Hanano-ego' (s) × *hemsleyanum*
H: K.-H. Hübbers (1990), G: K.-H. Hübbers (2001), N: K.-H. Hübbers (2011), REG: J. Westhoff (2011)
Fls up to 10/domed truss, open funnel-shaped, 50 × 70mm, with 6–7 wavy-edged lobes, white throughout, with two short splays of vivid yellow (13A) spots in the throat; filaments white, anthers pinkish; style and stigma pale yellow; slightly scented. Calyx minute. Lvs oblong-lanceolate, 120 × 60mm, margins slightly decurved, cordate at base, broadly acute and mucronate at apex, matt green above, with a dense brownish white indumentum above and below at first, with age only a few brown hairs remain below, especially near the midrib. Shrub 1.1 × 1.0m in 10 years. June. Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: German word for the pattern made by frost, literally "ice flower"

(r) **'Eispalast'**

cv. Elepidote rhododendron: 'Doctor Ans Heyting' (s) × 'Peucine' × unknown
H: K.-H. Hübbers (1993), G: K.-H. Hübbers (2001), N: W. Wüstemeyer (2011), REG: J. Westhoff (2011)
Fls 10/loose domed truss, broadly funnel-shaped, 80 × 70mm, with 5 very wavy-edged lobes, buds pale greenish yellow (10D) at the apex, shading to deep pink (48B) at the base, opening inside pale yellow (8D) inside and out, with light olive/dark greenish yellow (152B/C) spots forming a blotch in the dorsal throat (and base of the dorsal lobe) and extending around the base of the throat; filaments white, anthers red-brown; stigma greenish. Calyx 2mm long, light green. Lvs oblong, 160 × 55mm, margins flat, slightly oblique/cuneate at base, broadly acute and mucronate at apex, hairless. Shrub 1.1 × 1.0m in 10 years. June. Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: literal translation from the German "ice palace", in reference to the flower colour

(r) **'Eiswein'**

cv. Elepidote rhododendron: 'Doctor Ans Heyting' (s) × 'Peucine'
H: K.-H. Hübbers (1999), G: K.-H. Hübbers (2006), N: K.-H. Hübbers (2011), REG: J. Westhoff (2011)

Fls c.12/domed ball truss, open funnel-shaped, 60 × 80mm, with 7 wavy-edged lobes, buds moderate purplish red to dark purplish pink (186B–C), fading into cream, opening inside light greenish yellow (4C), outside light greenish yellow (4C), with darker veins; strong orange-yellow (163B) spots on two-thirds of the dorsal lobes. Calyx 3mm long, reddish. Pedicels violet. Lvs broadly elliptic, 110 × 60mm, margins slightly decurved, rounded at base, obtuse and mucronate at apex, matt green above, hairless. Shrub 1.2 × 1.1m in 10 years. June.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: named for the colour of the flowers and the raiser's fondness for this very special type of German wine

(r) **'Fay Dee'**

cv. Elepidote rhododendron: 'Diane's Delight' (s) × 'Cape White'

H: J. Doppel (2007), G: J. Doppel (2010), N: J. Doppel (2011), REG: J. Doppel (2011)

Fls 9/ball truss, broadly funnel-shaped, c.45 × 70mm, with 5 wavy-edged lobes, strong purplish red (67A) in bud, opening inside vivid purplish red (57C), with a prominent, dense, dorsal blotch of moderate purplish red (186A) spots; outside vivid red (57A). Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins downcurved, glossy and dark green above, hairless. Shrub 0.6 × 0.8m in 5 years. Mid-May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the daughter-in-law of the registrant

(a) **FEENKISSEN**

Trade designation for 'Hachkissen'

(a) **'Feuerköpfchen'**

cv. Deciduous azalea: 'Nabucco' (s) × 'Jack A. Sand'

H: Holger Hachmann (1998), G: Holger Hachmann (2003), N: Holger Hachmann (2008), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls double, 11–14/slightly loose, flat rounded truss, tubular funnel-shaped, 58–66mm wide, with wavy-edged lobes, bright red in bud, opening bright red, with apricot tones to the centre of the lobes; all stamens petaloid; outer lobes tapered and slightly reflexed, central lobes slightly irregular and more wavy-edged; delicately scented. Stems 1.1 × 1.1m in 10 years. Early May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **'Filigran'**

cv. Elepidote rhododendron: 'Graziella' (s) × 'Perlina' H: H. Hachmann (1995), G: H. Hachmann (2000), N: H. Hachmann (2001), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls 9–14/dense, rounded truss, broadly funnel-shaped, 85–95mm wide, with 5 flat to slightly wavy-edged lobes, intense pink outside in bud, opening inside bright pink, with a large bright red blotch of spots on the dorsal lobe; filaments and style purplish pink. Lvs linear, 70–75 × up to 17mm, margins flat, glossy

above, hairless. Shrub 0.6 × 0.9m in 10 years. May. Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **FINESSE**

Trade designation for 'Hachmann's Finesse'

(r) **FLUIDUM**

Trade designation for 'Hachmann's Fluidum'

(r) **'Forbidden Plateau'**

cv. Elepidote rhododendron: 'Courtenay Queen' (s) × *fortunei*

H: H.R. Wright (2005), G: H.R. Wright (2009), N: H.R. Wright (2011), REG: H.R. Wright (2011)

Fls 9/domed truss, open funnel-shaped, c.60 × 90mm, with 7 wavy-edged lobes, cream with a touch of pink in bud, opening inside white and changing to light pink with a pale yellow throat, a dark green eye and some inconspicuous light green spots on the dorsal lobe; outside white, changing to light pink. Calyx 5mm long, light green. Lvs oblong, 170 × 60mm, obtuse at apex, rounded at base, margins downcurved, semi-glossy green above, hairless. Shrub 0.6 × 0.3m in 6 years. Early May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the registrant is using names from the Comox Valley in British Columbia and the Forbidden Plateau is visible to the registrant from his garden. The flower colour (opening white and turning to pink) is reminiscent of a local legend about this plateau, where the local people sent their families to avoid frequent raids upon their homes; on one occasion the men returned to the plateau to retrieve their women and children, but although they searched until it snowed there was no sign of them. When it did snow the snow began to change from white to red and the place was declared evil and forbidden. Subsequently this phenomenon has been explained as a chemical reaction between the snow and a particular moss growing in that area

(r) **'Friesiane'**

cv. Elepidote rhododendron: 'Mareike' (s) × 'New Comet'

H: H. Hachmann (1993), G: H. Hachmann (1998), N: H. Hachmann (2002), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls 9–13/slightly loose, flat rounded truss, funnel-shaped, up to 80mm wide, with mostly 6 wavy-edged lobes, bright salmon pink in bud, opening inside bright yellow, outside yellow, with delicate bright red markings on the lower part of the dorsal lobe at first, but later these nearly fade away; filaments bright yellow; style greenish yellow, stigma green. Calyx with bright red margins and a green base. Pedicels reddish on adaxial surface, greenish on abaxial side. Lvs elliptic-orbicular, up to 100 × 65mm, obtuse and apiculate at apex, margins flat to decurved, glossy green above. Shrub 1.1 × 1.3m in 10 years. Late April to late May. Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) '**Glendoick Tanager**'

cv. *Lepidote rhododendron*: *brachyanthum* (s) × *viridescens*
H: P.A. & K.N.E. Cox (c. 1995), G: P.A. & K.N.E. Cox, N: P.A. & K.N.E. Cox, I: Glendoick Gardens Ltd, REG: P.A. & K.N.E. Cox (2011)
Fls 3–6/loose truss, campanulate, 18 × 20mm, with 5 wavy-edged lobes, pale greenish yellow (2D) in bud and both inside and out on opening, aging to light yellow-green (145D) on the outside; green spotting on the dorsal lobe; style green. Often has two inflorescences at each shoot apex. Calyx 3mm long, green. Lvs narrowly elliptic, up to 50 × 20mm, cuneate at base, obtuse at apex, margins flat, matt green above, with a scaly indumentum. Shrub 1.0 × 1.0m in 15 years. Late May to early June.
Trade designation: TANAGER

(a) '**Glühlicht**'

cv. *Deciduous azalea*: 'Betty Kelly' (s) × 'Fanal'
H: H. Hachmann (1967), G: H. Hachmann (1971), N: H. Hachmann (1973), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls single, 10–13/dense, rounded truss, tubular funnel-shaped, 58–85mm wide, with 5–6 flat-edged lobes, inside orange-red shading to bright orange in centre of lobes, with an orange-yellow blotch on the dorsal lobe, outside orange-red; filaments and style orange-red; style very long, exserted, stigma green; slightly sweet-scented. Lvs elliptic, with flat margins, glossy green above, reddish green when first emerging and dark brown-red in autumn. Shrub 1.0 × 1.2m in 10 years. May.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(a) '**Goldköpfchen**'

cv. *Deciduous azalea*: 'Annabella' (s) × 'Feuerwerk'
H: H. Hachmann (1986), G: H. Hachmann (1991), N: H. Hachmann (1993), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls single, 9–13/dense, rounded truss, tubular funnel-shaped, 50–70mm wide, with 5–6 flat to slightly wavy-edged lobes, buds bright orange-red, with conspicuous yellow-orange midveins, opening inside copper orange, with shades of orange-yellow. Lvs broadly elliptic, with flat margins, glossy green; reddish green at first in spring. Shrub 1.5 × 1.3m in 10 years. Late May to mid-June.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(a) '**Goldsinfonie**'

cv. *Deciduous azalea*: 'Annabella' (s) × 'Klondyke'
H: H. Hachmann (1986), G: H. Hachmann (1990), N: H. Hachmann (1993), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls single, 8–11/dense, rounded truss, tubular funnel-shaped (opening wide), 66–92mm wide, with 5–6, broad, very wavy-edged lobes, dark orange to bright red in bud, opening strong orange (25A), with traces of yellow-orange, aging to nearly brilliant

orange-yellow (21B); filaments and style bright orange, style exserted. Lvs elliptic, with flat margins, glossy green. Shrub 1.2 × 1.4m in 10 years. Mid- to late May.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) '**Gunter Dinger**'

cv. *Elepidote rhododendron*: 'Galaxis' (s) × 'Violette Funken'
H: H. Hachmann (1995), G: H. Hachmann (1999), N: Holger Hachmann (2004), I: Holger Hachmann (2004), REG: M. Oppenrieder (2011)
Fls 12–19/dense, rounded truss, funnel-shaped, 80–90mm wide, with 5 very wavy-edged, slightly reflexed lobes, purplish violet in bud, opening inside vivid reddish purple (78A), with a broad white stripe down the centre of each lobe, the midvein a slightly darker purplish violet, with a large blotch of nearly black spots on the dorsal lobe and parts of the adjacent lobes; filaments purple, anthers beige; style bright purple; both stamens and style exserted. Lvs elliptic, 120–140 × 30–45mm, margins decurved, glossy green above. Shrub 1.2 × 1.4m in 10 years. Mid-May to early June.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009). Etymology: named to celebrate the 70th birthday of the senior chief of the Dinger's Gartencenter, in Cologne, Germany

(r) '**Hachcher**'

cv. *Elepidote rhododendron*: 'Aeschbacher's Rubin' (s) × 'Busuki'
H: H. Hachmann (1995), G: H. Hachmann (2000), N: Holger Hachmann (2008), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 13–19/dense, flat-rounded truss, funnel-shaped, 70–80mm wide, with 5 wavy-edged lobes, bright red inside, red outside, with dark red spotting in the middle of the dorsal lobe. Lvs elliptic, up to 120 × 50mm, margins flat, glossy green above, hairless. Shrub 1.2 × 1.5m in 10 years. Mid-May to early June.
Syn: 'Calypso'. Trade designation: CHERRY KISS
Nomenclatural standard: colour print provided by the registrant (WSY). Originally published in Hachmann cat. 2009–2013 (2009) as 'Cherry Kiss', but this is now a registered trademark and had to be replaced as the epithet

(r) '**Hachhans**'

cv. *Elepidote rhododendron*: 'Midnight Mystique' (s) × 'Burgundula'
H: Holger Hachmann (1998), G: Holger Hachmann (2003), N: Holger Hachmann (2009), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 11–22/dense rounded truss, funnel-shaped, 45 × 70–85mm, with 5–7 flat-edged lobes, inside white with a slightly diffused transition to a broad marginal band of deep reddish purple (77A), dorsal lobe and parts of adjacent lobes with a dense fan of markings which are wine-coloured at the base and cinnamon near the apex. Lvs elliptic, 102–116 × 45–48mm, margins flat, dark glossy green above, hairless. Shrub 1.3 × 1.5m in 10 years. Mid-May to early June.

Syn: 'Night and Day'. Trade designation: HANS HACHMANN

Nomenclatural standard: colour print provided by the registrant (WSY). Originally published in Hachmann cat. 2009–2013 (2009) as 'Hans Hachmann', but this is now a registered trademark and had to be replaced as the epithet. Etymology: named after the raiser's father, the famous rhododendron breeder, Hans Hachmann

(a) '**Hachkissen**'

cv. Evergreen azalea: 'Melina' (s) × 'Panda'

H: Holger Hachmann (1997), G: Holger Hachmann (2000), N: Holger Hachmann (2008), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls double, 2–3/truss, broadly funnel-shaped, 50–55mm wide, with 5–11 wavy-edged lobes (with most stamens petaloid), buds white, with a greenish base, opening bright white throughout, unmarked; style white, stigma yellowish, ovary greenish. Lvs elliptic-ovate, margins flat, glossy green above, hairless. Flat-rounded shrub, up to 25 × 50cm in 10 years. Late May to mid-June.

Syn: 'Cita'. Trade designation: FEENKISSEN

Nomenclatural standard: colour print provided by the registrant (WSY). Originally published in Hachmann cat. 2009–2013 (2009) as 'Feenkissen', but the latter is now a trademark and had to be replaced as the epithet. Etymology: the trade designation alludes to the flower-covered plants being like small white cushions, "fee" meaning fairy and "kissen" meaning pillow

(r) '**Hachmann's Andrea**'

cv. Elepidote rhododendron: 'Fantastica' (s) × *williamsianum*

H: H. Hachmann (1986), G: H. Hachmann (1991), N: H. Hachmann (1993), I: H. Hachmann (2001), REG: M. Oppenrieder (2011)

Fls 6–8/loose truss, campanulate, 60–70mm wide, with 5 wavy-edged lobes, inside pinkish white, with broad pink margins, the outside richer pink throughout. Lvs rounded, up to 80 × 50mm, margins decurved, glossy dark green above, hairless. Broadly rounded shrub, 0.55 × 1.0m in 10 years. Late April to mid-May.

Trade designation: ANDREA

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2001–2005 (2001), under the unacceptable name 'Andrea'

(r) '**Hachmann's Attraktion**'

cv. Elepidote rhododendron: 'Andantino' (s) × 'Andantino'

H: H. Hachmann (pre 1994), G: H. Hachmann (1999), N: H. Hachmann (2005), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls 12–16/dense rounded truss, funnel-shaped, 44–60 × 80–105mm, with 5 wavy-edged lobes, strong red (53B) throughout, with a few darker red spots on the dorsal lobe; style and filaments reddish, with white hairs at the base, conspicuously held in a tight cluster in the centre of the flower; anthers dark brown. Lvs elliptic, 130–170 × 50–70mm, margins decurved, matt green above, hairless. Strong-growing, broadly erect shrub, 1.0 × 1.4m in 10 years. Late April to late May.

Trade designation: ATTRAKTION

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009) under the unacceptable name 'Attraktion'

(r) '**Hachmann's Denise**'

cv. Elepidote rhododendron: 'Abraxas' (s) × 'Campanile'
H: H. Hachmann (1998), G: H. Hachmann (2003), N: H. Hachmann (2003), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls 6–8/rather loose, flat-rounded truss, funnel-shaped, 67–86mm wide, with 5 flat-edged lobes, vanilla yellow ground, with broad border of purplish pink both inside and out, with numerous greenish spots on the lower part of the dorsal and adjacent lobes; filaments and style bright vanilla yellow; ovary greenish. Lvs elliptic, 150 × 65mm, margins decurved, rich glossy green above, hairless. Shrub 0.7 × 0.9m in 10 years. May.

Trade designation: DENISE

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009), under the unacceptable name 'Denise'

(r) '**Hachmann's Finesse**'

cv. Elepidote rhododendron: 'Gordian' (s) × 'Goldstück'

H: H. Hachmann (1997), G: H. Hachmann (2002), N: H. Hachmann (2003), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls 7–16/slightly loose, rounded truss, funnel-shaped, 65–80mm wide, with 5–7 wavy-edged lobes, rich primrose yellow throughout, with inconspicuous red markings at the base of the dorsal throat; filaments short, yellow; style nearly twice as long, greenish yellow, stigma bright green. Lvs elliptic, up to 76 × 45mm, margins wavy, matt green above, hairless. Shrub 1.0 × 0.9m in 10 years. Early May to early June.

Trade designation: FINESSE

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009) under the unacceptable name 'Finesse'

(r) '**Hachmann's Fluidum**'

cv. Elepidote rhododendron: 'Orangina' (s) × 'Belkanto'
H: H. Hachmann (1988), G: H. Hachmann (1993), N: H. Hachmann (1995), I: H. Hachmann (2001), REG: M. Oppenrieder (2011)

Fls 12–15/dense, rounded truss, funnel-shaped, up to 85mm wide, with 5 wavy-edged lobes, outside delicate rose with deeper coloured midribs, inside with a 10mm wide band of salmon pink surrounding a pale creamy yellow ground, the dorsal lobe with a large golden yellow flare which is copiously spotted with wine red, golden yellow and olive-green. Calyx intense salmon pink. Lvs narrowly elliptic, to 120 × 40mm, margins flat, glossy green above, hairless. Compact shrub, 1.4 × 1.6m in 10 years. Mid-May to early June.

Trade designation: FLUIDUM

Nomenclatural standard: colour print provided by the registrant (WSY). Originally published in Hachmann cat. 2001–2004 (2001) under the unacceptable name 'Fluidum'

(r) **'Hachmann's Mamamia'**

cv. *Elepidote rhododendron*: 'Sapporo' (s) × 'Osterschnee'

H: Holger Hachmann (1997), G: Holger Hachmann (2002), N: Holger Hachmann (2003), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls up to 16/ very dense, rounded truss, open funnel-shaped, up to 80mm wide, with 5 wavy-edged lobes, purplish pink in bud, opening bright pink inside and out, with a conspicuous, blackberry-coloured mark around the whole base of the corolla, extending up the dorsal side to cover up to half the dorsal lobe (and sometimes parts of an adjacent lobe), the mark breaking up into small spots towards the apex; filaments white; style nearly twice the length of the stamens, very thick, greenish white, with a bright green stigma. Lvs elliptic, up to 150 × 60mm, margins flat, glossy green above, hairless. Shrub 0.6 × 0.8m in 10 years. Mid-April to mid-May.

Trade designation: MAMAMIA

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009) under the unacceptable name 'Mamamia'

(r) **'Hachmann's Matador'**

cv. *Elepidote rhododendron*: 'Hachmann's Junifeuer' (s) × 'Aeschbacher's Rubin'

H: H. Hachmann (1994), G: H. Hachmann (1999), N: H. Hachmann (2000), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 8–15/round-domed truss, funnel-shaped, 73–84mm wide, with 5 wavy-edged lobes, deep red (60A) throughout, with numerous nearly black spots over the dorsal lobe and parts of the adjacent lobes; style and filaments reddish; anthers yellowish. Lvs elliptic, up to 145 × 67mm, with flat margins, glossy green above, hairless. Shrub 1.2 × 1.3m in 10 years. Mid-May to early June.

Trade designation: MATADOR

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009) under the unacceptable name 'Matador'

(r) **'Hachmann's Metallica'**

cv. *Elepidote rhododendron*: 'Marsalla' (s) × 'Peter Alan' H: Holger Hachmann (1994), G: Holger Hachmann (1999), N: Holger Hachmann (2000), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 14–21/dense, flat-rounded truss, open funnel-shaped, up to 100mm wide, with 5–6 wavy-edged lobes (the margins of the dorsal lobes often bent backwards), violet in bud, opening light purple (85A) throughout, with a large, conspicuous, dorsal marking of dark blackberry-coloured spots on a wine-coloured ground on the dorsal lobe and part of the adjacent lobes; filaments bright lilac; style reddish violet, stigma red. Lvs elliptic, margins flat, dark glossy green above, hairless. Shrub 1.4 × 1.6m in 10 years. Mid-May to mid-June.

Trade designation: METALLICA

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009) under the unacceptable name

'Metallica'

(r) **'Hachmann's Orakel'**

cv. *Elepidote rhododendron*: 'Pfauenauge' (s) × 'Galaxis' H: Holger Hachmann (1998), G: Holger Hachmann (2003), N: Holger Hachmann (2004), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 10–20/rounded truss, funnel-shaped, 70–87mm wide, with 5 wavy-edged lobes, rich violet inside and out, with a conspicuous blotch on the dorsal lobe and parts of the adjacent lobes, consisting of numerous blackberry-coloured spots on a wine-coloured ground; filaments pale purplish, anthers cream; style pink, khaki-coloured at the apex as is the stigma; base of filaments and ovary with white hairs. Lvs elliptic, margins flat, glossy mid-green, hairless. Shrub 1.0 × 1.1m in 10 years. Late May to mid-June.

Trade designation: ORAKEL

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009) under the unacceptable name 'Orakel'

(r) **'Hachmann's Saskia'**

cv. *Elepidote rhododendron*: 'Hachmann's Charmant' (s) × 'Andantino'

H: H. Hachmann (pre 1994), G: H. Hachmann (1999), N: H. Hachmann (2000), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 13–17/dense, rounded truss, funnel-shaped, 75–90mm wide, with 5 narrowly but very wavy-edged lobes, white ground, with a purplish pink margin and a blotch of, at first, greenish spots on a yellow ground, the spots aging to curry yellow; filaments and style white; anthers brown. Lvs elliptic, up to 180 × 60–70mm, margins decurved, glossy dark green above, hairless. Shrub 1.1 × 1.3m in 10 years. Early May to early June.

Trade designation: SASKIA

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009) under the unacceptable name 'Saskia'

(r) **'Hachmann's Tanaga'**

cv. *Elepidote rhododendron*: 'Hachmann's Kabarett' (s) × 'Schneespiegel'

H: Holger Hachmann (1991), G: Holger Hachmann (1996), N: Holger Hachmann (1998), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 10–23/dense, rounded truss, funnel-shaped, 76–90mm wide, with 5(–6) wavy-edged lobes, light purple (84B) inside, with the middle of the lobes paler, with a conspicuous, spotted, dorsal blotch covering most of the dorsal lobe and parts of the adjacent lobes – greenish yellow at first, aging to yellow and then orange to coppery red; outside darker in bud, opening like the inside, but with soft purplish red on the midveins; filaments white; style yellow at base, pinkish cream towards the apex. Lvs elliptic, margins flat, glossy dark green. Shrub 1.4 × 1.6m in 10 years. Early May to early June.

Trade designation: TANAGA

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann

cat. 2009–2013 (2009) under the unacceptable name ‘Tanaga’

(r) **‘Hachmann’s Veronika’**

cv. Elepidote rhododendron: ‘Trocadero’ (s) × ‘Lem’s Monarch’

H: H. Hachmann (1992), G: H. Hachmann (1997), N: H. Hachmann (1998), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls 7–17/rounded dome (sometimes slightly loose) truss, funnel-shaped, 90–100mm wide, with 5 wavy-edged lobes, vivid purplish red (67B) inside, with a much paler zone down the centre of each lobe and with a deep purple blotch at the base of the dorsal throat, breaking up apically into dots, only dots present around the rest of the base of the throat; filaments white; style greenish white below, white towards apex, strongly bent; ovary green. Lvs elliptic, margins flat, glossy dark green above, hairless. Shrub 1.2 × 1.4m in 10 years. Late May to mid-June.

Trade designation: VERONIKA

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009) under the unacceptable name ‘Veronika’

(a) **‘Hachneg’**

cv. Evergreen azalea: ‘Melina’ (s) × ‘Daphne’

H: Holger Hachmann (1998), G: Holger Hachmann (2001), N: Holger Hachmann (2008), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls single to semi-double, several at each shoot apex, broadly to tubular funnel-shaped, up to 50mm wide, with 5–10 flat-edged lobes (a few stamens of some flowers are petaloid), pinkish white, becoming darker pink at the margins and greenish yellow towards the base; sparse, tiny spots at the base of the dorsal lobe, greenish at first, aging to reddish; style white, stigma greenish at first, aging to yellow. Lvs elliptic to obovate, margins slightly upcurved, glossy green above, hairless. Shrub 0.5 × 0.8m in 10 years. Late May to mid-June.

Syn: ‘Dalina’. Trade designation: NEGLIGÉ
Nomenclatural standard: colour print provided by the registrant (WSY). Originally published in Hachmann cat. 2009–2013 (2009) as ‘Negligé’, but this is now a registered trademark and had to be replaced as the epithet. Etymology: the trade designation is suggested by the delicate colour of the flowers and is considered reminiscent of the delicate material and colours of such garments

(r) **HANS HACHMANN**

Trade designation for ‘Hachhans’

(a) **‘Helena Evelyn’**

cv. Deciduous azalea: hybrid derived from ‘Homebush’

H: W.G.T. Hyde (1970s), G: W.G.T. Hyde (1970s), N: Mrs R. Legrand (2011), I: Millais Nurseries (2011), REG: Mrs R. Legrand (2011)

Fls fully double, 10–14/tight ball truss, trumpet-shaped, 49 × 22mm, with 12 very wavy-edged lobes, deep pink (51B) in bud, opening inside and out light purplish pink (55C); sweetly scented. Calyx irregular, 4–6mm long, strong yellow-green (144B). Lvs elliptic,

90 × 33mm, cuneate at base, acute at apex, margins decurved, matt green above, hairless. Shrub 2.1 × 1.6m in at least 30 years. May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a granddaughter of the registrant

(v) **‘Hendrik’s Kanarie’**

cv. Lepidote rhododendron: ‘Satan’s Gift’ × ‘Flamenco Dancer’ (s) × ‘Vladimir Bukovsky’

H: H. Van Oost (2002), G: H. Van Oost (2004), N: H. Van Oost (2011), I: H. Van Oost (2011), REG: H. Van Oost (2011)

Fls 5–8/domed truss, tubular funnel-shaped, 82 × 65mm, with 5 wavy-edged lobes, buds yellow, opening with lobes deep yellow inside and out, throat and tube deep orange-yellow inside and out; unmarked; stigma pale green; anthers dark. Calyx vestigial. Lvs elliptic, 90 × 36mm, acute at apex, cuneate at base, with wavy edges, glossy green above, with silvery scales above and below. Slow-growing shrub, 0.7 × 0.5m in 10 years. March to June.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the epithet combines the registrant’s first name, combined with an allusion to the flower colour

(r) **‘Heritage Campfire Peach’**

cv. Elepidote rhododendron: parentage unknown

H: J.C. Cowles (1960s), G: Heritage Museums & Gardens (c 1970s), S: B. MacDonald (1986), N: Sandwich Club and Heritage Museums & Gardens (2010), I: Sandwich Club (1990s), REG: Sandwich Club (2011)

Fls 12/ball truss, open funnel-shaped, 55 × 90mm, with 6–7 wavy-edged lobes, strong pink (48D) in bud, opening inside pale yellow-green (155A), the lobes edged with moderate purplish pink (54D) and with twin, narrow, red rays on the dorsal lobe; outside pale yellow-green (155A), striped with strong pink (54C); stigma and style green; stamens rudimentary; slightly scented. Lvs oblong, 125 × 50mm, obtuse at apex, rounded at base, dull and moderate olive green (147A–B) above, hairless. Shrub with an open habit, 2.2 × 2.5m in 15 years. Late May (Cape Cod, Massachusetts, USA). Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY)

(r) **‘Heritage Pastel Perfume’**

cv. Elepidote rhododendron: parentage unknown

H: J.C. Cowles (1960s), G: Heritage Museums & Gardens (c 1970s), S: E.W. Collins (1985), N: Sandwich Club & Heritage Museums & Gardens (2010), I: Sandwich Club (1990s), REG: Sandwich Club (2011)

Fls 11/ball truss, open funnel-shaped, 55 × 90mm, with 6–7 wavy-edged lobes, light purplish pink (62C) in bud, opening inside yellowish white (158D), edged with pale purplish pink (62D) and with a strong greenish yellow (153B) flare on the three dorsal lobes; outside pale purplish pink (62D); strongly scented. Lvs oblong, 165 × 55mm, obtuse at apex, oblique at base, margins upcurved, semi-glossy and moderate olive green (146A) above, hairless. Shrub 2.2 × 2.5m in 15 years. Late May (Cape Cod, Massachusetts, USA).

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY)

(r) **'Heritage Snow Ruby'**

cv. Elepidote rhododendron: parentage unknown
H: J.C. Cowles (1960s), G: Heritage Museums & Gardens (c 1970s), S: E.W. Collins (1985), N: Sandwich Club and Heritage Museums & Gardens (2010), I: Sandwich Club (1990s), REG: Sandwich Club (2011)
Fls 12/domed truss, open funnel-shaped, 50 × 75mm, with 7 wavy-edged lobes, light purplish pink (62C) in bud, opening throughout pale purplish pink (62D) and maturing to white (155D), with prominent interior central markings of deep purplish red (59B) consisting of several irregularly shaped, flare-like marks extending upwards from the throat on to all the lobes, as well as moderate purplish red spotting on the three dorsal lobes. Lvs oblong, 140 × 50mm, obtuse at apex, rounded at base, semi-glossy and moderate olive green (146A) above, hairless. Shrub 2.2 × 2.5m in 15 years. Mid-May (Cape Cod, Massachusetts, USA).
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY)

(r) **'Holli'**

Syn. of 'Holli's Hope'

(r) **'Holli's Hope'**

cv. Elepidote rhododendron: ('Plum High' × 'Jonathan Shaw') F₂ (s) × 'Plum Passion'
H: J. Barlup (2000), G: J. Barlup (2005), N: J. Barlup (2010), REG: J. Barlup (2011)
Fls 16/ball truss, broadly funnel-shaped, 40 × 70mm, with 5 wavy-edged lobes, dark red (187A) in bud, opening inside light purple (76A), changing to strong reddish purple (72B) on dorsal lobes, with central lobe edged strong purplish red (72A) and with a dark red (187C) flare on the central dorsal lobe extending c.25mm from the base; outside light purple (76A), changing to strong reddish purple (72B) on dorsal lobes, with deep purplish red (71A) midveins; filaments and anthers white. Lvs lanceolate, 115 × 40mm, acute at apex, cuneate at base, margins flat, dull and olive green (147A) above, hairless. Shrub 0.3 × 0.5m in 5 years. Mid-May.

Syn: 'Holli'

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY)

(r) **'Holly's Choice'**

cv. Elepidote rhododendron: 'Blue Peter' (s) × 'White Dimples'
H: T. Ahern (1994), G: J. Doppel (1999), N: J. Doppel (2001), REG: J. Doppel (2011)
Fls 12/ball truss, broadly funnel-shaped, c.45 × 75mm, with 5 wavy-edged lobes, strong purplish red (67A) in bud, opening inside white, with a continuous band of very light purple (76B) at the margins and some strong greenish yellow (153C) spotting on the dorsal lobe; outside white, edged very light purple (76B). Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and dark green above, hairless. Shrub 1.8 × 1.2m in 16 years. Early May.
Nomenclatural standard: colour print of an electronic

image provided by the registrant (WSY). Etymology: named for the wife of the registrant

(r) **'Hōshun-in'**

cv. Elepidote rhododendron: *yakushimanum* (s) × *aureum*
H: H. Suzuki (1995), G: H. Suzuki (2007), N: H. Suzuki (2011), REG: H. Suzuki (2011)
Fls 10(–11)/hemispherical truss, open funnel-shaped, 50 × 70mm, with 5(–6) slightly wavy-edged lobes, buds yellowish white, suffused pale reddish orange, opening pale yellowish white throughout, with scattered, small, greenish brown spots on the dorsal lobe of some flowers. Lvs oblong, 100 × 30mm, acute to obtuse and apiculate at apex, cuneate at base, margins flat to slightly decurved, glossy green above, with a hairy indumentum below (white at first aging to brown). Shrub 50 × 40cm in 4 years (after grafting). Mid-May.
Nomenclatural standard: colour print provided by the registrant (WSY)

(a) **'Icacos'**

cv. Deciduous azalea: 'Washington State Centennial' (s) × 'Arouca'
H: K.-H. Hübbers (1994), G: K.-H. Hübbers (1999), N: Mrs L. Hübbers (2010), REG: J. Westhoff (2011)
Fls c.10/lax, domed truss, tubular funnel-shaped (opening out widely at the mouth of the tube), 60 × 80mm, with 5 very wavy-edged lobes, often recurved at the tips, moderate red (47A) in bud, opening inside light orange to light yellowish pink (29B/C), with darker veins, outside light orange-yellow (19B) and moderate reddish orange (35A/B), with red veins; slightly scented. Calyx 8mm long, light green, translucent. Lvs elliptic, 80 × 40mm, margins slightly upcurved, cuneate at base, broadly acute and mucronate at apex, glossy green above, with tiny white hairs on the margins and midvein. Shrub 0.8 × 0.7m in 10 years. Early May.
Nomenclatural standard: colour print of an electronic image provide by the raiser (WSY). Etymology: named for a village in Trinidad

(r) **'Ice Baller'**

cv. Elepidote rhododendron: *yakushimanum*, Exbury form (s) × 'Mrs J.G. Millais'
H: J. Barlup (1981), G: J. Barlup (1986), N: J. Barlup (2009), REG: J. Barlup (2011)
Fls 20/ball truss, saucer-shaped, 50 × 90mm, with 5 wavy-edged lobes, strong purplish red (64C) in bud, opening white (155D), with a densely spotted, moderate yellow (162A) dorsal flare extending c.30mm from the base; outside white (155D). Lvs elliptic, 145 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.5 × 2.4m in 29 years. Late May.

(r) **'Ilma'**

cv. Elepidote rhododendron: 'Catavbiense Grandiflorum' (s) × 'Mrs Furnivall'
H: R. Kondratovičs (1978), G: R. Kondratovičs (1984), N: R. Kondratovičs (2004), I: Rhododendron Nursery Babīte, University of Latvia (2004), REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 20/strongly domed truss, broadly funnel-shaped, 80 × 70mm, with 5 wavy-edged lobes, buds deep purplish pink (73A), opening inside moderate purplish pink (62B), outside strong purplish pink (68B), with discrete spots of dark red (59A) on the dorsal lobes; filaments light pink, anthers white; style pink, stigma purple. Calyx 2mm long, light green. Lvs elliptic, 125 × 60mm, rounded at base, pointed at apex, margins flat, matt green above, hairless. Shrub 2.2 × 3.0m in 20 years. Late May to early-June.

Trade designation: ILMA ČEPĀNE

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March 2009, p.27 as 'Ilma Čepāne', but granted PBR in Latvia as 'Ilma' so the latter must be used for this cultivar as the accepted name.

Etymology: named for a colleague of the registrant who is also a professor at the University of Latvia

(r) **ILMA ČEPĀNE**

Trade designation for 'Ilma'

(r) **'Imants'**

cv. Elepidote rhododendron: 'Midsummer' (s) × Elizabeth Group

H: R. Kondratovičs (1975), G: R. Kondratovičs (1981), N: R. Kondratovičs (2004), I: Rhododendron Nursery Babīte, University of Latvia (2004),

REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 15/strongly domed truss, open funnel-shaped, 90 × 90mm, with 5 narrowly very wavy-edged lobes, buds strong purple (77B), opening inside light purple (75B), outside strong purple (77B), with a large blotch of discrete of dark greenish yellow (152C) spots on the dorsal lobes; filaments light purple, anthers light grey; style pink, stigma cherry red. Calyx 7mm long, light green. Lvs elliptic, 140 × 60mm, rounded at base, pointed at apex, margins flat, glossy green above, hairless. Shrub 1.8 × 2.7m in 20 years. Early June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2011, p.29. Etymology: Imants is a Latvian male christian name

(r) **'Impromptu'**

cv. Elepidote rhododendron: 'Pinnacle', open-pollinated (s)

H: J.B. Brueckner (1981), G: J.B. Brueckner, N: C. Woodward (2010), REG: C. Woodward (2011)

Fls 10/conical truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, dark red (59A) and moderate purplish red (59C) in bud, the three dorsal lobes opening moderate purplish red (54A), with a spotted dark red (187A) blotch, the other lobes opening strong purplish red (64B); outside deep purplish red (71A), with a strong purplish red (64B) streak down the middle and along part of the edge; pistil and stamens white. Calyx miniscule, deep pink (160D). Lvs elliptic, 185 × 85mm, broadly acute at apex, rounded at base, the margins flat, dull and moderate olive green (147A and 137A) above, hairless. Shrub 1.5 × 1.5m in 29 years. Late May to early June. Nomenclatural standard: colour print of an electronic

image provided by the registrant (WSY). Etymology: the cross occurred without preparation or advance planning

(r) **'Italia 2011'**

cv. Elepidote rhododendron: parentage unknown, a plant found in the Parco Burcina, Biella, Italy

S: N. Furno (2000), N: N. Furno (2000),

REG: N. Furno (2011)

Fls 18–20/hemispherical truss, 25–32 × 58–65mm wide, with 5 moderately wavy-edged lobes, buds very pale purple (76D), with light purple (77C) margins, opening inside and out white (155A–B), with light purple (77C) margins; a very conspicuous dorsal blotch on three lobes, deep orange-yellow (163A) and dark orange (N163A) to moderate reddish orange (N172A); stamens 10, exerted from the corolla, anthers moderate yellowish pink (N170C), filaments white; pistil exceeding the corolla, but shorter than the stamens, style pale yellow (160D), with a moderate yellow-green (147B) apex, stigma moderate red (180B). Lvs lanceolate, 105–123 × 29–34mm, acute at base and apex, slightly twisted, matt moderate olive green (147A) above, hairless; petioles 18–22mm long. Shrub to at least 1.6 × 0.9m. May.

Etymology: named to commemorate the 150th anniversary of the unification of the Italian state

(r) **'Ivory Spice'**

cv. Elepidote rhododendron: 'Winter Spice' (s) × 'Muffy'

H: J. Barlup (2003), G: J. Barlup (2008), N: J. Barlup (2010), REG: J. Barlup (2011)

Fls 11/ball truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, white (155D) in bud, opening inside and out white (155D), with a large dorsal flare of deep purplish red (59B) spots extending c.35mm from the base; anthers dark red; pistil yellow. Calyx irregular, up to c.35mm long, white (155D), with deep purplish red (59B) spotting. Lvs elliptic, 115 × 50mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.5 × 0.9m in 7 years. Late May.

(r) **'Jane Bradish-Ellames'**

cv. Lepidote rhododendron: 'Tinkerbird' (s) × *lindleyi*

H: K.N.E. Cox (c. 2001), G: P.A. & K.N.E. Cox,

N: P.A. & K.N.E. Cox, I: Glendoick Gardens,

REG: K.N.E. Cox (2011)

Fls 3–4/open truss, more-or-less downfacing, broadly funnel-campanulate, 70 × 70mm, with 5 wavy-edged lobes, pale pink in bud, opening inside pure creamy white, outside also pure creamy white – although with some pink when first open; corolla green at base; scented. Calyx 8mm long, green. Pedicels long, green. Lvs broadly ovate, 70–90 × 30–40mm, acute at apex, margins flat, matt green above, ciliate, scaly. Shrub 0.6 × 1.9m in 10 years. April (under protection), May (grown outside).

Etymology: named for the wife of the registrant, who proposed the original cross

(a) **'Jessica Rose'**

cv. Deciduous azalea: hybrid derived from 'Homebush'

H: W.G.T. Hyde (1970s), G: W.G.T. Hyde (1970s), N: Mrs R. Legrand (2011), I: Millais Nurseries (2011), REG: Mrs R. Legrand (2011)
Fls double, 10–12/tight ball truss, trumpet-shaped, 44 × 25mm, with 15 wavy-edged lobes, moderate red (181A) in bud, opening inside strong red (51A); sweetly scented. Calyx irregular, 3–7mm long, strong yellow-green (144B). Lvs narrowly elliptic, 110 × 40mm, cuneate at base, acute at apex, margins decurved, matt green above, hairless. Shrub 2.1 × 1.6m in at least 30 years. May.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a granddaughter of the registrant

(r) **'Johann Darney'**

cv. *Elepidote rhododendron*: parentage unknown
G: C.O. Darney, S: C.O. Darney (2004), N: C.O. Darney (2011), REG: C.O. Darney (2011)
Fls 9/domed truss, broadly funnel-shaped, 65 × 40mm, with 6–7 wavy-edged lobes, light purple (close to 75A–76A) in bud, opening inside white, edged pale purplish pink (close to 65D), with a yellowish throat; outside white, tinged pale purplish pink; strongly scented. Calyx 3mm long, green. Lvs oblong, 200 × 90mm, broadly acute at apex, oblique at base, margins upcurved, moderate green with randomly distributed darker reddish markings above, hairless. Shrub 2.4 × 1.9m in 10 years. Mid-May.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the mother of the registrant

(r) **'John's Flare'**

cv. *Elepidote rhododendron*: 'Capistrano' (s) × 'Furnivall's Daughter'
H: J. Doppel (2004), G: J. Doppel (2008), N: J. Doppel (2011), REG: J. Doppel (2011)
Fs 12/ball truss, broadly funnel-shaped, c.50 × 65mm, with 5 wavy-edged lobes, light greenish yellow (8C) in bud, opening inside pale yellow-green (4D), with a large, prominent, vivid red (44A), spotted dorsal flare; outside pale yellow-green (4D). Lvs elliptic, 100 × 40mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and mid-green above, hairless. Shrub 0.36 × 0.4m in 7 years. Mid-May.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: derived in part from the registrant's christian name and from the most prominent character of the flowers

(r) **'Joseph Brueckner'**

cv. *Elepidote rhododendron*: *brachycarpum* (as subsp. *tigerstedtii*) (s) × *arboreum*
H: J.B. Brueckner (1978), G: J.B. Brueckner, N: M. Brueckner (2010), REG: C. Woodward (2011)
Fls 15–17/conical truss, funnel-campanulate, 43 × 60mm, with 5 very wavy-edged lobes, vivid red (57A) in bud, opening inside with a 6mm wide band of deep purplish pink (54B) at the lobe margins, then changing abruptly to deep purplish pink (66C) and blending to white at the base, with some randomly scattered deep red (53A) and strong red (53B) spots on the three dorsal lobes and with a surface depression at

the base of each lobe coloured pale purplish pink (56A); outside strong purplish red (54A) and deep purplish pink (54B) turning to light purplish pink (55C) towards the base, and finally to white. Calyx 2mm long, pale greenish yellow (1C). Lvs elliptic, 100–210 × 40–60mm, broadly acute at apex, rounded at base, the margins flat, dull and moderate olive green (147A) above, with a hint of 137A, hairless. Shrub 3.4 × 3.7m in 32 years. Mid-May.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the breeder, who raised a range of cold-hardy rhododendrons for the Canadian climate which retained aesthetic attributes of the parent plants

(r) **'Julistar'**

cv. *Elepidote rhododendron*: 'Solandra' × *fortunei* subsp. *discolor* × *catawbiense*
H: W. Wüstemeyer (1990), G: W. Wüstemeyer (1999), N: W. Wüstemeyer (2003), I: Baumschule Wüstemeyer (2011), REG: J. Westhoff (2011)
Fls up to 18/loose, rounded truss, funnel-shaped, 60 × 100mm, with 5–6 wavy-edged lobes, the centre of the flower sometimes with at least some partially petaloid stamens, buds moderate purplish red (58A), opening inside light purple to very light purple (75B/C), outside strong purplish pink (73B), with some darker patches; dorsal lobe with strong red (53C) spots. Calyx minute, light brown. Lvs oblanceolate, 170 × 70mm, margins flat to slightly upcurved, rounded at base, broadly acute at apex, hairless. Shrub 1.5 × 1.5m in 19 years. Late June to mid-July.
Nomenclatural standard: colour print derived from an electronic image provided by the registrant (WSY). Etymology: named from the star-like form of the flower and the late flowering period. First published in *Jahrbuch der Deutschen Rhododendron Gesellschaft*:45 (2003)

(r) **'Junistolz'**

cv. *Elepidote rhododendron*: 'Spätlese' (s) × 'San Sebastian'
H: H. Hesse (1997), G: H. Hesse (2003), N: H. Hesse, REG: H. Hesse (2011)
Fls 8–15/slightly loose, rounded truss, open funnel-shaped, 53–75 × 60–105mm, with 6–8 wavy-edged lobes, strong pink in bud, opening inside light pink, shading to whitish pink in the centre, with dark red markings at the base of the throat, outside purplish pink and partly cream-coloured. Calyx 1–3mm long, light green. Lvs elliptic to oblong-elliptic, 57–165 × 15–53mm, rounded at base, mucronate at apex, margins flat, slightly glossy and very dark green above, hairless. Shrub 1.1 × 1.3m in 14 years. Early to mid-June.
Etymology: named for its consistent flowering period and the large flower trusses ("Stolz" meaning pride or loftiness in German)

(r) **'Kamti'**

cv. *Elepidote rhododendron*: 'Rubicon' (s) × 'Hachmann's Polaris'
H: K.-H. Hübbers (1993), G: K.-H. Hübbers (1998), N: Mrs L. Hübbers (2007), REG: J. Westhoff (2011)
Fls c.14/lax, domed truss, campanulate, 40 × 50mm,

with 5 wavy-edged lobes, buds deep purplish pink (54B), opening inside white, flushed rose pink, outside pale orange-yellow (19D), flushed light pink (50D), with a dark red, bilobed blotch in the throat. Calyx 4mm long, green, rimmed with red. Lvs oblong, 80 × 35mm, margins flat, oblique at base, broadly acute and mucronate at apex, glossy green above, hairless. Compact slow-growing shrub, 0.9 × 0.9m in 10 years. Late April to early May.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: Kamti is a girl's name in Hindi

(v) **'Kane Boy'**

cv. Lepidote rhododendron: *konori* (s) × 'Doctor Hermann Sleumer' × (*macgregoriae* × *aurigeranum*)
H: R.A. Mitchell (pre 2008), G: R.A. Mitchell (2008),
N: R.A. Mitchell (2011), I: Pacific Island Nursery,
REG: R.A. Mitchell (2011)

Fls 7–10/conical truss, tubular funnel-shaped, 75 × 65mm, with 6 wavy-edged lobes, inside with a light greenish yellow (3D) centre, blending through yellowish pink to moderate purplish pink (54D) lobes; outside pink at base, shading to yellowish pink and finally to moderate purplish pink (54D) lobes; unmarked; filaments light yellow at base, shading into red, anthers brown; style yellowish green at base, shading into greenish pink, stigma light green; slightly scented. Lvs elliptic, 170 × 75mm, acuminate at apex, rounded at base, margins upcurved, glossy and moderate olive green (146A–147A) above, moderate yellow-green (146D) below, with tiny dark scales below along the midrib. Shrub with an open habit, 1.8 × 1.4m in 12 years. Once a year, in early April. Etymology: named for a male kitten belonging to the registrant. In Hawaii "Kane" is one way to identify the male sex, "Boy" is added for emphasis

(r) **'Kārlis Ulmanis'**

cv. Elepidote rhododendron: selection from *degronianum* (s)
H: R. Kondratovičs (1959), G: R. Kondratovičs (1965),
N: R. Kondratovičs (1988), REG: Rhododendron
Nursery Babīte, University of Latvia (2011)
Fls 8–9/slightly domed truss, open funnel-shaped,
70 × 70mm with 5 very wavy-edged lobes, buds deep
purplish pink (N57D), opening white (155C)
throughout, with discrete, strong yellowish green
(N144A) spots on the dorsal lobes; filaments white,
anthers light cream; style light pink, stigma pink. Calyx
1mm long, light green. Lvs elliptic, 140 × 50mm,
pointed at base and apex, margins flat to decurved,
glossy green above, with light brown hairs which age
to brown. Shrub 2.5 × 3.0m in 20 years. May.
Nomenclatural standard: colour print provided by
the registrant (WSY). First published in *Dārzs un
Drava*, December 2007, p.10. Etymology: named for
the former President of Latvia (1936–1940)

(r) **'Karl Naue'**

cv. Elepidote rhododendron: 'Hachmann's Junifeuer'
(s) × 'Aeschbacher's Rubin'
H: H. Hachmann (1994), G: H. Hachmann (1999),
N: Holger Hachmann (2007), I: Holger Hachmann

(2007), REG: M. Oppenrieder (2011)

Fls 13–20/dense, round-domed truss, funnel-shaped,
62–75mm wide, with 5 flat to wavy-edged lobes,
strong red (53B) throughout, with dark red (nearly
black) spots on the dorsal lobe; stamens tightly
grouped in the centre of the flower, filaments white;
style exerted, white, stigma red. Lvs elliptic, margins
flat to decurved; petioles reddish. Shrub 1.8 × 1.5m in
10 years. Mid-May to mid-June.

Nomenclatural standard: colour print provided by
the registrant (WSY). First published in Hachmann
cat. 2009–2013 (2009). Etymology: named for the
71st birthday of the head of the Dittersdorf
Baumschulen, Amtsberg OT Dittersdorf, Germany

(r) **'Keishun'**

cv. Elepidote rhododendron: open-pollinated
seedling from 'Hallelujah' (s)
G: H. Suzuki (2005), N: H. Suzuki (2005),
REG: H. Suzuki (2011)
Fls (8–)9(–10)/open truss, open funnel-shaped, 50 ×
70mm, with (5–)6(–7) smooth-edged lobes, deep pink
in bud, opening with a white line down the centre of
each pink lobe, throat white, with a conspicuous,
solid red basal blotch around the base of the throat,
breaking into short lines of spots on the dorsal side,
outside pink. Lvs oblong, 140 × 50mm, obtuse and
apiculate at apex, cuneate at base, margins flat, glossy
green above, hairless. Shrub with an open habit, 50 ×
30cm in 5 years (after grafting). Mid-May.
Nomenclatural standard: colour print provided by
the registrant (WSY). Seed collected by the registrant
in 1995

(r) **'Kiskadee'**

cv. Elepidote rhododendron: 'Doctor Ans Heyting'
(s) × 'Peucine'
H: K.-H. Hübbers (1999), G: K.-H. Hübbers (2005),
N: Mrs L. Hübbers (2011), REG: J. Westhoff (2011)
Fls up to 9/domed truss, open funnel-shaped, 40 ×
80mm, with 7(–8) wavy-edged lobes and with some
stamens (often several) partially petaloid, orange-
yellow in bud, opening inside brilliant yellow (13C),
with 8 lines of strong greenish yellow (151A) spots on
the dorsal lobe, outside brilliant to pale greenish
yellow (13C–D); stigma green. Calyx 2–3mm long,
yellowish green. Lvs elliptic, 130 × 50mm, margins
flat, wedge-shaped at base, broadly acute at apex,
glossy above when young, hairless. Shrub 1.1 × 1.9m
in 10 years. June.

Nomenclatural standard: colour print of an electronic
image provided by the raiser (WSY). Etymology: named
after a bird with yellow feathers found in Trinidad

(r) **'Kranenball'**

cv. Elepidote rhododendron: 'Rubicon' (s) ×
'Hachmann's Polaris'
H: K.-H. Hübbers (1993), G: K.-H. Hübbers (1998),
N: K.-H. Hübbers (2011), REG: J. Westhoff (2011)
Fls c.20/ball truss, campanulate, 40 × 30mm, with
5(–6) wavy-edged lobes, buds strong red (53B),
opening inside strong red (53D), shading to almost
white in the centre, outside strong red (53B/D), with
deep red (53A) spots on the three dorsal lobes. Calyx

2–20mm long, light green with a light pink rim or the larger lobes coloured like the corolla, but not conspicuous. Lvs elliptic, 90 × 30mm, margins flat, cuneate/rounded at base, broadly acute and mucronate at apex, matt green above, hairless. Shrub 0.9 × 0.9m in 10 years. Early May.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: named for the city of origin (Kranenberg) and the shape of the truss

(r) **‘Kranenflare’**

cv. *Elepidote rhododendron*: ‘Mondnacht’ (s) × ‘Hachmann’s Kabarett’

H: K.-H. Hübbers (2001), G: K.-H. Hübbers (2006), N: K.-H. Hübbers (2010), REG: J. Westhoff (2011) Fls c.12/flat-topped, domed truss, broadly funnel-shaped, 70 × 110mm, with 5 wavy-edged lobes, buds strong purplish red (63B), opening inside pale purplish pink (65C) with paler, creamy suffusions, outside light purplish pink (63D), the dorsal lobes with a conspicuous fan of dark red (59A) spots. Calyx 7mm long, reddish green. Lvs oblong, 130 × 55mm, margins slightly upcurved, cordate at base, broadly acute and mucronate at apex, matt green above, hairless. Shrub 1.4 × 1.3m in 10 years. Early to mid-May.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: named after the city of origin (Kranenberg) and the colour and marking of the flowers

(r) **‘Kranenkorn’**

cv. *Elepidote rhododendron*: ‘Peucine’ (s) × ‘Horizon Monarch’

H: K.-H. Hübbers (1993), G: K.-H. Hübbers (1999), N: K.-H. Hübbers (2011), REG: J. Westhoff (2011) Fls up to 12/ball truss, broadly funnel-shaped, 50 × 80mm, with 6 wavy-edged lobes, buds light greenish yellow (8B), with up to 8 deep pink (48A/B) lines along midveins, opening inside light yellow-green (2C), outside light greenish yellow (4C), dorsal lobes with a broad fan of dark red (59A) spots leading down to a blotch that encircles the base of the throat. Calyx 2–3mm long, light green, with a small red rim. Lvs oblong-elliptic, 150 × 65mm, margins flat, rounded at base, broadly acute and mucronate at apex, matt green above, hairless. Shrub 1.6 × 1.1m in 15 years. June.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: named after the city of origin (Kranenberg), the yellow colour which resembles grains of corn (‘korn’ in German) and the grain-like marking of the flowers

(r) **‘Lava Glow’**

cv. *Elepidote rhododendron*: ‘Odessa’ (s) × ‘Crimson Pippin’

H: K.-H. Hübbers (1992), G: K.-H. Hübbers (1998), N: K.-H. Hübbers (2011), REG: J. Westhoff (2011) Fls c.12/ball truss, open funnel-shaped, 45 × 70mm, with 5 wavy-edged lobes, buds strong red (46A), opening inside strong purplish red (63A), outside vivid red (46B), the centre of the dorsal lobe with a blotch of strong orange-yellow (22A) spots on a nearly white ground; not fading in the sun. Calyx 2mm

long, light green. Lvs elliptic-lanceolate, 110 × 30mm, margins slightly decurved, rounded at base, acute at apex, matt green above, hairless. Shrub 1.5 × 1.5m in 15 years. Mid-May.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: the colour of the flowers has the intensity of glowing lava

(r) **‘Lederstrumpf’**

cv. *Elepidote rhododendron*: ‘Peucine’ (s) × ‘Welse’

H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2006), N: K.-H. Hübbers (2011), REG: J. Westhoff (2011)

Fls c.8/loose truss, funnel-campanulate, 55 × 80mm, with 7 wavy-edged lobes, opening inside light greenish yellow (8B/C), with faint red stripes between the lobes, outside light yellow (12C), with faint red stripes and darker veins. Calyx 2mm long, green, with a brown rim. Lvs elliptic, 100 × 40mm, margins flat, rounded at base, broadly acute and mucronate at apex, hairless; very uniform and flat, leathery. Shrub 0.7 × 0.6m in 10 years. Mid-May.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: name suggested by the leathery texture of the leaves

(r) **‘Leitmotif’**

cv. *Elepidote rhododendron*: *aureum* var. *aureum* (s) × *brachycarpum* ‘Montanum’

H: J.B. Brueckner (pre 1994), G: J.B. Brueckner, N: M. Brueckner (2010), REG: C. Woodward (2011)

Fls 12/domed truss, funnel-campanulate, 40–45 × 50mm, with 5 very wavy-edged lobes, strong purplish red (64B) in bud, opening inside dark red (59A) and strong purplish red (59D) and fading to nearly white, the throat flushed light pink, with a blotch and spots of deep greenish yellow (153A) and strong greenish yellow (153B); outside with the upper half very pale purple (69B), the lower half deepening to strong purplish pink (63C) and with a vertical streak of moderate purplish pink (62B) reaching the tip of the lobe; on some lobes the stamen and pistil are enclosed at the base by two tiny, very pale purple (69B) petals. Calyx 2mm long, strong yellow-green (144B). Lvs orbicular, 100–110 × 60mm, obtuse at apex, cordate at base, with flat margins, glossy and moderate olive green (147A) above, hairless. Shrub with a dense habit, 1.8 × 3.1m in 41 years. Mid- to late May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the registrant indicates that the layer upon layer of identical flowers on this plant is reminiscent of recurring musical themes or leitmotifs

(r) **‘Lemon Charm’**

cv. *Elepidote rhododendron*: ‘Horizon Lakeside’ (s) × ‘Recital’

H: J. Barlup (1997), G: J. Barlup (2002), N: J. Barlup (2010), REG: J. Barlup (2011)

Fls 18/domed truss, broadly funnel-shaped, c.55 × 90mm, with 7 wavy-edged lobes, brilliant yellow (11A) in bud, the dorsal lobes opening light yellow (10C), blending to pale yellow (11D) at edges, with a spotted flare of strong red (53C) extending c.15mm on the central dorsal lobe and with additional, light,

random spotting of the same colour; other lobes opening pale greenish yellow (10D), fading to pale yellow (11D) at edges; outside opening light yellow (10C) blending to pale yellow (11D) on edges, the midribs with light yellow (10C) veining. Calyx c.15mm long, pale greenish yellow (10D), with strong red (53C) streaks. Lvs elliptic, 115 × 40mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.6m in 5 years. Early May.

(r) **'Leuchtpolster'**

cv. Elepidote rhododendron: 'Ovation' (s) × *forrestii* Repens Group

H: H. Hachmann (1990), G: H. Hachmann (1995), N: H. Hachmann (1997), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls 5–7/ loose, flatly rounded truss, campanulate, 43–55mm wide, with 5 flat-edged lobes, red throughout; filaments red, tightly clustered in the centre of the flower, shorter than the red style; conspicuous, very pale, floral bracts. Lvs obovate-elliptic, obtuse at apex, margins flat, dark green and glossy above, hairless. Low domed shrub, 0.4 × 0.8m in 10 years. Mid-April to early May.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **'Lil Clark'**

cv. Elepidote rhododendron: parentage unknown H: D.J. Dosser (2005), G: D.J. Dosser (2010), N: D.J. Dosser (2010), REG: D.J. Dosser (2011)

Fls 10/ball truss, broadly funnel-shaped, 110mm wide, with 5 wavy-edged lobes, buds moderate purplish red (185C), opening moderate red (179B) inside and out, with pale yellow (158A) spots on the three dorsal lobes; slightly scented. Calyx 10mm long, moderate purplish red (185C). Lvs oblanceolate, 120 × 40mm, acute at apex, cuneate at base, margins decurved, matt green above, hairless. Shrub 1.0 × 0.75m in 6 years. November (Australia).

Etymology: named for a friend of the registrant

(r) **'Limoncello'**

cv. Lepidote rhododendron: *keiskei* (s) × *fletcherianum*

H: J.B. Brueckner (1982), G: J.B. Brueckner, N: C. Woodward (2010), REG: C. Woodward (2011) Fls 7/lax truss, open funnel-shaped, 40–60 × 60mm, with 5 wavy-edged lobes, a blend of brilliant yellow-green (142B) and vivid yellow-green (154A) in bud, opening inside pale yellow-green (4D) to pale greenish yellow (2D) to light yellow-green (2C), with very small and faint brilliant yellow-green (154B) spotting; outside pale greenish yellow (2D) to light greenish yellow (1C). Calyx 1mm long, strong yellow-green (144B). Lvs elliptic, 40 × 20–30mm, broadly acute at apex, the margins flat, semi-glossy and moderate olive green (147A to 137A) above, hairless. Shrub with an open habit, 1.5 × 2.2m in 28 years. Mid-May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology:

named for the resemblance of the flower colour to that of the lemon-flavoured Italian liqueur of the same name

(a) **'Little Red'**

cv. Evergreen azalea: sport from 'Allotria'

H: H. Hachmann (1968), G: H. Hachmann (1993), N: H. Hachmann (1998), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls 2–3/truss, funnel-shaped, up to 50mm wide, with 5–7, flat-edged lobes, purplish rose pink inside and out; filaments and style purplish rose pink. Lvs elliptic, acute at apex, mid-green, with flat margins, hairless. Shrub 0.6 × 0.8m in 10 years. Late May to mid-June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(a) **MAJESTY**

Trade designation for 'Arismajesty'

(r) **'Malini'**

cv. Elepidote rhododendron: 'Mondnacht' (s) × 'Hachmann's Kabarett'

H: K.-H. Hübbers (2001), G: K.-H. Hübbers (2006), N: Mrs L. Hübbers (2011), REG: J. Westhoff (2011)

Fls c.7/loose, domed truss, broadly funnel-shaped, 60 × 100mm, with 7 wavy-edged lobes, inside cream (nearly pale orange-yellow, 19D), outside white, with dark red (59A) spots forming a fan on the three dorsal lobes; stigma green. Calyx 7mm long, green, with a red rim. Lvs elliptic, 140 × 65mm, margins wavy, rounded at base, obtuse and mucronate at apex, glossy and slightly rugose above, hairless. Shrub 0.6 × 0.7m in 5 years. Early to mid-May.

Somewhat resembles 'Sappho'

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: Hindi for a female gardener

(r) **MAMAMIA**

Trade designation for 'Hachmann's Mamamia'

(a) **'Maracas'**

cv. Deciduous azalea: 'Washington State Centennial' (s) × 'Arouca'

H: K.-H. Hübbers (1994), G: K.-H. Hübbers (1999), N: Mrs L. Hübbers (2010), REG: J. Westhoff (2011)

Fls at least 5/domed truss, tubular funnel-shaped (opening out widely at the mouth of the tube), 60 × 90mm, with 5(–6) very wavy-edged lobes, inside pale orange-yellow (19D), outside nearly white, with red veins and sporadic patches of moderate red (47A), with a vivid orange-yellow (23A) blotch on the dorsal lobe and half of each adjoining lobe; scented.

Calyx 7mm long (one lobe sometimes to 12mm), light green, translucent. Lvs obovate, 70 × 40mm, margins wavy, cuneate at base, broadly acute and mucronate at apex, with tiny white hairs along the margins and on the undersides; new leaves slightly tinged bronze. Early May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a seaside resort in Trinidad

(r) **'Martha Player'**

cv. *Elepidote rhododendron*: 'Mrs Sam' (s) × *hyperythrum*

H: L.M. Player (1998), G: L.M. Player (c. 2002), N: L.M. Player, REG: L.M. Player (2011)

Fls 20–26/ball truss, open funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, moderate purplish pink (62B) in bud, opening inside white, with a prominent double flare of deep purplish red (59B) spots on the dorsal lobe and a small green eye at the base; outside white. Calyx 3mm long, light yellow-green (145C). Lvs elliptic, 115 × 40mm, broadly acute at apex, cuneate at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.7 × 2.2m in 13 years. Mid-April. Heat tolerant (surviving well after a week of more of temperatures up to 38°C).

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the mother of the registrant

(r) **'Mary Watte'**

cv. *Elepidote rhododendron*: 'Miss B.L. Jones' (s) × 'Petra's Debut'

H: D.J. Dosser (2000), G: D.J. Dosser (2007), N: D.J. Dosser (2011), REG: D.J. Dosser (2011)

Fls 11/ball truss, broadly funnel-shaped, 75 × 50mm, with 5 wavy to strongly wavy-edged lobes, buds deep purplish pink (58D), opening strong purplish pink (63C) inside and out, with deep purplish red (59B) spots on the dorsal lobe; slightly scented. Lvs lanceolate, 125 × 42mm, margins decurved, matt green above, hairless. Shrub 1.5 × 1.0m in 11 years. November.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a friend of the registrant

(r) **MATADOR**

Trade designation for 'Hachmann's Matador'

(a) **'Mayo'**

cv. *Deciduous azalea*: 'Washington State Centennial' (s) × 'Arouca'

H: K.-H. Hübbers (1994), G: K.-H. Hübbers (1999), N: Mrs L. Hübbers (2010), REG: J. Westhoff (2011)
Fls c.10/domed truss, tubular funnel-shaped (opening out widely at the mouth of the tube), 70 × 100mm, with 5(–6) lobes, all the lobes wavy-edged but the dorsal lobe very wavy, buds vivid yellow (14A) at apex, brilliant yellow (14C) at base, with moderate red (47A) stripes, opening inside pale yellow (11C), with a light yellow (10C) dorsal lobe which is marked by a vivid orange-yellow (23A) blotch; outside pale yellow (11C), with a light yellow (10C) dorsal lobe and a vivid reddish orange (34B) stripe down the middle of each lobe; strongly scented. Calyx 6mm long, light green, translucent. Lvs obovate-elliptic, 80 × 40mm, margins flat, cuneate at base, broadly acute at apex, glossy green above, with tiny white hairs along the margins and along the midrib. Shrub 1.3 × 1.1m in 15 years. Early May.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology:

named for a village in Trinidad

(r) **METALLICA**

Trade designation for 'Hachmann's Metallica'

(r) **'Midnight Beauty'**

cv. *Elepidote rhododendron*: 'Frank Galsworthy' (s) × 'Jonathan Shaw'

H: Holger Hachmann (1996), G: Holger Hachmann (2001), N: Holger Hachmann (2002), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls up to 23/round, ball-like truss, broadly funnel-shaped, 95–110mm wide, with 5–6 wavy-edged, slightly recurved lobes, with a silky shine, nearly black in bud, opening dark wine red, with large black markings on the dorsal lobe and parts of the adjacent lobes; filaments wine-coloured, with white hairs at the base; style also wine-coloured, as long as the filaments. Lvs elliptic, margins flat, mid-green. Shrub 1.4 × 1.6m in 10 years. Mid- to late May.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **'Miks'**

cv. *Elepidote rhododendron*: 'Purple Splendour' (s) × 'Arnošt Silva-Tarouca'

H: R. Kondratovičs (1984), G: R. Kondratovičs (1991), N: R. Kondratovičs (2005), I: Rhododendron Nursery Babīte, University of Latvia (2005), REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 16/strongly domed truss, open funnel-shaped, 90 × 90mm, with 5–6 very wavy-edged lobes, buds strong purplish red (72A), opening inside light purple (77C), outside strong purple (77B), with discrete, dark greyish reddish brown (200A) spots on the dorsal lobes; filaments white, anthers white; style white, stigma cherry red. Calyx 3mm long, light green. Lvs elliptic, 170 × 50mm, rounded at base, pointed at apex, margins flat, glossy green above, hairless. Shrub 2.9 × 2.8m in 20 years. Late May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2011, p.28. Etymology: Miks is a Latvian male christian name

(r) **'Milderubin'**

cv. *Lepidote rhododendron*: *hirsutum* (s) × *ferrugineum*
G: Det Norske Arboret (1985), N: P.M. Jørgensen (2011), I: Gimle Planteskule (1990), REG: P.M. Jørgensen (2011)

Fls up to 15/flat truss, tubular-campanulate, 15 × 5mm, with 5 flat-edged lobes, crimson red in bud, opening rose red inside, deep rose outside; unmarked. Lvs narrowly elliptic, up to 40 × 5–8mm, tapering at base, pointed at apex, with rusty brown scales when young which become paler with age. Shrub 0.8 × 1.5m in 25 years, with a good habit and easily propagated and grown. Late June.

Nomenclatural standard: colour print provided by the registrant (WSY). A chance hybrid (1980) in the old Botanic Garden in Bergen, Norway, and subsequently moved to the Norske Arboret (Arboretet på Milde); the

name derived from the current location and the flower colour. Has been sold under the incorrect attribution *myrtifolium*. Has proved to be a good garden plant

(r) **'Mimi's Pride'**

cv. *Lepidote rhododendron*: PJM Group (s) × unknown
H: M.L. Wolin (1980), G: M.L. Wolin (1987),
N: M.L. Wolin (2011), REG: M.L. Wolin (2011)
Fls solitary, broadly funnel-shaped, 25 × 40mm, with 5 slightly wavy-edged lobes, deep reddish purple in bud, opening a rather vivid reddish purple throughout, slightly darker on the outside than the inside. Calyx c.5mm long, light green and tan. Lvs elliptic, 30 × 15mm, acute at apex, rounded at base, margins upcurved, dark green and glossy above, with tiny dark scales below along the midvein and on the pedicels. Shrub 1.4 × 1.8m in 30 years. Early April.

(r) **'Mone-hime'**

cv. *Elepidote rhododendron*: *brachycarpum*, double-flowered form × *degronianum* subsp. *degronianum* (s) × *arboreum*
H: H. Suzuki (1995), G: H. Suzuki (2004),
S: H. Suzuki (2004), N: H. Suzuki (2006),
REG: H. Suzuki (2011)
Fls double, (9–)10(–12)/loose, open-topped truss, open funnel-shaped, 40 × 50mm, with 5(–7) wavy-edged lobes in both the inner and outer whorls, red in bud, opening pinkish red throughout, unmarked. Lvs oblong, 120 × 35mm, margins flat, acute at apex, cuneate at base, with few hairs below. Shrub 0.4 × 0.3m in 4 years (after grafting). Mid-May. Nomenclatural standard: colour print provided by the registrant (WSY)

(r) **'Morning Melody'**

cv. *Elepidote rhododendron*: parentage unknown
H: unknown (1998), G: J. Doppel (2004),
N: J. Doppel (2009), REG: J. Doppel (2011)
Fls 14/ball truss, broadly funnel-shaped, c.60 × 75mm, with 5–7 wavy-edged lobes, vivid reddish orange (34B) in bud, opening inside light yellowish pink (26D), with a very large, prominent, spotted, strong red (41B) blotch, covering most of the central dorsal lobe, as well as portions of the adjacent lobes; outside light yellowish pink (26D). Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins downcurved, dull and mid-green above, hairless. Shrub 1.5 × 1.5m in 12 years. Early May.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: name coined to express the joy of seeing this plant in the morning

(r) **'Mrs June Sanderson'**

cv. *Elepidote rhododendron*: 'Miss B.L. Jones' (s) × 'Petra's Debut'
H: D.J. Dosser (2000), G: D.J. Dosser (2009),
N: D.J. Dosser (2011), REG: D.J. Dosser (2011)
Fls 16/conical truss, funnel-shaped, 100 × 55mm, with 5 wavy-edged lobes, strong purplish red (67A) in bud, opening strong purplish pink (67D), with a white flush, both inside and out; with deep purplish pink (67C) spots on the dorsal lobe; slightly scented. Lvs

oblanceolate, 150 × 50mm, acute at apex, truncate at base, margins decurved, matt green above, hairless. Shrub 1.5 × 0.75m in 10 years. November.
Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a friend of the registrant

(a) **'Naomi Legrand'**

cv. Evergreen azalea: parentage unknown
H: W.G.T. Hyde (1970s), G: W.G.T. Hyde (1970s),
N: Mrs R. Legrand (2011), I: Mrs R. Legrand,
REG: Mrs R. Legrand (2011)
Fls open trumpet-shaped, 51 × 25mm, with 6 wavy-edged lobes, moderate purplish red (59C) in bud, opening strong purplish red (59D) both inside and out, with prominent speckling on the dorsal lobes of dark red (59A). Calyx 6–10mm long, strong yellow-green (144A). Lvs oblanceolate, 47 × 15mm, narrowly cuneate at base, narrowly acute at apex, margins flat, matt green above, hairless. Shrub 0.9 × 0.9m in over 30 years. May.
Etymology: named to celebrate the 40th birthday of the registrant's daughter

(r) **'Narzissenbraut'**

cv. *Elepidote rhododendron*: 'Solandra' (s) × 'Edward Dunn'
H: W. Wüstemeyer (1995), G: W. Wüstemeyer (2002),
N: W. Wüstemeyer (2002), I: W. Wüstemeyer (2008),
REG: J. Westhoff (2011)
Fls up to 10/loose domed truss, broadly funnel-shaped, 55 × 90mm, with 6–7 very wavy-edged and slightly reflexed lobes, inside pale yellow to pale orange-yellow (19C/D), most stamens petaloid and strong yellowish pink (37B), with a patch of moderate yellow (160A) spots on the three dorsal lobes; outside light orange-yellow (19B), the lobes blushed pink; stigma prominent, light green. Calyx 7-lobed, 5 lobes about 10mm long, two up to 20mm long, light orange-yellow (19B). Lvs oblong-obovate, 130 × 50mm, margins flat, rounded at base, broadly acute at apex, matt green above, hairless. Shrub 1.2 × 1.5m in 10 years. Late June to mid-July. First published in *Jahrbuch der Deutsche Rhododendron Gesellschaft* 2003, p.47. Etymology: literally means "daffodil bride" and refers to the flowers having a certain resemblance to daffodils

(a) **NEGLIGÉ**

Trade designation for 'Hachneg'

(r) **'Night and Day'**

Syn. of 'Hachhans'

(r) **'Night Music'**

cv. *Elepidote rhododendron*: 'Jonathan Shaw' (s) × 'Black Adder'
H: J. Barlup (2003), G: J. Barlup (2009), N: J. Barlup (2010), REG: J. Barlup (2011)
Fls 18/ball truss, broadly funnel-shaped, 50 × 85mm, with 5 wavy-edged lobes, dark red (187A) in bud, opening inside with a moderate purplish red (59C) centre, c.15mm deep, and blending to dark red (59A) at the edges, with a dark red (187A) flare on the central dorsal lobe, this starting c.15mm from the base and

extending for 25mm; outside moderate purplish red (59C) at the base, blending to dark red (59A) at the edges, midribs dark red (187B); anthers close to white. Lvs elliptic, 125 × 50mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.8m in 6 years. Late May.

(r) '**Nimbu Pani**'

cv. *Elepidote rhododendron*: 'Doctor Ans Heyting' (s) × 'Pucine'

H: K.-H. Hübbers (1999), G: K.-H. Hübbers (2005), N: Mrs L. Hübbers (2011), REG: J. Westhoff (2011) Fls 6–8/domed truss, broadly funnel-shaped, 60 × 100mm, with 6(–7) wavy-edged lobes, buds brilliant greenish yellow (4A–5B), opening inside pale yellow-green (4D) at the margins, shading to light greenish yellow (4C) towards the throat, outside light greenish yellow (4B/C), strong to vivid red (46A/B) spots forming two bands in the lower quarter of the dorsal lobes and varying in intensity. Calyx 2–5mm long, yellow-green, with a red-dotted margin. Lvs broadly elliptic, 100 × 50mm, margins slightly decurved, cordate at base, obtuse and mucronate at apex, glossy green above when young, hairless. Shrub 1.1 × 1.0m in 10 years. June.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: in Hindi "nimbu" is the word for lemon and "pani" the word for water – the name is chosen because the flower colour looks like lemonade

(r) '**ORAKEL**'

Trade designation for 'Hachmann's Orakel'

(a) '**Orangenköpfchen**'

cv. *Deciduous azalea*: 'Orangeade' (s) × 'Narcissiflorum'

H: Holger Hachmann (1998), G: Holger Hachmann (2003), N: Holger Hachmann (2008), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011) Fls double, up to 18/dense, rounded truss, tubular funnel-shaped, 40–47mm wide, with up to 10 flat-edged lobes, inside strong orange (24A), with traces of salmon red, outside bright red, with orange stripes on the midveins; all the stamens petaloid, the outer lobes tapered and the flower as a whole star-shaped, the central lobes more irregular. Autumn leaf colour yellow-orange. Shrub 1.1 × 1.1m in 10 years. Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) '**Paradise Meadows**'

cv. *Elepidote rhododendron*: 'Haida Gold' (s) × 'Golden Star'

H: H.R. Wright (1991), G: H.R. Wright (1996), N: H.R. Wright (2011), REG: H.R. Wright (2011) Fls 12/domed truss, funnel-campanulate, c.50 × 70mm, with 6 wavy-edged lobes, close to strong to vivid purplish red (61B–C) in bud, opening inside light yellow; outside yellow, tinged with amber; light green spots on the dorsal lobe. Calyx 2.5mm long, green. Lvs oblong, 110 × 45mm, obtuse at apex,

oblique at base, margins flat, semi-glossy green above, hairless. Shrub 1.0 × 0.8m in 18 years. Early May. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the location known as Paradise Meadows can be seen from the registrant's garden

(r) '**Paul Dee**'

cv. *Elepidote rhododendron*: 'Calsap' (s) × Janet Blair' H: J. Doppel (2001), G: J. Doppel (2003), N: J. Doppel (2011), REG: J. Doppel (2011)

Fls 15/ball truss, broadly funnel-shaped, c.45 × 75mm, with 5 wavy-edged lobes, deep purplish pink (70C) in bud, opening inside white and shading to very light purple (75C) at margins, with a prominent, elongated, dark red (183A), spotted dorsal blotch, outside very light purple (75C). Lvs elliptic, 115 × 40mm, broadly acute at apex, rounded at base, margins lightly downcurved, semi-glossy and olive green above, hairless. Shrub with a dense habit, 0.8 × 1.1m in 10 years. Mid-May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a son of the registrant

(a) '**Pêrlite**'

cv. *Deciduous azalea*: a *Rustica azalea* (s) × 'Uguns'

H: R. Kondratovičs (1978), G: R. Kondratovičs (1985), N: R. Kondratovičs (2006), I: Rhododendron Nursery Babīte, University of Latvia (2006), REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls single, 7/slightly domed truss, tubular funnel-shaped, 45 × 45mm, with 5 wavy-edged lobes, buds white (155D), opening white (155D) throughout and with a bifurcated, vivid yellow (14A) blotch on the dorsal lobe; filaments white, anthers light brown; style light pink, stigma light green. Calyx 2mm long, green. Lvs oblanceolate, 70 × 25mm, pointed at base and apex, margins wavy, glossy green above, hairless. Shrub 1.4 × 1.0m in 10 years. Early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, December 2007, p.12

(r) '**Peter Chappell**'

cv. *Lepidote rhododendron*: probably *campylogynum* × *glaucophyllum*

R: P.G.G. Chappell (pre 2009), N: P.A. & K.N.E. Cox, REG: K.N.E. Cox (2011)

Fls 4–5/very open truss, tubular-campanulate, with 5 more-or-less smooth-edged lobes, buds deep reddish purple, opening reddish purple throughout, although slightly paler in the centre; unmarked; filaments pale purplish pink, anthers pale brown; style purplish pink, stigma pale yellowish at first, aging to purplish. Lvs lanceolate, 20–30 × 9–12mm, acute at apex, margins slightly decurved, slightly glossy green above, glaucous below, with widely spaced brown scales. Shrub 0.6 × 1.0m in 10–15 years. April to May.

(r) '**Pigalle**'

cv. *Lepidote rhododendron*: 'Sacko' (s) × *calostrotum* subsp. *riparioides*

H: H. Hachmann (1990), G: H. Hachmann (1994), N: H. Hachmann (1996), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 3–5/flat rounded truss, broadly funnel-shaped, up to 30mm wide, with 5 flat-edged lobes, dark blue violet throughout, with darker blue violet spots on the dorsal lobe; stamens violet; style purplish violet. Lvs elliptic, up to 20 × 10mm, margins flat, dark and glossy green above. Shrub 0.3 × 0.3m in 10 years. Early to mid-May.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(v) **‘Pink Organdy’**

cv. *Lepidote rhododendron*: parentage unknown
H: A. Rouse (c. 2006), G: A. Rouse (2010), N: A. Rouse (2011), REG: A. Rouse (2011)
Fls 2–3/open truss, tubular funnel-shaped, 38 × 50mm (tube 18mm, lobes 20mm long), with 5 wavy-edged lobes, inside light pink (49C) flushed to light pink (50D), outside similarly coloured; unmarked. Lvs narrowly elliptic, 25 × 10mm, obtuse to acute at apex, tapering at base, margins flat, with scales. Compact, much-branched shrub, 35 × 25cm in 5 years. August (Australia).
Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: name suggested by the flower colour and the flowers being reminiscent of the material organdy

(r) **‘Pink Titan’**

cv. *Elepidote rhododendron*: ‘Lem’s Monarch’ (s) × un-named seedling
H: K.-H. Hübbers (1990), G: K.-H. Hübbers (1997), N: K.-H. Hübbers (2010), REG: J. Westhoff (2011)
Fls c.12/conical truss, funnel-campanulate, 60 × 80mm, with 5 wavy-edged lobes, vivid purplish red (57D) in bud, opening inside light rosy purple, shading paler towards the centre and with darker veins (strong purplish pink, 62A), outside vivid purplish red (61C); two small strips of markings towards the base of the dorsal lobe. Calyx 2mm long, red, translucent. Lvs elliptic, 130 × 60mm, margins flat, rounded at base, broadly acute at apex, matt green above, hairless. Shrub 2.0 × 2.0m in 20 years. Mid-May.
Etymology: named for the size of the truss and the colour of the flowers

(r) **‘Protad’**

cv. *Elepidote rhododendron*: ‘Mondnacht’ (s) × ‘Hachmann’s Kabarett’
H: K.-H. Hübbers (2001), G: K.-H. Hübbers (2006), N: K.-H. Hübbers (2010), REG: J. Westhoff (2011)
Fls c.10/domed truss, broadly funnel-shaped, 50 × 90mm, with 6(–7) wavy-edged lobes, buds strong purplish red (60C/D) at apex, shading paler to base, opening inside pale yellow (11D) in the centre, the lobes pale purplish pink (62D) fading to almost white, outside pale yellow (11D) at the base, with very pale purple (69A) lobes; with a very conspicuous, large, dark red (187A–C) blotch on the three dorsal lobes which breaks up apically into spots; filaments white, anthers brown; style creamy white, stigma green. Calyx

2–6mm long, greenish red. Lvs obovate-elliptic, 130 × 70mm, margins slightly wavy, rounded at base, broadly acute and mucronate at apex, glossy green above, hairless. Shrub 0.6 × 0.7m in 5 years. Early May.
Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY)

(r) **‘Purple Embers’**

cv. *Elepidote rhododendron*: ‘Midnight Mystique’ (s) × ‘Jonathan Shaw’
H: J. Barlup (1999), G: J. Barlup (2005), N: J. Barlup (2010), REG: J. Barlup (2011)
Fls 16/ball truss, saucer-shaped, 50 × 90mm, with 5 wavy-edged lobes, dark red (187A) in bud, opening inside strong purple (77B), edged deep reddish purple (77A), with a dark red (163A) dorsal flare extending c.35mm from the base; outside deep reddish purple (77A), with dark purple (83A) midveins. Lvs elliptic, 125 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 11 years. Mid-May.

(r) **‘Raphaella’**

cv. *Elepidote rhododendron*: ‘Abraxas’ (s) × ‘Campanile’
H: H. Hachmann (1998), G: H. Hachmann (2003), N: H. Hachmann (2005), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 10–15/slightly loose, rounded truss, funnel-shaped, 70–87mm wide, with 5–6 wavy-edged lobes, opening with a broad margin of rich pink, the middle of the dorsal lobe and parts of the adjacent lobes rich yellow, the remainder of the lobe centres vanilla yellow, fading with age; greenish yellow spots on the darker yellow parts of the dorsal lobe; filaments creamy white; style yellow at base, purplish pink towards apex, stigma red. Lvs elliptic, margins flat, mid-glossy green above. Shrub 0.7 × 0.9m in 10 years. Mid-April to early May.

(a) **‘Rasma’**

cv. *Deciduous azalea*: parentage unknown
H: R. Kondratovičs (1989), G: R. Kondratovičs (1995), N: R. Kondratovičs (2008), REG: Rhododendron Nursery Babite, University of Latvia (2011)
Fls single, 7/slightly domed truss, tubular funnel-shaped, 70 × 70mm, with 5 wavy-edged lobes, buds deep yellowish pink (43C), opening inside pale purplish pink (65D), outside strong purplish pink (62A), with a conspicuous blotch of strong orange-yellow (17A) on the dorsal lobe (more-or-less covering the lobe in most flowers and the spots coalescing); filaments light pink, anthers white; style light pink, stigma green; slightly scented. Calyx 2mm long, green. Lvs elliptic, 120 × 35mm, pointed at base and apex, margins wavy, glossy green above, hairless. Shrub 2.0 × 1.5m in 10 years. Late May to early June.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2010, p.35. Etymology: Rasma is a Latvian female christian name

(r) **‘Raspberry Mist’**

cv. *Elepidote rhododendron*: ‘Violet Mist’ (s) × ‘Black Adder’

H: J. Barlup (2003), G: J. Barlup (2008), N: J. Barlup (2011), REG: J. Barlup (2011)

Fls 17/ball truss, broadly funnel-shaped, 45 × 65mm, with 5 very wavy-edged lobes, deep purplish red (71A) in bud, opening inside deep purplish pink (72D) in the centre and darkening through strong reddish purple (72B) to strong purplish red (72A) at the edges, and having a densely spotted, dorsal flare of dark greyish reddish brown (200A), starting c.10mm from the base and extending c.25mm; outside deep purplish pink (72D), with midveins and outer edges strong purplish red (72A). Lvs elliptic, 115 × 45mm, broadly acute at apex, rounded at base, with flat margins, dull and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.9m in 7 years. Early June.

(r) **'René de Milleville'**

cv. *Elepidote rhododendron*: 'Kilimanjaro' × ('Fred Wynniatt' × 'Ring of Fire') (s) × 'Red Riding Hood' H: M. Colombel (2002), G: M. Colombel (2008), N: M. Colombel (2011), REG: M. Colombel (2011) Fls 8–10/domed truss, broadly funnel-campanulate, 55–65 × 70–90mm, with 7 wavy-edged lobes, buds vivid purplish red (61C) at apex to vivid red (57A), opening inside light purplish pink (62C) at margins, shading to moderate orange-yellow (164C) and then deep red (185A) in the throat, outside light purplish pink (62C) at the margins, shading to moderate orange (168C); numerous small spots on the dorsal lobes (orange on the dorsal lobes, darker in the throat); very narrow moderate red (180A) stripes, both inside and outside, down the midveins; filaments mostly yellow, but ruby at the base; style red, nearly straight; ovary with white glands. Calyx 15–20mm long, moderate orange-yellow (164B), with narrow, moderate red (180A) stripes. Lvs elliptic, 115–130 × 50–55mm, rounded at base, obtuse at apex, margins flat, but leaf blade V-shaped and slightly reflexed in long axis, matt green above, hairless. Shrub 0.9 × 1.5m in 9 years. Early to mid-May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a man who lived for many years in Nepal, sent back seeds and wrote a book about the rhododendrons of Nepal

(r) **'Rhododendronpark Graal-Müritz'**

cv. *Elepidote rhododendron*: 'Hachmann's Kabarett' (s) × 'Peter Alan'

H: Holger Hachmann (1994), G: Holger Hachmann (1999), N: Holger Hachmann (2006), I: Holger Hachmann (2006), REG: M. Oppenrieder (2011) Fls 8–16/dense rounded truss, funnel-shaped, 75–90mm wide, with 5–6 very wavy-edged lobes, deep purple violet in bud, opening inside strong purple (83C), aging to strong purple (82B), with a large dark violet to nearly black blotch on the dorsal lobes and parts of the adjacent lobes, which later becomes purplish; filaments bright lilac, anthers white; apex of style and the stigma slightly purplish. Lvs elliptic, margins flat, dark glossy green above. Shrub 1.4 × 1.6m in 10 years. Late May to mid-June. Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009). Etymology: named for the

Rhododendronpark in Graal-Müritz, Mecklenburg, Germany

(a) **'RLH1-1P2'**

cv. Evergreen azalea: un-named selection (RLH-1900-RP) (s) × 'September Song'

H: R.H. & L.J. Head (1996), G: R.H. & L.J. Head, S: R.H. & L.J. Head (2000), REG: Greenleaf Nursery Company (2011)

Fls single to semi-double, with 2–4 fls/bud and 1–9 buds/shoot, 35–45 × 70–85mm, with a 5–6-lobed, outer corolla, with very broad, undulate and ruffled lobes and sometimes with 1–5 narrower, petaloid stamens, when fully opened near strong purplish red (61B), with spots and streaks of near deep red (60A); becoming closer to strong purplish red (60D) with age; similarly coloured outside; stamens 6–10, filaments near deep red (60A), anthers near dark greyish red (N186C); style near deep red (60A), stigma near dark purplish red (N79A). Calyx 5–6-lobed, 12–17mm long, ciliate and silky hairy, close to strong yellow-green (144A). Lvs 'obelliptic', 38–63 × 16–29mm, subacute and mucronulate at apex, cuneate to attenuate at base, margins revolute, ciliate, close to dark yellowish green (139A), with paler veins; petioles c.6–9mm long, green. Compact, upward-spreading, globose shrub, 30–45 × 25–35cm in 2 years. April to May and again July to October. Trade designation: BLOOM-A-THON RED AZALEA

(a) **'RLH1-2P8'**

cv. Evergreen azalea: un-named selection (RLH-1900-RP) (s) × 'Watchet'

H: R.H. & L.J. Head (1996), G: R.H. & L.J. Head, S: R.H. & L.J. Head (2000), REG: Greenleaf Nursery Company (2011)

Fls double, 3–6/cluster, broadly funnel-shaped, 50–65 × 90–100mm, with a 5-lobed outer corolla and a central whorl of 5–10 lobes, the lobes narrowly and strongly wavy-edged, the outer whorl opening inside near deep purplish pink (61D), with spots and speckles of near strong purplish red (61B), maturing to near deep purplish pink (N57D), with spots and speckles of strong purplish red (61B); outside opening near deep purplish pink (N57D) and remaining this colour; stamens 5–10, filaments near deep purplish pink (61D), anthers near brownish orange (N167A); style near strong purplish red (63A), stigma near deep purplish red (59B). Calyx 5–6-lobed, 4–4.5mm long, silky pubescent, ciliate, green. Lvs 'obelliptic' to elliptic, c.20–50 × 10–29mm, subacute to rounded and mucronulate at apex, cuneate to attenuate at base, margins slightly revolute, ciliate, hairy, darker than dark green (136A) above when mature; petioles 5–8mm long. Shrub 30–45 × 35–40cm in 2 years. April to May and July to October.

Trade designation: BLOOM-A-THON PINK DOUBLE AZALEA

(a) **'RLH1-3P3'**

cv. Evergreen azalea: un-named selection (RLH-1800-WS) (s) × 'September Song'

H: R.H. & L.J. Head (1996), G: R.H. & L.J. Head, S: R.H. & L.J. Head (2000), REG: Greenleaf Nursery

Company (2011)

Fls single, 2–3fls /bud with 1–3 buds /shoot, broadly funnel-shaped, 30–35 × 60–75mm, with 5 very broad, undulate and ruffled lobes, when fully open white (N155D) throughout, with spots and speckles close to strong yellow-green (N144C); stamens c.10, filaments white (N155C), anthers moderate orange-yellow (165C); style white (N155C), stigma near strong yellow-green (143C). Calyx 5-lobed, 2.8–4mm long, strong yellow-green (N144D). Lvs ‘obelliptic’, 10–32 × 4–15mm, rounded or obtuse and mucronate at apex, cuneate or attenuate at base, ciliate, hairy, dark yellowish green (139A) above, veins close to strong yellow-green (144A); petioles 3–5.5mm long, green. Compact, upwardly spreading, globose-shaped shrub, 25 × 35cm in 2 years. April to May and again July to October.

Trade designation: BLOOM-A-THON WHITE AZALEA

(a) ‘RLH1-4P19’

cv. Evergreen azalea: Un-named selection (RLH-1900-RP) (s) × un-named selection (RLH-19-PAF) H: R.H. & L.J. Head (1996), G: R.H. & L.J. Head, S: R.H. & L.J. Head (2000), REG: Greenleaf Nursery Company (2011)

Fls single, 3–6/cluster, broadly funnel-shaped, 40–45 × 70–80mm, with 5 wavy-edged lobes, opening inside at first near deep purple (N81A), with spots and speckles close to deep purplish red (71A), outside close to deep purple (N81A); maturing inside to strong reddish purple (N78A) and then N78B, with spots and speckles close to deep purplish red (71A), outside near strong reddish purple (N78A) becoming N78B; rarely with one petaloid present; stamens c.10, filaments near vivid purplish red (57B), anthers dark greyish red (N186C); style close to strong purplish red (61B). Calyx c.15–20mm long, 5-lobed, ciliate, hairy, green, but with a central strong purplish red (60D) blotch and streaks. Lvs oblanceolate to elliptic, 30–65 × 10–25mm, subacute and mucronulate at apex, cuneate to attenuate at base, margin slightly revolute, ciliate, hairy above and below, near dark yellowish green (139A) above when mature; petioles c.6–10.5mm long, green. Shrub 35–50 × 30–40cm in 2 years. April to May and again July to October.

Trade designation: BLOOM-A-THON LAVENDER AZALEA

(r) ‘Robis’

cv. Elepidote rhododendron: open-pollinated seedling from ‘Catawbiense Grandiflorum’ H: R. Kondratovičs (1967), G: R. Kondratovičs (1973), N: R. Kondratovičs (2006), REG: Rhododendron Nursery Babite, University of Latvia (2011) Fls 14–16/slightly domed truss, funnel-shaped, 80 × 80mm, with 5 wavy-edged lobes, buds deep purplish red (59B), opening inside strong purplish red (60D), outside strong purplish red (60D), with relatively inconspicuous, discrete greyish purple (N77A) spots on the dorsal lobes; filaments pink-purple, anthers white; style pink-purple, stigma dark purple. Calyx 8mm long, very light pink. Lvs elliptic, 140 × 65mm, rounded at base, pointed at apex, margins flat, glossy green above, hairless. Shrub 2.5 × 3.0m in 20 years.

Late May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, December 2007, p.10. Etymology: Robis is a Latvian male christian name

(r) ‘Rosandra’

cv. Elepidote rhododendron: ‘Peucine’ (s) × ‘Horizon Monarch’

H: K.-H. Hübbers (1993), G: K.-H. Hübbers (2000), N: W. Wüstemeyer (2011), REG: J. Westhoff (2011) Fls up to 15/conical truss, open funnel-shaped, 50 × 70mm, with 5 wavy-edged lobes, buds strong purplish red (58B/C), opening inside moderate to light purplish pink (62B/C), with a very pale yellow band down the centre of each lobe, outside strong purplish red (58B), the dorsal throat with numerous, faint, strong greenish yellow (151B) markings; stigma green. Calyx 3–5mm long, light green. Lvs oblong to narrowly oblong, 100 × 25–35mm, margins flat to slightly decurved, rounded at base, broadly acute at apex, matt green above, hairless. Shrub 1.6 × 1.4m in 10 years.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: a combination of the word rose and the female christian name Sandra; the name of a valley in Slovenia

(r) ‘Rose Dust’

cv. Elepidote rhododendron: ‘Winter Spice’ (s) × ‘Muffy’

H: J. Barlup (2003), G: J. Barlup (2007), N: J. Barlup (2011), REG: J. Barlup (2011)

Fls 15/ball truss, broadly funnel-shaped, 50 × 80mm, with 5 wavy-edged lobes, strong purplish red (67A) in bud, opening inside very pale purple (69B), with lobe edges deep purplish pink (70C), and with strong purplish red (71C) nectar pouches on all lobes extending c.10–15mm from the base, and strong purplish red (71C) spotting extending c.30mm on the dorsal lobe; outside very pale purple (69B), the lobes edged with deep purplish pink (70C) and with lines of the same colour along the midribs. Lvs elliptic, 125 × 55mm, broadly acute at apex, rounded at base, the margins flat, dull and moderate olive green (147A) above, hairless. Shrub with a dense habit, 0.6 × 0.9m in 7 years. Mid-May.

(a) ‘Rosenköpfchen’

cv. Deciduous azalea: ‘Nabucco’ (s) × ‘Jack A. Sand’

H: Holger Hachmann (1998), G: Holger Hachmann (2003), N: Holger Hachmann (2008), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls double, up to 14/slightly loose, flat rounded truss, tubular funnel-shaped, up to 75mm wide, with 10 wavy-edged lobes, purplish red in bud, opening inside bright purplish rose pink, with a yellow-orange blotch at the base of the dorsal lobe; all stamens petaloid, the outer lobes tapered and slightly reflexed, inner lobes more irregular and partly very wavy or twisted. Shrub 1.1 × 1.1m in 10 years. Mid-May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **'Rosy Ray'**

cv. Elepidote rhododendron: 'Rosilac' (s) × 'Peucine' × 'Desert Gold'
H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2008), N: K.-H. Hübbers (2011), REG: J. Westhoff (2011)
Fls c.16/ball truss, broadly funnel-shaped, 60 × 90mm, with 7 wavy-edged lobes, buds deep yellowish pink (39B), opening inside light greenish yellow (8C), shading to white on the lobes, outside light greenish yellow (8C), with faint apricot veins, with 7 strips of strong red (53B) spots arranged around the base of the throat (alternating with the lobes), producing a star-like pattern. Calyx 3mm long, green, with reddish tips. Lvs oblong-elliptic, 130 × 60mm, margins slightly decurved, rounded at base, obtuse and mucronate at apex, matt green above, hairless. Shrub 1.1 × 0.8m in 10 years. Early to mid-May. Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY). Etymology: the name describes the markings on the flowers

(r) **'Rudite'**

cv. Elepidote rhododendron: an open-pollinated seedling from seedling H-75/17 (itself an open-pollinated seedling from *smirnowii*)
H: R. Kondratovičs (1982), G: R. Kondratovičs (1988), N: R. Kondratovičs (1993), REG: Rhododendron Nursery Babīte, University of Latvia (2011)
Fls 10/slightly domed truss, open funnel-shaped, 90 × 90mm, with 5 narrowly but very wavy-edged lobes, buds deep purplish pink (N57D), opening inside white (155D), outside strong purplish pink (68B), with some fairly inconspicuous, discrete spotting of strong yellow (N144B) on the dorsal lobes; filaments light pink, anthers grey pink; style light pink, stigma red. Calyx 2mm long, light pink. Lvs elliptic, 160 × 70mm, pointed at base and apex, margins flat to decurved, glossy green above, with white hairs when young, which age to light grey. Shrub 2.5 × 3.0m in 20 years. Early June.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, December 2007. p.11. Etymology: Rudite is a Latvian female christian name

(r) **'Rudolf Heinemann'**

cv. Elepidote rhododendron: 'Belami' (s) × 'Goldiesel'
H: H. Hachmann (1994), G: H. Hachmann (1999), N: Holger Hachmann (2006), I: Holger Hachmann (2007), REG: M. Oppenrieder (2011)
Fls 8–16/dense, rounded truss, funnel-shaped, 63–80mm wide, with 5 flat-edged lobes, salmon pink in bud, opening creamy yellow, with a rich cherry red blotch at the base of the dorsal side, fanning out on to the lobe as red spotting; ovary greenish, style yellow. Lvs oblong-elliptic, margins flat, dark glossy green above. Shrub 0.8 × 0.8m in 10 years. Mid-April to mid-May.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009). Etymology: named for the 70th birthday for the head of the Grönfingers garden centre in Rostock, Germany

(r) **'Saralynn'**

cv. Elepidote rhododendron: 'Meela' (s) × 'Christina Dee'
H: J. Barlup (2004), G: J. Barlup (2008), N: J. Barlup (2010), REG: J. Barlup (2011)
Fls 14/ball truss, broadly funnel-shaped, 60 × 90mm, with 5 very wavy-edged lobes, strong red (51A) in bud, opening inside pale orange-yellow (19D), with pale yellow (19C) on the dorsal lobe, the outer edges changing to very pale purple (69C), with deep purplish red (59B) spotting on the dorsal lobes extending c.45mm from the base; outside pale yellow (19C), fading to very pale purple (69C) on the edges. Lvs elliptic, 115 × 60mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.75 × 0.9m in 6 years. Late May.
Etymology: named for a friend of the registrant's wife

(r) **'Sariska'**

cv. Elepidote rhododendron: 'Kalamaika' × 'Perlina' (s) × 'Goldstück'
H: H. Hachmann (1995), G: H. Hachmann (2000), N: H. Hachmann (2001), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 12–21/dense, rounded truss, funnel-shaped, 70–80mm wide, with 5 very wavy-edged lobes, buds salmon pinkish cream, with thin salmon pink margins and central vein, opening inside creamy yellow, with thin salmon pink margins and central vein; with a large red-brown, spotted blotch on the dorsal lobes and parts of the adjacent lobes; filaments white; style various shades of green. Lvs elliptic, margins wavy, mid-glossy green above, with yellow rims. Shrub 1.2 × 1.1m in 10 years. Early May to early June.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **SASKIA**

Trade designation for 'Hachmann's Saskia'

(r) **'Sasonade'**

cv. Elepidote rhododendron: 'Fantastica' (s) × *dichroanthum* subsp. *scyphocalyx* Herpesticum Group
H: Holger Hachmann (1993), G: Holger Hachmann (1998), N: Holger Hachmann (2000), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls 9–12/ loose, flat rounded truss, campanulate, up to 40mm wide, with 5, wavy-edged lobes, inside vivid reddish orange (44C) at the margins, otherwise vivid orange (16A), with vivid red (45B) spots towards the base; outside bright red, with areas of bright orange; filaments very short, clustered in the middle of the flower, salmon pink with white hairs on the basal part, anthers dark brown; style bright orange, stigma greenish. Lvs broadly elliptic, margins decurved, dark glossy green above, with a thin hairy indumentum below. Shrub 0.7 × 0.9m in 10 years. Late April to late May.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2011 (2009)

(a) **‘Schneeköpfchen’**

cv. Deciduous azalea: ‘Oxydol’ (s) × ‘Narcissiflorum’
H: Holger Hachmann (1998), G: Holger Hachmann (2003), N: Holger Hachmann (2008), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls double, 8–13/dense, flat rounded truss, tubular funnel-shaped, 38–45mm wide, with mostly 10 wavy-edged lobes, greenish creamy white in bud, opening white; all stamens petaloid, the lobes tapered and recurved, the flower star-shaped; intensely sweet-scented. Shrub 1.3 × 1.3m in 10 years. May.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **‘Schwitter’s Rubin’**

Syn. of ‘Aeschbacher’s Rubin’

(r) **‘Sen-no-izayoi’**

cv. Elepidote rhododendron: *brachycarpum*, double-flowered form (s) × *degronianum* subsp. *heptamerum* var. *kyomaruense*, double-flowered form
H: S. Kanke (1995), G: S. Kanke (2004), N: S. Kanke (2011), REG: S. Kanke (2011)
Fls double, (10–)11(–13)/ hemispherical truss, open funnel-shaped, 40 × 60mm, with (5–)6 wavy-edged lobes in each whorl (an average of 6/whorl) and with irregularly petaloid stamens, pink/white in bud, opening inside very pale pink, outside deeper pink on the tube, much paler on the lobes; greenish dorsal spotting. Lvs oblong, 100 × 35mm, cuneate to narrowly rounded at base, broadly acute to obtuse and apiculate at apex, margins decurved, glossy above, with a hairy indumentum below which is white when young, aging to brown. Shrub 50 × 30cm in 6 years (after grafting). Mid-May.
Nomenclatural standard: colour print provided by the registrant (WSY)

(r) **‘Seraglio’**

cv. Elepidote rhododendron: ‘Evening Glow’ (s) × ‘Fred Winter’ × ‘Fashion Plate’
H: D. Mac Mullan (2005), G: D. Mac Mullan, N: D Mac Mullan (2010), REG: D. Mac Mullan (2011)
Fls in a conical truss, broadly funnel-shaped, c.75 × c.75mm, with 6 very wavy-edged lobes, orange/yellow in bud, opening inside mid-yellow at base, merging into paler yellow and then dark ivory at the margins, with three rays of orange spots on the dorsal lobe extending from base to mid-lobe; outside similar but unmarked. Calyx 20mm long, light to mid-yellow. Lvs elliptic, 65 × 30mm, broadly acute at apex, rounded at base, with flat margins, semi-glossy and mid-green above, hairless. Shrub 1.8 × 1.1m in 5 years. Early to mid-season.

(r) **‘Sharon Rose Smith’**

cv. Elepidote rhododendron: parentage unknown
H: P. Smith (c. 2000), G: P. Smith (2006), N: P. Smith, REG: P. Smith (2011)
Fls 12/ball truss, open funnel-shaped, c.70 × 100mm, with 7 wavy-edged lobes, strong red (53D) in bud, opening inside brilliant yellow (9C) to pale greenish yellow (9D) in the centre, with light yellowish pink

(36C) edges on all lobes and sparse vivid yellow (14A) spotting on the three dorsal lobes; outside light orange-yellow (16C) at base, shading to light orange-yellow (16B) at outer edges; slightly scented. Pedicels red. Calyx 3–6mm long, green, with red on some edges. Lvs oblong, 150 × 65mm, obtuse at apex, rounded at base, margins flat, dull and moderate olive green (146A) above, hairless. Dense shrub, 1.8 × 1.5m in 10 years. Early April.

Etymology: named for the wife of the registrant

(r) **‘Silvia Marie’**

cv. Elepidote rhododendron: ‘Janet Blair’ (s) × ‘Purple Lace’
H: J. Doppel (2001), G: J. Doppel (2005), N: J. Doppel (2011), REG: J. Doppel (2011)
Fls 20/ball truss, broadly funnel-shaped, c.50 × 75mm, with 5 wavy-edged lobes, vivid reddish purple (74A) in bud, opening inside light reddish purple (74C) at margins and fading lighter towards midveins and base, with a dorsal blotch of small, discrete, strong greenish yellow spots; outside light reddish purple (74C). Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and mid-green above, hairless. Shrub 0.9 × 0.9m in 10 years. Late May.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the registrant’s daughter-in-law

(r) **‘Simple Jane’**

cv. Elepidote rhododendron: ‘Admiral Vanessa’ (s) × ‘Welse’
H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2009), N: K.-H. Hübbers (2011), REG: J. Westhoff (2011)
Fls 10–12/domed truss, open funnel-shaped, 50 × 90mm, with 6 wavy-edged lobes, opening inside strong pink (50C), with pale yellow (8D) on the dorsal lobe, outside strong pink (50C), with darker veins (strong red, 51A); deep orange-yellow (163A) spots on two-thirds of the dorsal lobe. Calyx 2mm long, light green. Lvs oblong-elliptic, 100 × 40mm, margins flat, cuneate at base, broadly acute at apex, matt green above, hairless. Shrub 0.9 × 0.8m in 10 years. June.

(r) **‘Singing Sun’**

cv. Elepidote rhododendron: ‘Sun Blush’ (s) × ‘Goldprinz’
H: J. Barlup (2002), G: J. Barlup (2007), N: J. Barlup (2010), REG: J. Barlup (2011)
Fls 18/domed truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, light yellow (10C) in bud, opening inside and out pale yellow (11C), with a prominent dorsal flare of deep red (185A) spots extending c.45mm from the base and two rays extending c.15mm; in addition there are three small rays of the same colour on the lower lobes, extending c.10mm from the base. Lvs elliptic, 125 × 50mm, broadly acute at apex, rounded at base, margins downcurved, glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 8 years. Mid-May.

(a) **‘Siparia’**

cv. Deciduous azalea: ‘Washington State Centennial’ (s)

× ‘Arouca’

H: K.-H. Hübbers (1994), G: K.-H. Hübbers (1999), N: Mrs L. Hübbers (2010), REG: J. Westhoff (2011)
Fls up to 9/loose domed truss, tubular-funnel-shaped (opening out widely at the mouth of the tube), 60 × 90mm, with 5–6 very wavy-edged lobes (and with some petaloid stamens), strong reddish orange (31A) in bud, opening light yellowish pink (36B/C) to pale yellowish pink (27D), with darker red veins; outside white, with deep pink (48B) patterns; dorsal lobes (and some of the petaloid stamens) with a large vivid orange-yellow (23A) blotch; filaments and style pink; prominent light green stigma; scented. Calyx 6mm long, light green, translucent. Lvs elliptic, 90 × 40mm, margins flat, cuneate at base, broadly acute and mucronate at apex, with tiny white hairs on the margins and midribs. Shrub 1.1 × 0.7m in 10 years. Early May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a town in Trinidad

(r) ‘Sonlal’

cv. Elepidote rhododendron: ‘Rosilac’ (s) × ‘Peucine’ × ‘Desert Gold’
H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2006), N: Mrs L. Hübbers (2010), REG: J. Westhoff (2011)
Fls c.16/loose domed to flat truss, broadly funnel-shaped, 50 × 70mm, with 7 wavy-edged lobes, vivid reddish orange (34B) in bud, opening inside light greenish yellow (8B), with pinkish red stripes running down from the sinuses; outside light greenish yellow (8B), with pinkish red stripes down the midveins; stigma green; colour sun-resistant; slightly scented. Calyx 5mm long, greenish. Lvs broadly elliptic, 110 × 60mm, margins decurved, cordate at base, obtuse and mucronate at apex, matt green above, hairless. Shrub 1.1 × 0.8m in 10 years. Early to mid-May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the name means golden treasure in Hindi

(a) ‘Sonnenköpfchen’

cv. Deciduous azalea: ‘Orangeade’ (s) × ‘Narcissiflorum’
H: Holger Hachmann (1998), G: Holger Hachmann (2003), N: Holger Hachmann (2008), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)
Fls double, 18–21/dense, flat rounded truss, tubular funnel-shaped, 45–51mm wide, with 10 flat-edged lobes, greenish yellow in bud, opening inside rich yellow, with a golden yellow dorsal blotch and a trace of pink on the outer ring of lobes when first open; all stamens petaloid, the lobes all tapered and slightly recurved, the flower star-shaped; with an intense sweet scent. Shrub 1.3 × 1.3m in 10 years. May.
Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) ‘Stead’s Best’

cv. Lepidote rhododendron: *nuttallii* (s) × *lindleyi*
H: E.F. Stead (pre-1950), G: E.F. Stead, N: E.F. Stead, REG: University of Canterbury (New Zealand) (2011)

Fls 4/open truss, broadly funnel-campanulate, 140 × 120mm, with 5 wavy-edged lobes, buds red at tips, shading through green to cream at the base, opening inside white, with a yellow throat which is darker on the dorsal side, outside white; scented. Calyx 25mm long, green. Lvs oblong, 160 × 60mm, margins flat, narrowly rounded and obtuse at apex, cuneate at base, with red scales below when young, which later become red-brown on a silvery surface. Shrub 4.0 × 2.5m in 60 years, with papery bark. November (New Zealand). Nomenclatural standard: colour print provided by registrant (WSY)

(r) ‘Tālavija’

cv. Elepidote rhododendron: open-pollinated seedling from *smirnowii* – (H-75/17) (s) × ‘Marka’
H: R. Kondratovičs (1985), G: R. Kondratovičs (1991), N: R. Kondratovičs (2008), REG: Rhododendron Nursery Babīte, University of Latvia (2011)
Fls 8–10/strongly domed truss, open funnel-shaped, 80 × 80mm, with 5 wavy-edged lobes, buds strong purplish red (63B), opening inside moderate purplish pink (68C), outside deep purplish pink (68A), with conspicuous dorsal markings of dark red (59A); filaments very short, light pink, anthers white; style pink, stigma cherry red. Calyx 6mm long, light green. Lvs elliptic, 170 × 55mm, rounded at base, pointed at apex, margins flat, glossy green above, with white hairs when young, aging to grey. Shrub 3.1 × 2.3m in 20 years. Late May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2011, p.30. Etymology: named for a students’ association in the University of Latvia

(r) TANAGA

Trade designation for ‘Hachmann’s Tanaga’

(r) TANAGER

Trade designation for ‘Glendoick Tanager’

(r) ‘The Minahan’

cv. Elepidote rhododendron: ‘Janet Blair’ × ‘Crest’ (s) × ‘Yellow Balloon’
H: D. Mac Mullan (1989), G: D. Mac Mullan, N: D. Mac Mullan (2010), REG: D. Mac Mullan (2011)
Fls 12/conical truss, funnel-shaped, c.70 × 95mm, with 6 wavy-edged lobes, medium yellow in bud, opening strong medium yellow in the tube, paling towards the lobes; outside medium/light pure yellow; medium yellow speckles on the dorsal lobe. Lvs elliptic, 90 × 25mm, broadly acute at apex, rounded at base, semi-glossy and mid-green above, hairless. Shrub 1.2 × 1.2m in 7 years. Midseason.
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named after Joe Minahan, a friend of the registrant and another enthusiastic hybridiser of rhododendrons

(a) ‘Thomas David’

cv. Deciduous azalea: hybrid derived from ‘Homebush’
H: W.G.T. Hyde (1970s), G: W.G.T. Hyde (1970s), N: Mrs R. Legrand (2011), I: Millais Nurseries (2011), REG: Mrs R. Legrand (2011)

Fls double, 10–12/tight ball truss, trumpet-shaped, 47 × 29mm, with 9–10 wavy-edged lobes, deep red (185A) in bud, opening inside vivid reddish orange (42A), outside moderate red (47A); with a faint blotch on dorsal lobes of strong orange (25B); sweetly scented. Calyx irregular, 4–6mm long, strong yellow-green (144B). Lvs elliptic, 95 × 45mm, cuneate at base, acute at apex, margins decurved, matt green above, hairless. Shrub 2.1 × 1.6m in at least 30 years. May. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for a grandson of the registrant

(a) **'Tom Hyde'**

cv. Evergreen azalea: parentage unknown

H: W.G.T. Hyde (1970s), G: W.G.T. Hyde (1970s), N: Mrs R. Legrand (2011), I: Mrs R. Legrand, REG: Mrs R. Legrand (2011)

Fls open-trumpet-shaped, 45 × 25mm, with 7 wavy-edged lobes (the few petaloids becoming more noticeable as the flower matures), strong red (46A) in bud, opening vivid red (45A) both inside and out, with faint speckling of strong red (46A) on the three dorsal lobes. Calyx 5mm long, strong yellow-green (144A). Lvs narrowly elliptic, 30 × 13mm, cuneate at base, acute at apex, margins upcurved, glossy above, dark red (187A) and especially striking through the winter, more green in summer but still with a burnished appearance, hairless. Shrub 0.9 × 0.9m in 30 years or more. May.

Etymology: named for the grandfather of the registrant

(r) **'Toms'**

cv. Elepidote rhododendron: open-pollinated seedling from 'Eduards Smilģis'

H: R. Kondratovičs (1985), G: R. Kondratovičs (1991), N: R. Kondratovičs (2009), REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 17/strongly domed truss, open funnel-shaped, 70 × 60mm, with 5 wavy-edged lobes, buds strong purplish red (72A), opening inside light purple (84C), outside strong reddish purple (N78B), with a conspicuous blotch of dark greyish purple (N186B) on the dorsal lobes; filaments violet, anthers light brown; style purple, stigma cherry red. Calyx 6mm long, light green. Lvs elliptic, 130 × 55mm, pointed at base and apex, margins flat, matt green above, hairless. Shrub 1.8 × 1.8m in 20 years. Late May to early June.

Nomenclatural standard: colour print provided by the registrant. First published in *Dārzs un Drava*, March/April 2010, p.36. Etymology: Toms is a Latvian male christian name

(r) **'Triple A'**

cv. Lepidote rhododendron: *minus* Carolinianum Group, pink form (s) × *impeditum*, Cumerford selection

H: J.B. Brueckner (1970), G: J.B. Brueckner, N: A. Brueckner, A. Keefe & A. Woodward, (2011), REG: C. Woodward (2011)

Fls 3–5/domed truss, campanulate, 50–65 × 30–40mm, with 5 very wavy-edged lobes, strong purple (83C) or vivid violet (87A) or vivid purple (87B) in bud, opening inside vivid violet (87A) and

light purple (87C), unmarked; outside vivid violet (87A) and light purple (87C). Calyx 1–2mm long, strong yellow-green (145A). Lvs elliptic, 15 × 10mm, broadly acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, scaly below. Shrub with a dense habit, 0.6 × 0.6m in 40 years. Mid-May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the epithet acknowledges the connection between the raiser and his three grandchildren, whose names all begin with the letter A

(r) **'VERONIKA'**

Trade designation for 'Hachmann's Veronika'

(r) **'Viesturs'**

cv. Elepidote rhododendron: 'Carawbiense Grandiflorum' (s) × 'Purple Splendour'

H: R. Kondratovičs (1974), G: R. Kondratovičs (1980), N: R. Kondratovičs (1997), REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 19/slightly domed truss, open funnel-shaped, 70 × 60mm, with 5 wavy-edged lobes, buds dark purple (83A), opening inside strong purple (N81B), outside strong reddish purple (N78A), with discrete light olive (152B) spots on the dorsal lobes; filaments violet, anthers light violet; style red, stigma red. Calyx 2mm long, light green. Lvs elliptic, 155 × 55mm, rounded at base, pointed at apex, margins flat, matt green above, hairless. Shrub 1.4 × 1.8m in 10 years. Late May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, March/April 2010, p.35. Etymology: Viesturs is a Latvian male christian name

(a) **'Vija'**

cv. Deciduous azalea: 'Uguns' (s) × *luteum*

H: R. Kondratovičs (1978), G: R. Kondratovičs (1986), N: R. Kondratovičs (1998), I: Rhododendron Nursery Babīte, University of Latvia (1998), REG: Rhododendron Nursery Babīte, University of Latvia (2011)

Fls 15/slightly domed truss, tubular funnel-shaped, 70 × 70mm with 5 wavy-edged lobes, buds strong orange (168A), opening inside light orange-yellow (16B), outside light orange-yellow (16C), with a vivid orange-yellow (23A) blotch on the dorsal lobes; filaments yellow, anthers white; style yellow, stigma light green; slightly scented. Calyx 1mm long, green. Lvs elliptic, 140 × 45mm, pointed at base and apex, margins wavy, glossy green above, hairless. Shrub 1.5 × 1.5m in 10 years. Late May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in *Dārzs un Drava*, December 2007, p.17. Etymology: Vija is a Latvian female christian name

(r) **'Violet Breeze'**

cv. Elepidote rhododendron: 'Fancy' × *yakushimanum* (s) × 'Frank Galsworthy' × 'Purple Splendour'

H: J. Barlup (1995), G: J. Barlup (2000), N: J. Barlup (2010), REG: J. Barlup (2011)

Fls 13/ball truss, broadly funnel-shaped, 40 × 65mm, with 5 wavy-edged lobes, strong purplish red (72A) in bud, opening inside very pale purple (76C), changing to strong purple (77B) on the tips of the lobes, and with prominent dark red (183A), densely spotted flares on the two dorsal lobes, extending c.25mm from the base; outside very pale purple (76C), changing to strong purple (77B) on midribs and on the tips of the lobes. Lvs elliptic, 115 × 40mm, broadly acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, hairless. Shrub with a dense habit, 0.8 × 1.2m in 11 years. Late May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY)

(r) **‘Violet Touch’**

cv. *Elepidote rhododendron*: ‘Violet Mist’ (s) × ‘Plum Passion’

H: J. Barlup (2001), G: J. Barlup (2006), N: J. Barlup (2010), REG: J. Barlup (2011)

Fls 21/ball truss, broadly funnel-shaped, 50 × 65mm, with 5 very wavy-edged lobes, deep purplish red (71A) in bud, opening inside light purple (77D), with a dark red (167A) flare starting c.15mm from the base and extending upwards c.20mm on the dorsal lobe, and with the tips of all the lobes strong purple (77B); outside light purple (77D), with midribs and lobe tips strong purple (77B). Lvs elliptic, 115 × 50mm, broadly acute at apex, rounded at base, margins upcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 9 years. Early May.

(v) **‘Vuurdoop’**

cv. *Lepidote rhododendron*: ‘Apassionata’ (vireya) (s) × ‘Leonore Frances’

H: H. Van Oost (2002), G: H. Van Oost (2004), N: H. Van Oost (2010), I: H. Van Oost (2011), REG: H. Van Oost (2011)

Fls 4–7/domed truss, tubular funnel-shaped, 85 × 56–60mm, with 5 flat-edged lobes, lobes broadly margined orange inside and out, centre of lobes and the whole of the tube yellow both inside and out, unmarked; stigma green; anthers dark. Calyx vestigial. Lvs elliptic, 90 × 40mm, cuneate at base, obtuse and apiculate at apex, margins flat, matt green and with silvery scales above, with greenish brown scales below. Shrub 0.8 × 0.8m in 10 years. Flowering mostly in spring and autumn, but occasionally in winter and summer.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: the epithet means baptism of fire and refers to the colour. Resembles ‘Anatta Gold’ in colour but with smaller corollas and with a plant habit that is more compact and upright growing

(r) **‘Weissenburg’**

cv. *Elepidote rhododendron*: ‘Whitestone’ (s) × *yakushimanum*, dwarf form

H: J. Nicolella (1992), G: K. Heger (2000), N: K. Heger (2003), I: Holger Hachmann (2009), REG: M. Oppenrieder (2011)

Fls up to 19/rounded, dense to slightly loose truss,

fully double, funnel-shaped, 60–70mm wide, with up to 12 lobes (all the stamens petaloid), outer lobes wavy, some of the inner lobes very wavy, pale pink in bud (like little pink rosebuds), opening inside delicate pinkish white, becoming all white with a slightly greenish base and greenish spots on the dorsal lobe; outside pinkish white; style white, stigma yellowish green. Lvs elliptic, 117–120 × 40mm, glossy green above, with a thin, hairy indumentum below. Shrub 0.7 × 1.0m in 10 years. May.

Nomenclatural standard: colour print provided by the registrant (WSY). First published in Hachmann cat. 2009–2013 (2009)

(r) **‘Wild Berry’**

cv. *Elepidote rhododendron*: ‘Violet Mist’ (s) × ‘Plum Passion’

H: J. Barlup (2001), G: J. Barlup (2005), N: J. Barlup (2010), REG: J. Barlup (2011)

Fls 14/ball truss, broadly funnel-shaped, 40 × 65mm, with 5 very wavy-edged lobes, deep purplish red (71A) in bud, opening inside light purple (77D), changing to strong purple (77B), with a moderate red (183C) dorsal flare extending upwards c.30mm from the base; outside light purple (77D), changing to strong purple (77B), with the tips of the lobes deep reddish purple (77A) and with strong purplish red (71B) midveins. Lvs elliptic, 115 × 45mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.6m in 9 years. Mid-May.

(a) **‘Zoe Elizabeth Stoltz’**

cv. Evergreen azalea: ‘Carol Kittel’ (s) × ‘Girard’s Fuchsia’

H: J.F. Klimavicz (1998), G: J.F. Klimavicz (2000), N: K. Stoltz (2011), REG: J.F. Klimavicz (2011)

Fls semi-double, 2–3/ball truss, saucer-shaped when fully open, 40 × 70mm, with 13 very wavy-edged lobes, buds vivid reddish purple (78A), with white along the midveins, opening inside strong reddish purple (78B) at the edges of each lobe, fading to white in the centre of the corolla, with a slight spotted blotch of moderate purplish red (58A); outside strong reddish purple (78B), with white along the midveins. Calyx 10mm long, light yellow-green (145D). Lvs elliptic, 45 × 20mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above, hairless. Dense shrub, 0.5 × 0.5m in 5 years. Early May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY). Etymology: named for the deceased daughter of Ken Stoltz, who won the right to name this plant in a charity auction sponsored by the National Institutes of Health for the Children’s Inn

Names and addresses

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

* = deceased

Ahern, T.
Bethlehem, Pennsylvania, USA

Aris Horticulture, Inc.
Barberton, Ohio, USA

Barlup, J.
Bellevue, Washington, USA

Baumschule Wüstemeyer
Schermbach, Germany

Bergmann, W.R.
Lehigh Acres, Florida, USA

*Brueckner, J.B.
Mississauga, Ontario, Canada

Chappell, P.G.G.
Boldre, Lymington, Hampshire, England

Collins, E.W.
Hendersonville, North Carolina, USA

Colombel, M.
Menez-Rohou, Fouesnant, France

*Cowles, J.C.
Stow, Massachusetts, USA

Cox, K.N.E. & P.A.
Glendoick Gardens, Glencarse, Perthshire, Scotland

Darney, C.O.
Parkesburg, Pennsylvania, USA

Doppel, J.
Lenhartsville, Pennsylvania, USA

Dosser, D.J.
Lower Wilmot, Tasmania, Australia

Furno, N.
Parco Burcina, Biella, Italy

Garbutt, T.A.
Oamaru, North Otago, New Zealand

George, R.E.
Sammamish, Washington, USA

Gimle Planteskule
Nordheimsund, Norway

Glendoick Gardens
Glencarse, Perthshire, Scotland

Greenleaf Nursery Company
Park Hill, Oklahoma, USA

Hachmann, H. (Hans Hachmann)
Baumschule H. Hachmann, Barmstedt, Germany

Hachmann, Holger
Baumschule H. Hachmann, Barmstedt, Germany

Head, R.H. & L.J.
Seneca, South Carolina, USA

Heger, K.
Krefeld, Germany

Heritage Museum and Gardens
Sandwich, Massachusetts, USA

Hesse, H.
Hamersen, Germany

High Beeches Gardens Conservation Trust, The
Handcross, West Sussex, England

Hübbers, K.-H. & Mrs L.
Kranenburg, Germany

*Hyde, W.G.T.
Ferndown, Dorset, England

Jørgensen, P.M.
Det Norske Arboret, Hjelstad, Norway

Kanke, S.
Yaizu-district, Fukushima-prefecture, Japan

Keefe, A.
Mississauga, Ontario, Canada

Klimavicz, J.F.
Vienna, Virginia, USA

Kondratovičs, R.K.
Riga, Latvia

Le Duigou, A. & L.
Raden, Scaer, France

Leggett, J.
Waianakarua, North Otago, New Zealand

Legrand, Mrs R.
West Moors, Dorset, England

MacDonald, B.
Hanson, Massachusetts, USA

Mac Mullan, D. Hamburg, Pennsylvania, USA	University of Canterbury Ilam Gardens, Christchurch, New Zealand
McDonald, S.F. Le-Mac Nurseries, Hampton, Virginia, USA	Van Oost, H. Kruishouten, Belgium
Millais Nurseries, Farnham, Surrey, England	Westhoff, J. Bremen, Germany
Mitchell, R.A. Volcano, Hawaii, USA	Wolin, M.L. East Brunswick, New Jersey, USA
Nicolella, J. Hampton Bays, New York, USA	Woodward, C. & A. Mississauga, Ontario, Canada
Norske Arboret, Det Hjellestad, Norway	Wright, H.R. Courtenay, British Columbia, Canada
Oppenrieder, M. Baumschule H. Hachmann, Barmstedt, Germany	Wüstemeyer, W. Baumschule Wüstemeyer, Schermbeck, Germany
Pacific Island Nursery Kea'au, Hawaii, USA	
Player, L.M. Fayetteville, North Carolina, USA	
*Pratt, M.C. Pulborough, West Sussex, England	
Rhododendron Nursery Babīte, University of Latvia Riga, Latvia	
Rouse, A. Hawthorn East, Victoria, Australia	
Sandwich Club Sandwich, Massachusetts, USA	
Shellenberger, R. Snohomish, Washington, USA	
Sim, Mrs A. Oamaru, New Zealand	
Smith, P. Santa Monica, California, USA	
*Stead, E.F. Ilam, Christchurch, New Zealand	
Stoltz, K. USA	
Suzuki, H. Koriyama-city, Fukushima-prefecture, Japan	

Corrections to the International Rhododendron Register and Checklist

The following list indicates further significant additions and corrections that need to be made to the printed text of the *Register and Checklist* published in 2004 or to the six subsequent Supplements.

The Registrar is grateful to all those who have brought such errors to his notice and welcomes knowledge of further amendments that may need to be made.

Minor changes are not noted here but such changes have been made to the database and will be available in any future electronic version. The database will also be taking into account additional information that comes to light relating to any individual entry, as well as including further entries to the Checklist.

In the following list the page number refers to the page in the *Register and Checklist* (or one of its Supplements) on which the entry concerned is to be found.

‘Chariots of Fire’ (Fourth Supplement, p.4)

This should be recorded as a deciduous, not an evergreen azalea

‘Rhodunter 48’ (p.1159)

Add trade designations: INKARHO-DUFHECKE, WEISS and INKARHO FRAGRANT HEDGE, WHITE

‘Rhodunter 149’ (Third Supplement, p.21)

Add trade designations: INKARHO-DUFHECKE, LILA and INKARHO FRAGRANT HEDGE, LILAC

‘Pukeiti Skyrocket’ (p.1116)

The origin of this plant had been confused by the registrant: it had in fact originally been collected by Norman Cruttwell (pre-1986) on Mt Suckling, Papua New Guinea and taken to his Lipizauga Botanical Sanctuary on Mt Gahavisukar, from where it had been obtained by Graham Smith. It has sometimes been known as ‘Starburst’ and may be a selection from *archboldianum*

‘Starburst’

Add as a synonym of ‘Pukeiti Skyrocket’ (above)

‘Wesley Hayes’ (p.1441)

The flower colour should be amended from white to read: scarlet red, aging to light red. It is probably an *arboreum* hybrid.

ISBN 9781907057311

