



The International Rhododendron Register and Checklist (2004)

Fifth Supplement



© 2009 The Royal Horticultural Society
80 Vincent Square, London SW1P 2PE, United Kingdom
www.rhs.org.uk

International Rhododendron Registrar:
Dr A C Leslie

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781907057007

Printed and bound in the UK by Page Bros, Norwich

Cover: *Rhododendron* 'Mrs Charles E. Pearson' by Alfred John Wise

The International Rhododendron Register and Checklist (2004)

Fifth Supplement

Introduction	page 2
Names registered, January–December 2008	page 3
Names and addresses	page 34
Corrections	page 37

Introduction

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first four Supplements included names registered from 1 January 2003 to 31 December 2007. This Fifth Supplement includes all those registered in 2008. A further list of corrections is also included here.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7–10 of the Introduction.

It may be helpful here to repeat that the following abbreviations are employed:

(a) = azalea
(az) = azaleodendron
cv. = cultivar
G = grown to first flower by...
gp = Group
H = hybridised by...
I = introduced by...
N = named by...
(r) = rhododendron
R = raised by...
REG = registered by...
S = selected by...
(s) = seed parent
(v) = vireya

Where possible colour references are given to the *RHS Colour Chart* (1966, 1986, 1995 & 2001) with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS website (www.rhs.org.uk/plants/registerpages.Rhodoform.PDF) or from one of the national or regional registrars as follows:

North America	Mrs J.W. Murray, 21 Squire Terrace, Colts Neck, New Jersey 07722-1021, USA
Australia	Mr K. Gillanders, Woodbank, 2040 Huon Road, Longley, Tasmania 7150, Australia
New Zealand	Mr B.J. Coker, 42 Massey Crescent, St Albans, Christchurch 8014, New Zealand
Japan	Dr K. Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken, Japan 768-0021

Dr A.C. Leslie
International Rhododendron Registrar
109 York Street
Cambridge CB1 2PZ
England

(email: alanleslie@rhs.org.uk)

Names registered January–December 2008

(r) 'Abby Kate'

cv. *Elepidote rhododendron* : 'Cup Day' (s) × 'Samantha Sawers'

H: D.J. Dosser (1995), G: D.J. Dosser (2003), N: D.J. Dosser (2007), I: B.S. Coates (2007), REG: D.J. Dosser (2008)

Fls 13/ball truss, funnel-shaped, 65 × 85mm, with 5 wavy-edged lobes, buds vivid purplish red (66A), opening vivid purplish red (66A) inside and out, with moderate purplish red (58A) spots on the dorsal lobe. Calyx 25mm long, vivid purplish red (66A). Lvs lanceolate, 150 × 35mm, margins wavy and decurved, glossy above, obtuse at base, cuspidate at apex, hairless. Shrub 1.5 × 1.5m in 10 years. November (Australia).

(Etymology: named for a granddaughter of the introducer. Nomenclatural standard: colour print provided by the registrant (WSY0101030))

(v) 'Abby Lauren'

cv. *Lepidote rhododendron* : 'Teddy's Best' (s) × 'Pastenello'

H: G.L.S. Snell (2000–2001), G: K. Hagan (2006), N: K. Hagan (2008), REG: K. Hagan (2008)

Fls 10/lax truss, tubular funnel-shaped, 75 × 60mm, with 5 flat-edged lobes, inside light yellow (15D), with light orange (29B) ventral lobes, outside light yellow (15D); unmarked. Calyx 1mm long, light green. Lvs oblanceolate, 110 × 50mm, acute at apex, cuneate at base, margins flat, glossy green above. Shrub 0.9 × 0.8m in 7 years.

(Nomenclatural standard: colour print provided by the registrant (WSY0101961). Etymology: named for a granddaughter of the registrant)

(r) 'Adrien Le Fur'

cv. *Elepidote rhododendron* : 'Haloceanum' (s) × 'Coronation Day'

H: J.F. Saint-Jalm (1996), G: J.F. Saint-Jalm (2001), N: J.F. Saint-Jalm (2008), REG: J.F. Saint-Jalm (2008)

Fls 11–12/loose ball truss, broadly funnel-shaped, 65 × 125mm, with 6 wavy-edged lobes, buds strong purplish red (63B), opening inside very pale purple (69D), fading to white, outside strong purplish red (63B); a deep purplish red (59B) blotch surrounds the base of the throat and there are numerous small, but relatively inconspicuous spots chiefly on the dorsal lobe/dorsal throat; stamens 14, filaments white; style white, stigma light yellow-green (145C). Calyx 1mm long, strong purplish red (63B). Lvs elliptic, 100 × 70mm, rounded at base, broadly acute at apex, hairless. Shrub 2.2 × 1.7m in 11 years. Mid April to early May.

(Nomenclatural standard: colour print provided by the registrant (WSY0101331). Etymology: named for a great enthusiast for rhododendrons and other plants)

(r) 'Alberto Maria De Agostini'

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)

Fls 12–14/hemispherical truss, 35–40 × 60–65mm, with 5 slightly wavy-edged lobes, buds strong purple (77B), opening inside and out light purple (76A), with a lighter centre (very light purple, 76B), with strong orange-yellow (163B) and strong yellow (153B) markings on the dorsal lobe which just spread on to the adjacent lobes; stamens 10, anthers greyish purplish red (N77B) and strong reddish purple (N78A), filaments moderate purplish pink (N74D); pistil not or only just exerted from the corolla, up to 4mm longer than the stamens, style moderate purplish pink (62B) with a paler base (62C), stigma moderate red (185B). Lvs lanceolate, 106–114 × 39–42mm, flat, matt green above, cuneate at base, obtuse at apex, moderate olive green (147A) above, moderate yellow-green (146D) below, hairless; petioles 15–20mm long. Shrub 2.5m. May.

(Nomenclatural standard: colour print from an electronic image provided by the registrants (WSY0101262). Etymology: named for a member of the Salesian Order (during the 19/20th century) who travelled in Patagonia: his brother Giovanni De Agostini founded the De Agostini Geographic Institute)

(r) 'Alfonso Sella'

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)

Fls 17–21/domed truss, 30–35 × 57–65mm, with 5 smooth-edged lobes, strong reddish purple (70B) in bud, opening deep to moderate purplish pink (N74C-D), with spots of moderate brown (165A) to moderate reddish brown (166B) in the throat and on the three dorsal lobes; stamens 10, held inside the corolla, anthers light purple (76A), filaments light purple (N78D); pistil exerted from the corolla, exceeding the stamens by 10–12mm, style strong purplish red (63B), stigma deep red (185A). Lvs lanceolate, 132–166 × 45–51mm, margins flat, cuneate at base, acute at apex, matt moderate olive green (147A) above, moderate yellow-green (146D) below, hairless; petioles 13–18mm long. Shrub 3.0m. May. (Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101264). Etymology: named for an Italian botanist from Biella, Italy)

(v) 'Alicia Kaye'

cv. *Lepidote rhododendron* : 'Gardenia Odyssey' (s) × 'Australia II'

H: G.L.S. Snell (2002), G: K. Hagan (2007), N: K. Hagan (2008), REG: K. Hagan (2008)

Fls 9/lax truss, tubular funnel-shaped, 70 × 105mm, with 5 wavy-edged lobes, light yellow-green (142C) in bud, opening inside white (155A), flushed moderate yellowish pink (37C) in the throat, outside white (155A), flushed moderate yellowish pink (37C), with

the base of the tube light yellow (20B); unmarked; scented. Lvs elliptic, 100 × 55mm, broadly acute at apex, cuneate at base, margins upcurved, glossy green above. Shrub 25 × 30cm in 5 years. Mid spring to autumn.

(Nomenclatural standard: colour print provided by the registrant (WSY0101959). Etymology: named for a granddaughter of the registrant)

(r) **'Angel Eyes'**

cv. *Elepidote rhododendron* : 'Anna' (s) ×

yakushimanum 'Koichiro Wada'

H: R.A. Furman (1979), G: R.A. Furman (1984),

N: R.A. Furman (2008), REG: R.A. Furman (2008)

Fls 12/ball truss, open funnel-shaped, c.70–75 × 115–125 mm, with 6 very wavy-edged lobes, strong purplish pink (55B) in bud, opening white (155D) throughout, with a basal blotch of strong red (46A) and strong red (53C), waxy and of heavy substance. Lvs elliptic, 75–175 × 25–50mm, obtuse at apex, rounded at base, margins downcurved, semi-glossy and dark green above, hairy below, the hairs white when young aging to tan. Shrub 3.0 × 2.7m in 29 years. Mid May.

(r) **'Anna Holm'**

cv. *Elepidote rhododendron* : parentage unknown

H: D.A. Griswold (1970), G: Griswold Nursery,

N: Griswold Nursery, I: Griswold Nursery,

REG: Griswold Nursery (2008)

Fls 16/ball truss, funnel-campanulate, 65 × 75mm, with 7 wavy-edged lobes, light greenish yellow (1C) in bud, opening pale greenish yellow (1D) throughout, with short, red, vertical markings in the throat of the dorsal lobes; slightly scented. Lvs elliptic, 140 × 50mm, broadly acute at apex, cordate at base, with wavy margins, moderate olive green (147A) above, hairless; petioles 25mm long, reddish orange, visibly hairy. Upright shrub, 1.8 × 1.2m in 37 years. May. (Etymology: named for the mother-in-law of Linda Malland's sister. Linda Malland compiles the registration data for Griswold Nursery)

(a) **'Ann Lee McPhail'**

cv. Evergreen azalea : 'Chikyu-no-haru' (s) ×

'Surprise' (Glenn Dale)

H: J.O. Harris (c.1978), G: J.O. Harris (c.1980),

N: J.O. Harris (c.1981), I: J.O. Harris (c.1981),

REG: J.O. Harris (2008)

Fls single, 5/domed truss, funnel-campanulate, c.75 × 70mm, with 6 wavy-edged lobes, opening inside light purplish pink (55C), edged deep purplish pink (55A), outside pale purplish pink (56D); unmarked. Calyx c.13mm long, brilliant yellow-green (149B). Lvs ovate, 45 × 22mm, obtuse at apex, rounded at base, margins upcurved, dull and deep yellowish green (141B) above, hairless. Shrub 1.1 × 0.8m in 10 years. Early April.

(Etymology: named for a member of the former Ralph Pennington chapter of the American Azalea Society. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.64)

(r) **'April Morn'**

cv. *Elepidote rhododendron* : 'Nancy Evans' (s) × 'Tia'

H: J. Barlup (2000), G: J. Barlup (2005), N: J. Barlup

(2008), REG: J. Barlup (2008)

Fls 14/domed truss, funnel-campanulate, 50 × 70mm, with 5 wavy-edged lobes, strong red (53B) in bud, opening inside and out with broad light pink (49C) margins, blending to pale greenish yellow (10D) towards the centre; with a strong red (53C) basal blotch extending upwards to c.15mm on the dorsal lobe and c.5mm on the other lobes. Calyx vestigial. Lvs elliptic, 95 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and dark yellowish green (139A) above, hairless. Shrub with a dense habit, 0.5 × 0.8m in 8 years. Late April.

(r) **'Armonica'**

cv. *Elepidote rhododendron* : parentage unknown

G: Parco Burcina, S: E. Accati & N. Furno (2000),

N: E. Accati & N. Furno (2008), REG: E. Accati &

N. Furno (2008)

Fls 12–15/rather loose, hemispherical truss, 40–50 × 65–70mm, with 5 rounded, smooth-edged lobes, deep purplish pink (68A) in bud, opening light to deep purplish pink (75B, N66D) inside and out, with a white centre; dark red to deep purplish red (187A, B, C, D) spots mainly on the dorsal lobe and just extending onto the adjacent lobes; stamens 10, held within the corolla, anthers moderate purplish pink (186D) and pale yellow (162D), filaments white; pistil 5–10mm longer than the stamens but still held within the corolla, style pale greenish yellow (2D), stigma strong reddish orange (169A). Lvs lanceolate, 120–140 × 50–60mm, acute at base and apex, margins flat, upper surface wrinkled, matt and moderate olive green (147A) above, moderate yellow-green (146C) below; petioles 20–25mm long. Shrub 3.0m. May. (Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101266). Etymology: named after one of the farmsteads in the Parco Burcina)

(a) **'Balbo'**

cv. Evergreen azalea : parentage unknown

S: A. Balbo (2002), G: A. Balbo (2004), N: A. Balbo

(2008), I: A. Balbo (2008), REG: A. Balbo (2008)

Fls single, 1–3/hemispherical truss, funnel-shaped, 45–50 × 25–28mm, with 5 smooth to very slightly wavy-edged lobes, strong purplish pink (62A) in bud and throughout on opening, with strong purplish red (71B) spots on the three dorsal lobes; stamens 5, exerted from the corolla, filaments white at base, shaded strong purplish pink (73B) above, anthers deep orange-yellow (163A); style exceeding the stamens, light yellow-green (142D) at base, shading to white and then to light purplish pink (62C) at the apex, stigma strong purplish pink (62A). Calyx 12mm long, moderate yellow-green (146C). Lvs lanceolate, 20–30 × 10mm, acute at base and apex, margins flat, glossy and moderate yellow-green (137C) above and moderate yellow-green (146C) below when mature, with some hairs. Shrub 0.8 × 0.6m in 5 years. May. (Etymology: the family name of the registrant)

(v) **'Barbara Campbell'**

cv. *Lepidote rhododendron* : *konori* hybrid (s) × (*macgregoriae* × *aurigeranum*)
H: R.A. Mitchell (2002), G: R.A. Mitchell (2007),
N: R.A. Mitchell (2008), REG: R.A. Mitchell (2008)
Fls 14/domed truss, tubular funnel-shaped, 80 × 80mm, with 8 wavy-edged lobes, opening inside pale purplish pink (65C), outside pale purplish pink (65D); unmarked; slightly scented. Lvs elliptic, 105 × 70mm, acuminate at apex, rounded at base, margins upcurved, dull and strong yellow-green (144B) above, scaly (more densely so below). Shrub 0.9 × 0.75m in 6 years. Early July.
(Etymology: named for the instigator in founding the ARS chapter in Hawaii)

(r) **'Beni-yaku-yama'**

cv. *Elepidote rhododendron* : 'Yaku-yama-zakura' (s) × 'Yaku-yama-zakura'
H: Y. Yamamoto (2001), G: Y. Masuda (2007),
S: Y. Yamamoto (2008), N: Y. Masuda (2008),
REG: Y. Masuda (2008)
Fls 9–12/loose open-topped truss, broadly funnel-campanulate, 40 × 60mm, with (6–)7 slightly wavy-edged lobes, buds red, opening inside pink, outside deep pink, with orange spotting in the dorsal throat. Calyx 10mm long, pale brown. Lvs narrowly elliptic to narrowly oblanceolate, 120 × 25mm, narrowly obtuse and mucronate at apex, margins flat, glossy green above, with white scales below when young, later with no indumentum. Shrub 0.2 × 0.3m in 7 years (pot grown). Late April.
(Nomenclatural standard: colour slide provided by the registrant (WSY0101459). Etymology: "beni" refers to the deep pink colour and "yaku-yama" to the parental cultivar)

(r) **'Bill Fee'**

cv. *Elepidote rhododendron* : ((*arboreum* × Elizabeth Group) × *arboreum*) (s) × 'Red Majesty'
H: E.W. Moyles (1988), G: E.W. Moyles (1993),
N: D.S. Wallace (2008), I: Singing Tree Gardens (2008), REG: Singing Tree Gardens (2008)
Fls 18/domed truss, funnel-campanulate, 40 × 50mm, with 5 very wavy-edged lobes, vivid red (46B) in bud, opening vivid red (44A) throughout, unmarked. Lvs lanceolate, 100 × 25mm, acute at apex, cuneate at base, margins downcurved, dull and strong yellowish green (141C) above, hairy below on the leaf blade, midrib and on the petiole (the hairs moderate yellow (160A) when young and when mature). Shrub 1.8 × 1.8m in 10 years. Early February (continuing with some flowers for several months).
(Etymology: named for a former logger, turned nurseryman, of Fort Bragg, California, USA, who gave this previously un-named seedling to the nominant)

(r) **'Brea Renee'**

cv. *Elepidote rhododendron* : 'Ross Maud' (s) × 'Samantha Sawers'
H: D.J. Dosser (1990), G: D.J. Dosser (1996),
N: D.J. Dosser (2007), REG: D.J. Dosser (2008)
Fls 15/ball truss, funnel-shaped, 50 × 100mm, with 5 very wavy-edged lobes, strong purplish red (63B) in

bud, opening light purplish pink (65B) inside and out, with spots of strong purplish pink (68B) on two dorsal lobes; slightly scented. Lvs lanceolate, 130 × 65mm, matt green above, truncate at base, acute at apex, hairless. Shrub 2.0 × 2.0m in 17 years. October to November.

(Nomenclatural standard: colour print provided by the registrant (WSY0100800). Etymology: named for the deceased daughter of a customer of the registrant)

(r) **'Brittany's Oso Fine'**

cv. *Elepidote rhododendron* : ((*caucasicum* × (Fabia Group × (Dido Group × 'Lem's Goal')) × 'Crest') (s) × 'Pridenjoy'
H: L.L. Newcomb (1986), G: B. Reading (1994),
N: C. Blankenship (2007), I: Chuck's Nursery & Hammond Nursery (2009), REG: C. Blankenship (2008)
Fls 9–11/domed truss, open funnel-shaped, 65 × 90mm, with 6–8 wavy-edged, sometimes imbricated lobes, buds pale yellowish pink (29D) and moderate yellowish pink (31D) with a light yellow (11B) base, opening inside with a brilliant greenish yellow throat and brilliant yellow (11A) centre, edged with a 25mm wide band of strong pink (49A) and a 6mm outer margin of light yellowish pink (36C); with a speckled dorsal flare of strong red (46A); outside with base brilliant yellow-green (154C), extending up 25mm, blending to strong pink (49A), with a narrow edge of light yellowish pink (36C). Calyx 6–20mm long, brilliant yellow (11A). Lvs elliptic, 65–125 × 25–75mm, obtuse at apex, cuneate at base, margins flat, semi-glossy and moderate olive x (147A) above, hairless. Dense shrub, 0.9 × 1.2m in 7 years. Early May.
(Etymology: Brittany is the christian name of a friend of the registrant)

(r) **'Burgundy Mist'**

cv. *Elepidote rhododendron* : 'Midnight Mystique' (s) × 'Jonathan Shaw'
H: J. Barlup (1999), G: J. Barlup (2004), N: J. Barlup (2006), REG: J. Barlup (2008)
Fls 17/ball truss, broadly funnel-shaped, 45 × 75mm, with 5 very wavy-edged lobes, dark reddish purple (77A) in bud, opening inside light purple (77D), with a margin of deep reddish purple (77A), c.15mm wide, and with a deep purplish red (61A) flare, extending c.6mm from the base and changing to dark red (183B) spots 25mm from the base; outside deep reddish purple (77A), with deep purplish red (61A) midribs and a ring c.20mm wide at base. Calyx vestigial. Lvs elliptic, 125 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 9 years. Late May.

(a) **'Caerhays Lavender'**

cv. Evergreen azalea : parentage unknown
R: J.C. Williams (1930s), I: Burncoose Nurseries (1983), REG: C.H. Williams (2008)
Fls single, 2–3/truss, light purple (N78D) in bud, opening lavender. Calyx long hairy. Later lvs

oblanceolate, c.27 × 11mm, tips submucronate, long hairy, lighter than moderate olive green (147A) above, lighter than moderate yellow-green (147B) below; spring leaves oblanceolate, c.40 × 15mm, light greenish yellow (6D), long hairy. Remarkable for the startling white new growth, which slowly turns to green after 4–6 weeks (but will last longer in full or partial shade). Shrub 1.0 × 1.5m in 20 years. April to May. (Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.191)
AM 2008

(a) **‘Caparo’**

cv. Deciduous azalea : ‘Washington State Centennial’ (s) × (‘Raimunde’ × ‘Twilight Sky’)
H: K.-H. Hübbers (1994), G: K.-H. Hübbers (1999), N: K.-H. Hübbers (2006), REG: J. Westhoff (2008)
Fls up to 10/slightly flattened, globe truss, open funnel-shaped, 50 × 100mm, with 5 very wavy-edged lobes, buds strong reddish orange (42B), opening inside a mixture of vivid yellow (14B), vivid reddish orange (33B) and deep yellowish pink (47C), outside a mixture of pale orange-yellow (24D), strong yellowish pink (37A) and deep yellowish pink/deep pink (47C/D); with a vivid orange-yellow (23A) blotch on the dorsal lobe; slightly scented. Calyx 10mm long, light green, translucent. Lvs elliptic, 90 × 30mm, margins flat, cuneate at base, acute and slightly mucronate at apex, matt green above. Shrub 0.8 × 0.7m in 10 years. Mid May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101567). Etymology: named for a city in Trinidad, the birthplace of the raiser’s wife)

(r) **‘Catherine’s Lemon Meringue’**

cv. Elepidote rhododendron : ‘Lemonade’ (s) × ‘Lem’s Cameo’
H: C. Weeks (1990), G: C. Weeks (1995), N: C. Weeks (2008), I: Westgate Gardens Nursery (2008), REG: C. Weeks (2008)
Fls 12–13/domed truss, broadly funnel-shaped, 55 × 115mm, with 6 wavy-edged lobes, light greenish yellow (1C) in bud, opening the same colour throughout and with red speckling on the three dorsal lobes. Calyx variable in length, from negligible to 20mm long, light greenish yellow (1C). Lvs obovate, 175 × 75mm, broadly acute at apex, oblique at base, margins flat, semi-glossy and deep yellowish green (141B) above, hairless. Shrub 0.8 × 1.1m in 10 years. Mid April to mid May.

(r) **‘Chantal de Kersulec’**

cv. Elepidote rhododendron : ‘Mrs Charles E. Pearson’ (s) × *grande*
H: R. Grall (1994), G: R. Grall (2001), N: Mme C. Bordaïs (2008), REG: Société Bretonne du Rhododendron (2008)
Fls 18–21/ball truss, funnel-campanulate, 50–55 × 60mm, with 5 slightly wavy-edged lobes, buds strong purplish red (60B), opening inside deep purplish pink (N74C), outside deep purplish pink (N74C) and with stripes; greyish purple (77A) spots on the dorsal

lobe, coalescing into a narrow solid mark towards the base of the throat; stamens 10–11, all upcurved; style white at first soon changing to light pink. Calyx 7–8mm long, very pale green (136D). Lvs elliptic, 155 × 55mm, margins decurved, cuneate at base, broadly acute at apex, matt dark green (136A) above, with a silvery indumentum when young, which gets thinner and becomes grey with age; petioles conspicuously yellow-green. Shrub 1.35 × 1.5m in 13 years. April to May.

(Nomenclatural standard: colour print provided by the registrant (WSY0101191). Etymology: named after Madame Chantal Bordaïs and the weekend cottage she and her husband use at Scaër: kersulec in Breton means «small house for Sundays»)

(r) **‘Cherry Surprise’**

cv. Elepidote rhododendron : ‘Exbury Calstocker’ (s) × ‘Rubicon’
H: C. Weeks (1998), G: C. Weeks, N: C. Weeks (2008), I: Westgate Gardens Nursery (2008), REG: C. Weeks (2008)
Fls 20/domed to ball truss, funnel-campanulate, 65 × 75mm, with 5 wavy-edged lobes, strong purplish red (54A) in bud, opening strong purplish red (54A) throughout and aging to strong red (53B), the three dorsal lobes with a moderately spotted, strong purplish red (54A) blotch; centre very dark red. Lvs elliptic, 180 × 50mm, acuminate at apex, cuneate at base, margins downcurved, semi-glossy and dark bluish green (133A) above, hairless. Shrub 0.9 × 1.5m in 9 years. Early to mid March.

(a) **‘Cille Shaw’**

cv. Evergreen azalea : ‘Amagasa’ (s) × ‘Grace Freeman No. 2’
H: J.O. Harris (c.1978), G: J.O. Harris (c.1980), N: J.O. Harris (c.1981), I: J.O. Harris (c.1981), REG: J.O. Harris (2008)
Fls single, solitary, saucer-shaped, c.40 × 65mm, with 5 very wavy-edged lobes, very pale purple (69A) in bud, opening inside and out very pale purple (69A) in the centre, shading to strong reddish purple (70B) at the edges; faint pale yellow-green (4D) blotch on the three dorsal lobes. Calyx c.5mm long, strong yellow-green (142A). Lvs obovate, 45 × 22mm, broadly acute at apex, rounded at base, margins flat, hairless. Shrub 0.5 × 0.7m in 10 years. Late April.
(Etymology: named for a member of the former Ralph Pennington chapter of the American Azalea Society. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.252)

(r) **‘Città di Biella’**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 26–29/hemispherical-conical truss, funnel-shaped, 48–54 × 48–60mm, with 5–7 wavy-edged lobes, buds strong reddish purple (72C), opening deep purplish pink (N74C) throughout, with strong purplish red (61B) spots on a paler ground on the

three dorsal lobes; stamens 10, included within the corolla, anthers greyish purplish red (N77D), filaments very light purple (75C); pistil 10–12mm longer than the stamens and exceeding the corolla, style deep purplish pink (N57D), stigma strong red (53B). Lvs lanceolate to ovate, 136–155 × 56–61mm, acute at base and apex, margins flat, upper surface matt moderate olive green (147A), moderate yellow-green (146C) below, hairless; petioles 18–22mm long. Shrub 3.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101346). Etymology: named for the capital city of the Italian province (Biella, Piedmont) in which the park is cited)

(r) **‘Cloudburst’**

cv. *Elepidote rhododendron* : ‘Wizard’ (s) × *irroratum* ‘Polka Dot’
H: D.S. Wallace (1995), G: D.S. Wallace (2000), N: D.S. Wallace (2008), I: Singing Tree Gardens (2008), REG: Singing Tree Gardens (2008)
Fls 17/domed truss, broadly funnel-shaped, 40 × 65mm, with 5 wavy-edged lobes, pale yellow (19C) at base in bud, shading to strong yellowish pink (43D) at apex, opening inside pale yellow (18D) in throat which extends into the middle of each lobe, with light yellowish pink (29C) lobe margins and with prominent spotting of strong red (53D) radiating outwards from the centre onto the lobes, the spots growing larger towards the edges; strong red (46A) nectar pouches; outside pale yellow (18D), shading to light yellowish pink (29C) along margins. Lvs lanceolate, 115 × 25mm, acute at apex, cuneate at base, margins downcurved, dull and vivid yellowish green (134A) above, hairless. Shrub 1.8 × 1.8m in 10 years. Early season.

(a) **‘Coronado Red’**

cv. *Evergreen azalea* : ‘Chikyu-no-haru’ (s) × ‘Surprise’ (Glenn Dale)
H: J.O. Harris (c.1979), G: J.O. Harris (c.1981), N: J.O. Harris (c.1982), I: J.O. Harris (c.1982), REG: J.O. Harris (2008)
Fls single, 2–3/domed truss, broadly funnel-shaped, c.60 × 75mm, with 5 flat-edged lobes, deep red (53A) in bud, opening deep red (53A) throughout. Calyx c.5mm long, strong yellow-green (142A). Lvs obovate, 45 × 22mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and strong yellow-green (143A) above, hairless. Shrub 1.2 × 1.2m in 10 years. Late April.
(Etymology: Coronado is the name of a housing subdivision near Lawrenceville (the home town of the registrant))

(r) **‘Cosmic Drops’**

cv. *Elepidote rhododendron* : ‘Nancy Evans’ (s) × ‘Lem’s Cameo’
H: C. Weeks (1995), G: C. Weeks (2003), N: C. Weeks (2008), I: Westgate Gardens Nursery (2008), REG: C. Weeks (2008)
Fls 18/domed truss, funnel-shaped, 65 × 90mm, with 6–8 wavy-edged lobes, buds brilliant yellow (10A) at

base, shading to strong yellowish pink (37A) at apex, opening inside brilliant yellow (10A) at base and strong yellowish pink (37A) at margins with the intermediate area very pale pink, close to white; outside colour similar to inside; prominent blotch of discrete, strong red (46A) spots arranged in a pattern of vertical lines on the three dorsal lobes. Lvs obovate, 150 × 50mm, broadly acute at apex, oblique at base, margins flat, semi-glossy and moderate olive green (137A) above, hairless. Shrub 0.9 × 0.8m in 6 years. Mid April to May.

(r) **‘Cowles Gold’**

cv. *Elepidote rhododendron* : parentage unknown
H: J.C. Cowles (pre 1996), G: B. Feller (2000), N: B. Feller (2008), REG: B. Feller (2008)
Fls 15/domed truss, open funnel-shaped, 65 × 115mm, with 7 wavy-edged lobes, strong red (46A) in bud, opening inside light yellow (11B), with a prominent blotch of dark red (59A) spots concentrated in the base, superimposed by a rather open dorsal flare of the same colour spots; outside light yellow (11B). Lvs oblong, obtuse at apex, rounded at base, semi-glossy and moderate olive green (147A) above. Shrub 1.2 × 1.2m in 10 years. Early May.
(The epithet deliberately uses Cowles in adjectival form, so this should not be corrected to the possessive form – Cowles’s)

(r) **‘Dad’s Bee’**

cv. *Elepidote rhododendron* : ‘Rosevallon’ (s) × ‘Grenadier’
H: W.E. Berg (1995), G: W.E. Berg (2000), N: C. Berg (2008), I: Dover Nursery, REG: C. Berg (2008)
Fls c.15/domed truss, funnel-campanulate, c.40 × c.40mm, with 5 wavy-edged lobes, strong red (46A) in bud, with some black spotting at the tips, opening the same colour (46A) throughout, with black nectar pouches and the throat spotted black. Calyx c.13mm long, strong red (46A). Lvs elliptic, 75 × 40mm, heavy-textured, broadly acute at apex, rounded at base, margins recurved, glossy and dark green above, dark reddish purple below (and as the plant tends to hold its leaves vertically this is a conspicuous feature), hairless. Shrub with an open habit, 0.9 × 0.9m in 10 years. Early March.

(r) **‘Dancing Embers’**

cv. *Elepidote rhododendron* : ‘Lem’s Tangerine’ (s) × ‘Fiery Orange’
H: J. Barlup (1998), G: J. Barlup (2003), N: J. Barlup (2007), REG: J. Barlup (2008)
Fls 13/domed truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, vivid red (45A) in bud, opening inside strong red (39A) at margins, blending to pale orange-yellow (16D) in the throat; outside strong red (39A) at margins, blending to pale orange-yellow (16D) towards the base; two moderate reddish orange (42D) flares on the dorsal lobe, 40mm from the base. Calyx 45mm long, strong red (39A), blending with pale orange-yellow (16D), with the three flares moderate reddish orange (42D), c.15mm from the base. Lvs elliptic, 110 × 40mm, broadly

acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 8 years. Late April.

(r) **'December Red'**

cv. *Elepidote rhododendron* : 'Ross Maud' (s) × 'Tortoiseshell Wonder'
H: D.J. Dosser (1990), G: D.J. Dosser (1998),
N: D.J. Dosser (2008), REG: D.J. Dosser (2008)
Fls 10/domed truss, funnel-shaped, 75 × 85mm, with 5 rounded, broadly overlapping, wavy-edged lobes, strong red (45D) in bud, opening strong red (45D) inside and out, unmarked. Lvs oblanceolate, 150 × 50mm, acute at apex, cuneate at base, margins decurved, matt green above, with light brown hairs below. Shrub 2.0 × 1.5m in 18 years. December (Australia).
(Nomenclatural standard: colour print provided by the registrant (WSY0101958))

(r) **'Dora Remotti'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 17–24/flattened hemispherical truss, funnel-shaped, 45 × 55–75mm, with 5 wavy-edged lobes, strong purplish red (61B) in bud, opening deep purplish pink (N74C), with a lighter (light purple, 75B) centre and dark red (59A) spots mostly on the dorsal lobe; stamens 10, included within the corolla, anthers strong reddish purple (72B), filaments white; pistil equalling or just exceeding the stamens and included within the corolla, style strong purplish red (58C) with a white apex, stigma dark red (187A). Lvs lanceolate, 120–140 × 40–45mm, acute at base and apex, margins flat, matt moderate olive green (147A) above, moderate yellow-green (146C) below; petioles 25–30mm long. Shrub 1.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101267). Etymology: named for the biological sciences graduate who carried out the research into safeguarding of the germplasm resources in the Parco Burcina)

(r) **'Dorothy Bishop'**

cv. *Elepidote rhododendron* : 'Hachmann's Charmant' (s) × 'Lem's Cameo'
H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2008), REG: J. Barlup (2008)
Fls 19/ball truss, broadly funnel-shaped, 75 × 80mm, with 5 very wavy-edged lobes, deep red (60A) in bud, opening inside to a blend of strong purplish pink (68B) and white (155D), with the white extending from the base to within c.15mm of the 68B border; outside strong purplish pink (68B), with white (155D) extending upwards c.15mm from the base; spotted dorsal flare of greyish red (184A) extending c.20mm from the base and additional spotting of moderate purplish red (184D) extending another 20mm. Calyx 25mm long, greyish red (184A), blending to white (155D), with strong purplish pink (68B) tips. Lvs elliptic, 105 × 45mm, broadly acute at

apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.4 × 1.8m in 12 years. Late May.

(Etymology: named for a Mrs Dorothy Bishop (aged 95), from England; the registrant and his wife are friends of Mrs Bishop's son and daughter-in-law)

(a) **'Dorothy Clark'**

cv. *Evergreen azalea* : 'Grace Freeman No. 2' (s) × 'Amagasa'
H: J.O. Harris (c.1978), G: J.O. Harris (c.1980),
N: J.O. Harris (c.1981), I: J.O. Harris (c.1981),
REG: J.O. Harris (2008)
Fls single, 2/flat truss, open funnel-shaped, c.55 × 65–115mm, with 5 wavy-edged lobes, strong purplish red (58B) in bud, opening very pale purple (69A), with a broad band of strong purplish red (58B) around the margins both inside and out; a blotch of light purplish pink (68D) spots extends along the inner edge of the 58B band on the dorsal lobe. Calyx c.8mm long, strong yellow-green (142A). Lvs obovate, 35 × c.10mm, broadly acute at apex, rounded at base, margins slightly upcurved, semi-glossy and bright green above, hairless. Shrub 0.9 × 0.9m in 9 years. Late April.
(Etymology: named for a friend of the registrant. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.367)

(r) **'Dotella'**

cv. *Elepidote rhododendron* : 'Dopey' (s) × 'Tarantella'
H: K.-H. Hübbers (1990), G: K.-H. Hübbers (1995),
N: K.-H. Hübbers (2002), REG: J. Westhoff (2008)
Fls 10/flat roundish truss, tubular funnel-shaped, 50 × 80mm, with 5 slightly wavy-edged lobes, inside deep to strong red (53A/B), outside deep red (185A), unmarked; style and filaments also red; the flower colour does not fade or burn in the sun. Calyx 2mm long, vivid red (46C). Lvs oblong-ovate, 100–150 × 30–60mm, mucronate at apex, truncate at base, margins slightly upcurved, matt green above, hairless. Shrub 1.1 × 1.1m in 10 years. Early May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101571). Etymology: the epithet is a combination of parts of the parental epithets. Introduced in 2006 by Glendoick Gardens (UK), and by Baumschule Neumann and Baumschule Ernst Stöckmann in Germany)

(r) **'Double Besse'**

cv. *Elepidote rhododendron* : 'Besse Howells', treated with colchicine
H: Briggs Nursery, G: Briggs Nursery (2005),
N: A. Emery (2008), I: Briggs Nursery (2005),
REG: Briggs Nursery (2008)
Fls hose-in-hose, 11–12/ball truss, open funnel-shaped, 40 × 65mm, with 4–5 very wavy-edged lobes in each of the two corollas, dark red (187B) in bud, opening inside deep red (60A), with a dark red (187A) blotch on the dorsal lobe; outside deep purplish red (187D). Lvs elliptic, 90–115 × 30–40mm, broadly

acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Shrub 0.9 × 0.9m in 10 years. Early May.

(a) **'Double Parfait'**

cv. Deciduous azalea : *occidentale* (SM 28-3) × *occidentale* 'Crescent City Double' (SM 28-2)
H: F.D. Mossman (1972), G: R.W. Cavender (1977), N: R.W. & K. Cavender & D. Meyer (2007), I: Briggs Nursery (2009), REG: R.W. Cavender (2008)
Fls 10–15/flat truss, tubular funnel-shaped, 40 × 50–55mm, varying from single (with 5 wavy-edged lobes) to semi-double or double (with 1–5 petaloid stamens, varying from fully petaloid to extremely deformed as well as narrow), buds variable in colour from deep purplish pink (55A) to pale yellow with deep purplish pink (55A) ribs, opening inside with very variable colouring: white, with dorsal lobe strong orange-yellow (17A) and other lobes with varying amounts of light purplish pink (55C); fully formed petaloid stamens (which may be the size of a normal corolla lobe) white, with 1–3 dorsal lobes strong orange-yellow (17A), the remaining petaloids white; outside white, the base of the tube deep purplish pink (55A), the dorsal lobe white and with some pale yellow and pink on the ribs, other lobes variable with deep purplish pink (55A) on the ribs and some 55A on all or on portions of the lobes; strongly scented. All flower forms may appear in the same truss but not necessarily on the same branch from year to year; the occurrence of the petaloids varies and they may not appear until the plant has flowered for several years. No normal calyx present. Lvs elliptic, 50–55 × 20–25mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above, hairless. Shrub 2.1 × 0.9m in 30 years. Mid May to mid June, with usually 5–6 fls open at any one time over a period of 7–10 days. (The SM numbers represent collections made by B.M. Smith and F.D. Mossman)

(r) **'Ed Hillary'**

cv. Elepidote rhododendron : *williamsianum* (s) × *grande*
H: B.W. Campbell (pre 1980), G: B.W. Campbell (1985), N: Dunedin Rhododendron Group (2008), I: Blue Mountain Nurseries (2008), REG: Dunedin Rhododendron Group (2008)
Fls 14/rounded truss, campanulate to broadly campanulate, 30–40 × 40–50mm, with 5 slightly wavy-edged lobes, buds vivid red (52A), opening inside and out ivory flushed strong red (52D) – or sometimes with the flushing just as a picotee edge; strong red (52A), or darker, blotch and spotting at the base of the dorsal throat. Calyx 2mm long, pale green. Lvs ovate to oblong, 85 × 35mm, cordate at base, obtuse and mucronate at apex, margins flat, matt green above, with a brown indumentum below when young which ages to light green; new growth often copper/brown. Shrub 1.5 × 1.5m in 10 years. September to October (New Zealand). (Nomenclatural standard: colour print provided by the registrant (WSY0101453). Etymology: named for the New Zealand mountaineer Sir Edmund Hillary who died in 2008)

(a) **'Edith Henderson'**

cv. Evergreen azalea : 'Banka' (s) × 'Target'
H: J.O. Harris (c.1978), G: J.O. Harris (c.1980), N: J.O. Harris (c.1981), I: J.O. Harris (c.1981), REG: J.O. Harris (2008)
Fls single, 1–2/flat truss, broadly funnel-shaped, c.85 × 95–115mm, with 5 wavy-edged lobes, deep yellowish pink (44D) in bud, opening inside deep yellowish pink (43C), with a dorsal blotch of strong red (46A) spots near the base of three lobes; outside strong yellowish pink (43D). Calyx c.10mm long, brilliant yellowish green (134C). Lvs elliptic, 65 × 25mm, between acute and broadly acute at apex, cuneate at base, margins flat, glossy and dark yellowish green (139A) above, hairless. Shrub with a dense habit, 1.1 × 1.1m in 9 years. Early May. (Etymology: named for a landscape architect from Atlanta, Georgia, USA. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.390)

(r) **'Elena Garibaldi'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: N. Furno (2008), REG: N. Furno (2008)
Fls 11–13/hemispherical truss, funnel-shaped, 33 × 50–53mm, with 5 slightly wavy-edged lobes, moderate purplish red to strong reddish purple (70A-B) in bud, opening deep purplish pink (68A), fading to white with age, with a conspicuous blotch and spots of dark red (187A) and deep purplish red (59B); stamens 10, exserted from the corolla, anthers moderate orange-yellow (164C), filaments white; pistil 4mm longer than the stamens, style light purplish pink (68D), stigma dark red (183A). Lvs lanceolate, 90–104 × 32–34mm, margins flat, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 13–14mm long. Shrub 1.0m. May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101321). Etymology: named for the Professor at the Faculty of Agriculture, Turin, who supervised the research to safeguard the rhododendrons of the Parco Burcina; the epithet uses her married name as there is already a plant named for her as 'Elena Accati')

(a) **'Ellie Harris'**

cv. Evergreen azalea : 'Sherwoodii' (s) × 'Fedora'
H: J.O. Harris (c.1971), G: J.O. Harris (c.1972), N: J.O. Harris (c.1973), I: J.O. Harris (c.1973), REG: J.O. Harris (2008)
Fls hose-in-hose, 3–6/domed truss, broadly funnel-shaped, c.60 × 65mm, with 5 wavy-edged lobes, strong purplish pink (62A) in bud, opening inside light purplish pink (62C) and changing to pale purplish pink (62D), with a faint dorsal blotch of small, strong purplish red (63D) spots; outside variable, light and pale purplish pink (62C-D); filaments pale pink, anthers dark; style bright pink, stigma yellow. Calyx c.10mm long, strong yellow-green (141D). Lvs elliptic, c.40 × c.20mm, obtuse at apex, cuneate at base, margins slightly upcurved, semi-glossy and dark bluish green (131A) above,

hairless. Shrub 1.1 × 0.9m in 10 years. Mid April. (Etymology: named for the wife of the registrant. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004. p.404)

(r) **'Emily Jayde'**

cv. Elepidote rhododendron : 'Doctor Peter Hewitt' (s) × 'Percy Wiseman'
H: D.J. Dosser (2001), G: D.J. Dosser (2006),
N: D.J. Dosser (2007), I: B.S. Coates (2007),
REG: D.J. Dosser (2008)
Fls 15/domed truss, tubular funnel-shaped, 75 × 100mm, with 5 wavy-edged lobes, buds vivid purplish red (57C), opening inside vivid purplish red (57C) and turning to white when fully out, outside very pale purple (69A); small, deep purplish pink (68A) spots on the dorsal lobe; stigma dark. Lvs oblanceolate, 125 × 50mm, matt green above, truncate at base, acute at apex, hairless. Shrub 0.6 × 0.6m in 5 years. October to November (Australia).
(Nomenclatural standard: colour print provided by the registrant (WSY0101032). Etymology: named for a granddaughter of the introducer)

(r) **'Emma and May'**

cv. Elepidote rhododendron : selection from *fortunei* S: Briggs Nursery (2004), G: Briggs Nursery (2007), N: Briggs Nursery (2007), I: Briggs Nursery (2008), REG: Briggs Nursery (2008)
Fls 4–12/domed truss, broadly funnel-shaped, 55 × 90mm, with 6–8 wavy-edged lobes, strong purplish pink (63B) in bud, opening very pale purple (75D) on lobes, fading to white; outside similar, with strong purplish pink (73B) veining; unmarked; scented. Lvs elliptic, 100–150 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (137A) above, hairless. Shrub 1.2 × 0.9m in 7 years. Mid May.
(Etymology: named for Emma Smith DeVoe and May Arkwright Hutton who helped Washington State women get the vote in 1910. Published in RareFind Nursery cat. (2008))

(r) **'Emma Williams'**

cv. Lepidote rhododendron : *yunnanense* (s) × Trewithen Orange Group
H: F.J. Williams (1968), G: P. Tregunna (1970s), N: F.J. Williams (c. 1985), I: Burncoose Nurseries (c.1987), REG: C.H. Williams (2008)
Fls 6–8/ rounded truss, greenish white in bud, opening light pinkish mauve throughout (in some years if the flowers open early they can be almost white or cream, only flushing pinkish mauve later). Shrub 2.0–2.5 × 2.0m in 20 years. April to May. Semi-deciduous in colder climates.
(Etymology: named for the former wife of Charles H. Williams)

(r) **'Enzo Piacenza'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)

Fls 18–21/hemispherical truss, funnel-shaped, 21 × 42–47mm, with 5 slightly wavy-edged lobes, deep purplish pink (72D) in bud, opening light purple to very pale purple (75B-D), with conspicuous, large spots of deep orange-yellow (163A) and deep greenish yellow (153A) on the dorsal lobe and just extending onto the adjacent lobes; stamens 10, exserted from the corolla, anthers dark purplish pink (186C), filaments white; pistil about 3mm longer than the stamens, style pale purplish pink (65D), stigma deep red (60A). Lvs oblanceolate, 90–115 × 47–50mm, margins not recurved, surface slightly undulate, matt moderate olive green (147A) above, lower surface moderate yellow-green (146D), with indumentum along the mid vein below; petioles 15–19mm long. Shrub 1.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101338). Etymology: named for the son of Felice Piacenza, the latter being the founder of the Parco Burcina)

(r) **'Ermanno De Biaggi'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 19–23/hemispherical truss, funnel-shaped, 30–40 × 55–60mm, with 5 smooth to slightly wavy-edged lobes, vivid purplish pink (N66A) in bud, opening deep purplish pink (N57C) inside and out, with deep greenish yellow (153A) spots on all 5 lobes but mostly on the dorsal lobes; stamens 9–10, included within the corolla, anthers moderate purplish red (64A), filaments vivid purplish red (67B); pistil 15mm longer than the stamens and exserted from the corolla, style strong purplish pink (63C) with a white base, stigma deep red (53A). Lvs lanceolate, 110–125 × 35–40mm, acute at base and apex, margins flat, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 15mm long. Shrub 3.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101270). Etymology: named for the city manager for Piedmont who has provided funds for research in the Parco Burcina)

(a) **'Fascination'** Syn. of 'Harris Fascination'

(r) **'Favaro'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 22–24/hemispherical truss, funnel-shaped, 37–50 × 47–60mm, with 5 slightly wavy-edged lobes, buds dark purple to strong purplish red (79B, 72A), opening strong reddish purple (72C) inside and out, with numerous dark purple (79A) spots mostly on the dorsal lobe; stamens 10, held inside the corolla, anthers light reddish brown (177B), filaments light purple (75B); pistil 8mm longer than the stamens and exserted from the corolla, style strong purplish red (71D), stigma dark red (187A). Lvs lanceolate to obovate, 102–124 × 30–42mm, margins flat but leaf surface

concave, acute at base and apex, matt dark yellowish green (139A) above, moderate olive green (147C) below; petioles 14–16mm long. Shrub 3.0m. May. (Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101272). Etymology: named for a place in Pollone, the village where the Parco Burcina is situated)

(r) **'Fay Norman'**

cv. Elepidote rhododendron : *campylocarpum* × *griffithianum*

H: Lady Norman (1937), G: Lady Norman, N: Mrs M. Gunn (2007), REG: Mrs M. Gunn (2008) Fls 8/open-topped truss, funnel-shaped, with 7–8 flat-edged, spreading lobes, 70–80 × 100mm, buds pale yellow (20D), with strong yellowish pink (37B) at apex and on veins, opening greenish white (157D) throughout, with a yellow-green flush towards the base; anthers moderate orange-yellow (c.164C); style pale yellow-green (1D), stigma strong yellowish green (N144A); scented. Calyx disk-shaped, green, with a slight glaucous hue. Pedicels strong red (47B) to moderate purplish red (59C), with a distinct glaucous bloom. Bud scales dark red (59A). Lvs ovate-elliptic, 120 × 45mm, margins flat, rounded at base, obtuse and slightly cuspidate at apex, hairless. Shrub 4.5–5.5 × 6m in 60 years. Mid May.

(Nomenclatural standard: herbarium specimen provided by the registrant (WSY0101114). Etymology: named for the registrant's grandmother, Lady Norman, daughter of the 1st Lord Aberconway).

(r) **'Felice Piacenza'**

cv. Elepidote rhododendron : parentage unknown G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)

Fls 10–15/conical truss, open funnel-shaped, 30–40 × 60–65mm, with 5 slightly wavy-edged lobes, buds light purple (75A), opening white, shading to very light purple (76B) at the margins, with a prominent blotch and spots of dark red (59A) mostly on the dorsal lobe; stamens 10, projecting from the corolla, anthers moderate orange-yellow (164C), filaments yellowish white (155D); pistil as long as the stamens, style pale greenish yellow (2D), stigma strong orange (N25A). Lvs lanceolate, 120–130 × 40–50mm, slightly twisted, margins flat, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (146D) below, hairless; petioles 15mm long. Shrub 2.5m. May.

(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101274). Etymology: named for the founder of the Parco Burcina)

(r) **'First Impression'**

cv. Elepidote rhododendron : 'Independence Day' (s) × 'Mrs Furnivall'

H: J. Barlup (1994), G: J. Barlup (1999), N: J. Barlup (2007), I: Nelson's Nursery (2006), REG: J. Barlup (2008)

Fls 21/ball truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, light purplish pink (62C) in

bud, opening inside white (155D), lightly tinged moderate purplish pink (62B) on margins, with a large, prominent flare of strong purplish red (60B) spots, mainly on the dorsal lobe, extending 40mm upwards from the base; outside white (155D), with midribs of dorsal lobe moderate purplish pink (62B) and outer edges strong purplish pink (62B). Calyx 15mm long, white (155D), edged moderate purplish pink (62B). Lvs elliptic, 125 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.8m in 12 years. Mid May.

(r) **'Forest Lane'**

cv. Elepidote rhododendron : 'Nancy Evans' (s) × 'Lem's Cameo'

H: K. Jones (1992), G: K. Jones (1997), N: K. Jones (2000), I: Forest Lane Nursery (2009), REG: C. Love (2008)

Fls 18–25/domed truss, funnel-campanulate, 50 × 75mm, with 5 wavy-edged lobes, moderate red (180C) in bud, opening inside light yellow (18B), with vivid reddish orange (30B) dorsal spotting; outside light yellow (18B), flushed with strong yellowish pink (32D). Calyx 25mm long, light yellow (14D), tipped with strong reddish orange (32B). Lvs elliptic, 125 × 50mm, broadly acute at apex, cuneate at base, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.1m in 10 years. Mid May.

(r) **'Fragrant Sensation'**

cv. Elepidote rhododendron : 'Horizon Monarch' (s) × 'Gargantua'

H: J. Barlup (1999), G: C. Blankenship (2004), N: C. Blankenship, I: Chuck's Nursery and Hammond Nursery, REG: C. Blankenship (2008)

Fls 16/domed truss, saucer-shaped, 70 × 100mm, with 7 very wavy-edged lobes, buds moderate yellow (160A), becoming light yellow (160B) with strong pink (50C) on outer edges, opening inside brilliant yellow (10A) in the throat, the lobes pale yellowish pink (27D), edged pale yellowish pink (36D); outside light yellow (10B) from the base to halfway up the lobes, with the apical half of the lobes light yellowish pink (36C); small, brush-like markings of vivid reddish orange (34B) at the base of the dorsal lobe; very thick and heavy substance; stigma fasciated in c.30% of fls; strongly scented. Calyx c.5–15mm long, brilliant yellow-green (154B), edged vivid reddish orange (34B). Lvs elliptic, 125–175 × 40–65mm, obtuse at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.4 × 1.8m in 9 years. Mid May.

(r) **'Frostlight'**

cv. Elepidote rhododendron : 'Janet Blair' (s) × 'Recital'

H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2008), REG: J. Barlup (2008)

Fls 18/ball truss, broadly funnel-shaped, 55 × 75mm, with 7 wavy-edged lobes, pale yellow (11D) in bud, opening inside and out white (155D), lightly tinged pale greenish yellow on the dorsal lobes; basal rays of

moderate red (181B), some extending as far as 20mm from the base. Calyx 40mm long, pale greenish yellow (10D), fading to white (155D), with a moderate red (181B) tinge extending c.15mm from the base. Lvs elliptic, 105 × 55 mm, broadly acute at apex, rounded at base, with wavy margins, dull and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.5m in 12 years. Mid May.

(r) **'Galinit'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)

Fls 14–17/tight hemispherical truss, funnel-shaped, 30–40 × 50–55mm, with 5 wavy-edged lobes, vivid reddish purple (N74A) in bud, opening light purple (76A) throughout, with numerous strong greenish yellow (151B) spots on the dorsal lobes; stamens 10, held within the corolla, anthers very light purple (75C), filaments strong purplish pink (68B); pistil equalling or shorter than the stamens, style light purplish pink (55C), stigma strong red (47B). Lvs lanceolate, 150–170 × 55–60mm, margins flat, acute at base and apex, matt moderate olive green (147A) above, strong yellow-green (144A) below, hairless; petioles 10–15mm long. Shrub 2.5m. May.

(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101276). Etymology: named after a site within the Parco Burcina)

(a) **'Georgia Giant'**

cv. Evergreen azalea : 'Moonbeam' × 'Lilacinum'
H: J.O. Harris (c.1971), G: J.O. Harris (c.1972),
N: J.O. Harris (c.1973), I: J.O. Harris (c.1973),
REG: J.O. Harris (2008)

Fls single, 2/flat truss, broadly funnel-shaped, c.100 × up to 140mm, with 5, very wavy-edged, reflexed lobes, white (155D) in bud, opening white (155D) throughout, with an inconspicuous dorsal blotch of pale yellow-green (157A) spots; slightly scented. Calyx c.10mm long, brilliant yellowish green (140B). Lvs elliptic, 50 × c.25mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and dark bluish green (133A) above, hairless. Shrub 1.4 × 1.1m in 8 years. Late April.

(Etymology: named for the size of the flowers and the State in which it originated. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.501)

(r) **'Giuseppe Bozzalla'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)

Fls 19–20/tight hemispherical truss, open funnel-shaped, 40 × 60–70mm, with 5 slightly wavy-edged lobes, buds strong purple (77B), opening light purple (84B) throughout, the dorsal lobe almost covered in dark red (187B-C) spots which just spread to the adjacent lobes; stamens 9–10, held within the corolla, anthers white, filaments strong reddish purple

(N78B); pistil 10mm longer than the stamens and exerted from the corolla, style white, stigma vivid purplish pink (N66A). Lvs elliptic-obovate, 100–120 × 40mm, margins flat, rounded to acute at base, broadly acute apex, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 15mm long. Shrub 2.0m. May.

(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101278). Etymology: named for a painter who resided in Biella (1874–1958))

(a) **'Gloria Still'**

cv. Evergreen azalea : 'Albert-Elisabeth' (s) × 'Fedora'
H: J.O. Harris (c.1978), G: J.O. Harris (c.1980),
N: J.O. Harris (c.1981), I: J.O. Harris (c.1981),
REG: J.O. Harris (2008)

Fls hose-in-hose, 4–8/truss, funnel-shaped, c.60 × 75mm, with up to 16 very wavy-edged lobes, moderate purplish pink (66D) in bud, opening inside and out deep purplish pink (66C), variegated randomly with white (155B); unmarked. Calyx c.10mm long, brilliant yellowish green (140B). Lvs elliptic, 40 × c.15mm, broadly acute at apex, cuneate at base, margins slightly upcurved, dull and strong yellow-green (143A) above, hairless. Shrub 0.6 × 0.9m in 6 years. Late April.

(Reported to be easy to force for Christmas.

Etymology: named for a friend of the registrant.

Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.519)

(r) **'Grand Recital'**

cv. Elepidote rhododendron : 'Mindy's Love' (s) × 'Jessie's Song'

H: J. Barlup (1999), G: J.D. Winberg (2006),
N: J. Barlup (2007), REG: J. Barlup (2008)

Fls 25/ball truss, broadly funnel-shaped, 65 × 95mm, with 5 wavy-edged lobes, strong pink (48C) in bud, opening inside and out pale greenish yellow (10D), with light yellow (10C) veining on the three dorsal lobes; with two speckled flares of deep purplish red (59B) on the dorsal lobe, extending upwards 25mm from the base. Calyx lobes irregular, 13–40mm long, pale greenish yellow (10D) to light yellow (10C) with deep purplish red (59B) flares 13–25mm long. Lvs elliptic, 135 × 65mm long, broadly acute at apex, rounded at base, margins flat, glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.1m in 6 years. Late April.

(r) **'Guido Piacenza'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)

Fls 12–20/hemispherical truss, funnel-shaped, 40 × 50–55mm, with 5 wavy-edged lobes, buds vivid purplish pink (N66A), opening deep purplish pink (N66D) throughout, with a dorsal blotch and spots of dark red (187C) extending from the base of the throat to half way up the dorsal lobe and just extending onto the adjacent lobes; stamens 10, held within the

corolla, anthers pale greenish yellow (1D), filaments white; pistil equalling or exceeding the corolla and 10–15mm longer than the stamens, style white, stigma moderate reddish orange (35B). Lvs oblanceolate, 195–215 × 50–70mm, margins flat, cuneate at base, acute to broadly acute and mucronate at apex, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 35mm long. Shrub 2.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101280). Etymology: named for the current president in charge of the Parco Burcina, a direct descendent of the founder)

(a) **‘Harris Fascination’**

cv. Evergreen azalea : ‘Grace Freeman No. 2’ (s) × ‘Amagasa’
H: J.O. Harris (c.1971), G: J.O. Harris (c.1972), N: J.O. Harris (c.1973), I: J.O. Harris (c.1973), REG: J.O. Harris (2008)
Fls single, 3/domed truss, broadly funnel-shaped, c.80 × 75–115mm, with 5 very wavy-edged lobes, deep purplish pink (58D) in bud, opening inside and out moderate purplish pink (62B), edged deep purplish pink (58D); unmarked. Calyx c.6mm long, brilliant yellowish green (140B). Lvs elliptic, 35 × c.15mm, broadly acute at apex, cuneate at base, semi-glossy and dark bluish green (131A) above, hairless. Shrub with a dense habit, 0.8 × 0.9m in 9 years. Late April.
Syn: ‘Fascination’
(Revised entry: previously listed as ‘Fascination’ (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.438. The prefix Harris is used here in an adjectival not possessive form)

(a) **‘Harris Parfait’**

cv. Evergreen azalea : ‘Sherwoodii’ (s) × ‘Fedora’
H: J.O. Harris (c.1970), G: J.O. Harris (c.1972), N: J.O. Harris (c.1973), I: J.O. Harris (c.1973), REG: J.O. Harris (2008)
Fls single, 4–6/flattened ball truss, open funnel-shaped, 45 × 55mm, with 5 flat-edged lobes, light purple (75B) in bud, opening inside and out white (155B), slightly tinged light purplish pink (63D), with a dorsal blotch of large, deep purplish pink (61A) spots on three lobes; slightly scented. Calyx c.20mm long, strong yellow-green (142A). Lvs elliptic, 30–40 × c.10mm, broadly acute at apex, cuneate at base, with slightly downcurved margins, glossy and dark yellowish green (139A) above, hairless. Shrub 1.1 × 1.1m in 10 years. Mid April.
Syn: ‘Parfait’
(Revised entry: previously listed as ‘Parfait’ (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.1035. The prefix Harris is used here in an adjectival not possessive form)

(a) **‘Harris Vibrant’**

cv. Evergreen azalea : ‘Grace Freeman No. 2’ (s) × ‘Amagasa’
H: J.O. Harris (c.1971), G: J.O. Harris (c.1972), N: J.O. Harris (c.1973), I: J.O. Harris (c.1973), REG: J.O. Harris (2008)
Fls single, 2/flat truss, funnel-shaped, c.60 × 55–100mm, with 5 wavy-edged lobes, opening white (155B) throughout, with a broad band of vivid purplish red (57B) around the margins, both inside and out; marked with a faint blotch of tiny, deep purplish pink (58D) speckles on the dorsal lobe. Calyx c.10–15mm long, light olive (152B). Lvs elliptic, 45 × 25mm, broadly acute at apex, cuneate at base, margins slightly upcurved, semi-glossy and strong yellow-green (144A) above, hairless. Shrub 0.6 × 0.9m in 9 years. Mid April.
Syn: ‘Vibrant’
(Revised entry: previously listed as ‘Vibrant’ (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.1414. The prefix Harris is used here in an adjectival not possessive form)

(a) **‘Harry’s Ruby’**

cv. Evergreen azalea : parentage unknown
R: unknown (pre 1988), REG: RHS Garden, Rosemoor (2008)
Fls 2(–3)/upright truss, funnel-shaped, 30 × 34mm, with 5 flat to slightly wavy-edged lobes, deep to strong red (53A–C) throughout, with inconspicuous red dorsal markings; filaments and style deep to strong red (53A–C); filaments with short white hairs in basal half. Calyx 3mm long, yellow-green, hairy. Pedicels up to 9mm long, flushed red. Lvs mostly elliptic to ovate (but very variable), up to 30 × 15mm, margins flat, subacute or obtuse and slightly apiculate at apex, mostly cuneate at base (but some rounded or oblique) slightly glossy to matt green above, with a few hairs on the midrib below. Broadly upright shrub, up to 1.45 × 1.7m in c.20 years. April to May.
(Nomenclatural standard: herbarium specimen provided by the registrant (WSY0101168).
Etymology: believed to have been named for Harry Smith (1911–1971) who was a gardener at Rosemoor under Lady Anne Palmer (now Lady Anne Berry))

(r) **‘Hey There’**

cv. Elepidote rhododendron : ((Tally Ho Group × *yakushimanum*) × (*yakushimanum* × ‘Noyo Chief’)) (s) × ‘The Honourable Jean Marie de Montague’
H: R.A. Furman (1986), G: R.A. Furman (1990), N: R.A. Furman (2008), REG: R.A. Furman (2008)
Fls 21/ball truss, open funnel-shaped, 60 × 75mm, with 6 wavy-edged lobes, strong red (53B) in bud, opening inside with a strong purplish red (54A) edge, shading to strong purplish pink (55B) and with strong red (55B), dorsal spotting; outside deep purplish pink (55A); slightly scented. Lvs elliptic, 50–115 × 25–40mm, broadly acute at apex, cuneate at base, margins wavy, glossy and dark green above, hairs on the lower surface white when young, aging to light tan. Shrub with an exceptionally dense habit, the foliage hugging the ground, 1.5 × 1.8m in 22 years. Early to mid May.

(r) **'Holi'**

cv. *Elepidote rhododendron* : 'Admiral Vanessa' (s) × 'Welse'

H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2007), N: Mrs L. Hübbers (2008), I: Baumschulen Wüstemeyer, REG: J. Westhoff (2008)

Fls 12/nearly rounded truss, funnel-shaped, 60 × 100mm, with 5 wavy-edged lobes, inside very pale purple (76C) in the throat, shading to very light purple (76B) at the margins, outside very pale purple (69A/B); densely spotted dark red (187A-C), mostly on four fifths of the dorsal lobes, with small lines of dense spots in the middle of all the lobes. Calyx 25mm long, cut into several irregular lobes, coloured like the corolla and densely spotted. Lvs oblong, 110 × 50mm, cuneate at tip with a mucronate apex, roundish at base, margins downcurved, matt green above, hairless. Shrub 0.5 × 0.4m in 5 years. Late May to early June.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101574). Etymology: the epithet is the name of a Hindu holiday)

(r) **'Hong Yun'**

cv. *Elepidote rhododendron* : *irroratum* (s) × *arboreum* subsp. *delavayi*

H: C. Zhang & B. Feng (1985), G: C. Zhang & B. Feng (2002), N: C. Zhang, B. Feng & L. Gao (2002), REG: C. Zhang, B. Feng & L. Gao (2008)

Fls 8–10/rounded, racemose-umbellate truss, tubular-campanulate, 40 × 40mm, with 5 slightly wavy-edged lobes, baby pink in bud, opening inside and out creamy yellow on the tube and pink on the lobes, with irregular, mauve, dorsal spotting. Calyx 3mm long, light green. Lvs elliptic to elliptic-lanceolate, 115 × 38mm, obtuse at apex, broadly cuneate at base, margins flat, hairy below (both gland-tipped and eglandular hairs). Shrub 0.6–1.0 × 1.0–1.5m in 20 years. March to April.

(Nomenclatural standard: colour print of an electronic image supplied by the registrants (WSY0101887))

(r) **'Ibykus'**

cv. *Elepidote rhododendron* : 'Paprika Spiced' (s) × 'Märchenland'

H: K.-H. Hübbers (1990), G: K.-H. Hübbers (1997), N: O. Tschetsch (2008), REG: J. Westhoff (2008)

Fls 20/flattened globe truss, funnel-shaped, 65 × 90mm, with 5 flat-edged lobes, strong purplish red (54A) in bud, opening inside white (I55C) and outside white (I55A); the whole of the throat densely spotted strong red (53C), more intense in the dorsal part. Calyx 2–5mm long, reddish. Lvs oblong, 100–170 × 35–60mm, acute at apex, cuneate at base, margins flat, matt green above, hairless. Shrub 1.0 × 1.0m in 10 years. Early to mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101576). Etymology: alludes to the ballade by Friedrich Schiller "Die Kraniche des Ibykus" – a word play with the name of the registrant's city, Kranenburg, and kranich (meaning stork): Ibykus was a Greek poet)

(a) **'Isabella Maria'**

cv. Evergreen azalea : 'Carol Kittel' (s) × 'Girard's Fuchsia'

H: J.F. Klimavicz (1998), G: J.F. Klimavicz (2000), N: J.F. Klimavicz (2007), REG: J.F. Klimavicz (2008)

Fls 3/ball truss, saucer-shaped, 40 × 50mm, with 5 wavy-edged lobes, strong to light purple (N80B-C) in bud, opening inside white, with broad, strong to light purple (N80B-C) margins and slight dorsal spotting of the same colour; outside similarly coloured. Calyx 6mm long, moderate yellow-green (138B). Lvs obovate, 50 × 20mm, obtuse at apex, cuneate at base, margins flat, moderate olive green (146A) above, hairless. Shrub with a dense habit, 0.45 × 0.45m in 5 years. Mid May.

(Etymology: named for a member of the registrant's family)

(a) **'Itty Bitty'**

cv. Evergreen azalea : ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'

H: W.L. Clagett (1996), G: W.L. Clagett (2002), N: W.L. Clagett (2008), I: White's Nursery (2009), REG: W.L. Clagett (2008)

Fls hose-in-hose and semi-double, 3/flat truss, broadly funnel-shaped, 30 × 30mm, with 15–19 wavy-edged lobes, deep purplish pink (68A) in bud, opening inside deep purplish pink (68A) at edge of each lobe and fading at centre of lobes to paler pink or to nearly white; outside deep purplish pink (68A), with paler pink or nearly white midribs; some red spotting on the dorsal lobes. Lvs elliptic, 25 × 15mm, broadly acute at apex, cuneate at base, with flat margins, glossy and moderate olive green (147A) above, with fine white hairs below when young, later hairless. Shrub 0.4 × 0.3m in 11 years. Mid May.

(r) **'James Colville'**

cv. *Elepidote rhododendron* : (parentage unknown, but possibly an *arizelum* × *maccabeanum* derivative)

H: J.A.R. Moffat (c. 1970), G: J.A.R. Moffat (c.1980), N: Mrs F.E. Colville (2008), REG: Mrs F.E. Colville (2008)

Fls 10–15/rounded truss, ventricose-campanulate, 40 × 35mm, with 8 wavy-edged lobes, pale yellow in bud, opening very pale lemon yellow both inside and out, with purple rays at the base of the corolla (longest and broadest around the dorsal side of the corolla). Calyx 4mm long, brownish green. Lvs oblanceolate, up to 260 × 100mm, margins decurved, matt green above, cuneate at base, obtuse and very shortly mucronate at apex, lower surface with a two layered hairy indumentum, the lower layer greyish, the upper layer light brown when young and aging to rusty brown. Shrub 4.5 × 4.5m in 30 years. March to April.

(Nomenclatural standard: colour print provided by the registrant (WSY0101024). Etymology: named for the husband of the registrant)

(a) **'James Scott'**

cv. Evergreen azalea : 'Mary Ellen Thomsen' (s) × 'Florence Waldman'
H: J.F. Klimavicz (1997), G: J.F. Klimavicz (1999), N: J.F. Klimavicz (2007), REG: J.F. Klimavicz (2008)
Fls 2–3/ball truss, saucer-shaped, hose-in-hose and double, 30 × 50mm, with 19–24 wavy-edged lobes, buds white with sectors and flecks of deep purplish pink (N74C), opening inside and out white, with many deep purplish pink (N74C) sectors and flecks, the throat pale yellow-green (157B) and unmarked. No normal calyx present. Lvs elliptic, 30 × 13mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above, hairless. Shrub with a dense habit, 0.45 × 0.45m in 5 years. Mid May.
(Etymology: named for a member of the registrant's family)

(a) **'Jay Valentine'**

cv. Evergreen azalea : 'Martha Hitchcock' (s) × 'Amagasa'
H: J.O. Harris (c.1995), G: J.O. Harris (c.1996), N: J.O. Harris (c.1998), I: J.O. Harris (c.1998), REG: J.O. Harris (2008)
Fls single, 2/truss, open funnel-shaped, c.50 × 70mm, with 5 wavy-edged lobes, vivid red (45A) in bud, opening vivid red (45A) throughout; dorsal blotch of small, strong red (46A) spots. Calyx c.6mm long, moderate yellow-green (139D). Lvs elliptic, c.30 × 20mm, broadly acute at apex, cuneate at base, margins slightly upcurved, glossy and deep yellowish green (141B) above, hairless. Shrub 0.9 × 1.5m in 6 years. Early May.
(Etymology: named for the grandson of James and Ellie Harris who was born on Valentine's Day)

(r) **'Jean E. Potente'**

cv. Elepidote rhododendron : 'Anne Hardgrove' (s) × *yakushimanum*, Exbury form
H: R. Murcott (1971), G: R. Murcott (1980), N: R.J. Potente (2008), REG: R.J. Potente (2008)
Fls 15/conical truss, open funnel-shaped, 40 × 40mm, with 6 wavy-edged lobes, vivid reddish orange (42A) in bud, opening inside with a vivid reddish orange (43A) throat, shading to deep pink (48C) on the lobes; outside deep pink (48C); unmarked. Lvs elliptic, 140 × 40mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and brilliant green (128A) above, hairless. Shrub 1.5 × 1.5m in 9 years. Midseason.
(Etymology: named for the late wife of the registrant)

(r) **'Jean Leggett'**

cv. Elepidote rhododendron : 'Charlotte de Rothschild' (s) × 'Alice Hennessy'
H: T.A. Garbutt (1994), G: J. Leggett (2006), N: T.A. Garbutt (2007), REG: T.A. Garbutt (2008)
Fls 13/ large, compact truss, funnel-shaped, 70 × 120mm, with 7 wavy-edged lobes, pink in bud, opening white flushed pink around the margins both inside and out; unmarked, but with yellow nectaries; not scented. Calyx 3mm long, pink. Lvs lanceolate, 170 × 70mm, acute at apex, rounded at base, margins

wavy, matt green above. Shrub 1.5 × 1.0m in 5 years. November (New Zealand).

(Nomenclatural standard: colour print provided by the registrant (WSY0101021). Etymology: named for a friend of the registrant)

(r) **'Jessie E. Lewis'**

cv. Elepidote rhododendron : 'Anna' (s) × ('Holy Moses' × Albatross Group)
H: E.F. Newcomb (1981), G: L.L. & E.F. Newcomb (1985), N: C. Blankenship (2007), I: Chuck's Nursery and Hammond Nursery (2009), REG: C. Blankenship (2008)
Fls 10–13/ball truss, open funnel-shaped, 70–75 × 120–145mm, with 7 very wavy-edged lobes, strong purplish red (58B) in bud, opening inside strong purplish pink (63C) at margins, changing to pale purplish pink (62D) towards the centre, with a moderate purplish red (59A) basal blotch, surrounded by a continuous, strong red flare, radiating about 25mm into all lobes; outside strong purplish pink (63C); strongly scented. Calyx c.0.1mm long, vivid red (45A). Lvs elliptic, 115–165 × 50–65mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.5 × 2.0m in 10 years. Early May.
(Etymology: named for the late grandmother of the registrant)

(a) **'Jiao Yan'**

cv. Evergreen azalea : sport from *simsei* 'Zi Yan'
G: C. Zhang & L. Gao (1999), N: C. Zhang, L. Gao & Y. Huang (2006), I: Stock Co. of Bioscience and Technology of Yunnan Green Land (2007), REG: C. Zhang, L. Gao & Y. Huang (2008)
Fls 3–5/truss, shaped like a camellia, with petaloid stamens, 38 × 80mm, with 5 flat-edged outer lobes (divided almost to base), peach pink in bud, opening peach pink inside and out, with brick red spots. Calyx petaloid. Lvs elliptic-ovate, 35–60 × 15–20mm, shortly acuminate at apex, cuneate at base, hairy (hairs pale brown at first, aging to deep brown). Shrub 0.4–0.6 × 1.0m in 8 years. April to May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101889). Sport first seen in 1997, propagated that year and first flowered as an independent plant in 1999)

(a) **'Jin Zhi Zhu'**

cv. Deciduous azalea : seedling from *molle*
G: C. Zhang (2003), N: C. Zhang & L. Gao (2006), REG: C. Zhang & L. Gao (2008)
Fls 7–13/ open, racemose-umbellate truss, funnel-shaped, 40–50 × 40–60mm, with 5 flat-edged lobes, pale yellow in bud, opening yellow inside and saffron orange outside, with red spots on the dorsal lobe; scented. Calyx 5–7mm long, green. Lvs oblong, 50–100 × 20–40mm, mucronate at apex, cuneate at base, matt green above, hairy (hairs grey white at first, aging to grey). Shrub 0.8–1.0 × 0.5m in 23 years. April to May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101888))

(r) **'Joan Elder'**

cv. *Elepidote rhododendron* : 'Lem's Cameo' (s) × 'Autumn Gold'
H: T.A. Garbutt (1994), G: J. Elder (2006),
N: T.A. Garbutt (2007), REG: T.A. Garbutt (2008)
Fls 23/fairly compact, hemispherical truss, funnel-shaped, 66 × 80mm, with 7 very wavy-edged lobes, red in bud, opening inside deep cream flushed red, outside deep pink with broad red stripes; with no markings but with red nectaries; not scented. Calyx 10mm long, red. Lvs obanceolate, 110 × 45mm, margins flat, cuneate at base, acute at apex, hairless. Shrub 0.75 × 0.75m in 6 years. Mid November (New Zealand).
(Nomenclatural standard: colour print provided by the registrant (WSY0101022). Etymology: named after a friend of the registrant)

(r) **'Judy's Sunrise'**

cv. *Elepidote rhododendron* : 'Deidre Hunt' (s) × 'Bryan H. Tonkin'
H: D.J. Dosser (2000), G: D.J. Dosser (2007),
N: D.J. Dosser (2007), I: J. Barratt (2007),
REG: D.J. Dosser (2008)
Fls 11/ball truss, funnel-shaped, 75 × 75mm, with 6 very wavy-edged lobes, buds strong purplish red (64C), opening inside pale purplish pink (62D) and turning to white, outside light purplish pink (62C); dorsal lobe with scattered, small, strong purplish red (64C) spots; dark ring around the base of the tube; style pale green, stigma large and conspicuous; slightly scented. Lvs oblanceolate, 150 × 64mm, glossy green above, truncate at base, broadly acute at apex, hairless. Shrub 0.8 × 0.48m in 7 years.
(Nomenclatural standard: colour print provided by the registrant (WSY0100797). Etymology: named for Judy Barratt who won the plant in a charity raffle held by the registrant)

(r) **'Kala Daag'**

cv. *Elepidote rhododendron* : 'Mondnacht' (s) × 'Hachmann's Kabarett'
H: K.-H. Hübbers (2001), G: K.-H. Hübbers (2006),
N: Mrs L. Hübbers (2007), I: Baumschule Ernst Stöckmann and Baumschulen Wüstemeyer (2008),
REG: J. Westhoff (2008)
Fls 15/roundish, flattened truss, funnel-shaped, 60 × 100mm, with 7 wavy-edged lobes, pink in bud, opening inside white (155B) and outside white (155C), with a dark red (187B) blotch in the throat shading out into dense spotting that covers three quarters of the three dorsal lobes; stigma pale green. Calyx 5–20mm long, moderate orange-yellow (168D). Lvs oblong-elliptic, 90–150 × 30–65mm, mucronate at apex, cuneate/rounded at base, margins slightly wavy-edged, glossy green above, hairless. Shrub 0.6 × 0.6m in 5 years. Early to mid May.
(Nomenclatural standard: colour print of an electronic image supplied by the registrant (WSY0101582). Etymology: the epithet means "black spot" in Hindi)

(r) **'Kali'**

cv. *Elepidote rhododendron* : 'Nova Zembla' (s) × 'Rubicon'
H: K.-H. Hübbers (1993), G: K.-H. Hübbers (1998),
N: Mrs L. Hübbers (2005), I: Glendoick Gardens & Baumschule Neumann (2006), REG: J. Westhoff (2008)
Fls 15/nearly rounded truss, funnel-shaped, 50 × 60mm, with 5 smooth-edged lobes, dark red (187A) in bud, opening inside deep purplish red (61A) in the throat and shading out to dark red (187A) at the margins, outside dark red (187C); dense spotting of strong pink (49A) extends from the throat and covers two-thirds of the dorsal lobe; yellowish white anthers. Lvs elliptic, 70–110 × 20–40mm, acute at apex, cuneate at base, margins flat, matt green above, hairless. Shrub 1.0 × 0.9m in 10 years. Early to mid May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101583). Etymology: named for a Hindu goddess)

(r) **'Kenneth Wilson'**

cv. *Elepidote rhododendron* : 'John Dosser' (s) × 'Petra's Debut'
H: D.J. Dosser (1992), G: D.J. Dosser (2004),
N: D.J. Dosser (2007), REG: D.J. Dosser (2008)
Fls 18/ball truss, funnel-shaped, 40 × 75mm, with 5 very wavy-edged lobes, buds vivid purple (80A), opening inside and out strong purple (80B), with a conspicuous, large, deep purplish red (59B) blotch on the dorsal lobe. Lvs lanceolate, 125 × 40mm, glossy green above, cuneate at base, acute at apex, hairless. Shrub 1.5 × 2.1m in 15 years. October to November (Australia).
(Nomenclatural standard: colour print provided by the registrant (WSY0100799). Etymology: named for a cousin of the registrant)

(a) **'Kita-no-hakuhō'**

cv. Evergreen azalea : (*kinsianum*, white-flowered × 'Karafune') (s) × 'Morioka-koyuki'
H: T. Saitō (1999), G: T. Saitō (2004), N: T. Saitō (2006), REG: T. Saitō (2008)
Fls hose-in-hose, 2–4/open truss, funnel-shaped, 28–30 × 25mm, with 5+5 smooth-edged lobes, buds white, flushed green, opening inside pure white, flushed green in the throat at first, outside light creamy white; unmarked. Calyx as corolla. Lvs oblong-elliptic, 37 × 18mm, margins flat, obtuse at base and apex, matt green above, with persistent white hairs. Shrub 40 × 30cm in 7 years. Mid May.
(Nomenclatural standard: colour print provided by the registrant (WSY0101036))

(r) **'Kranenburg'**

cv. *Elepidote rhododendron* : 'Lem's Monarch' (s) × 'Odessa'
H: K.-H. Hübbers (1987), G: K.-H. Hübbers (1992),
N: K.-H. Hübbers (2000), REG: J. Westhoff (2008)
Fls 10–12/flat, roundish truss, funnel-shaped, 70 × 110mm, with 5 wavy-edged lobes, buds vivid red (52A), opening inside strong purplish red (58C) in the throat, shading to vivid purplish red (57D) at the margins,

outside vivid purplish red (57B-D); vivid reddish orange (40A) spots in two lines on the dorsal part of the throat. Calyx 1–3mm long, vivid red (44A). Lvs ovate, 85–150 × 45–60mm, mucronate at apex, slightly cuneate at base, margins flat, matt green above, hairless. Shrub 1.5 × 1.5m in 10 years. Mid May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101589). Etymology: named after the city in the Lower Rhine region of Germany where the registrant lives)

(a) **‘Kranenfee’**

cv. Deciduous azalea : ‘Raimunde’ (s) × ‘Twilight Sky’

H: K.-H. Hübbers (1989), G: K.-H. Hübbers (1995), N: K.-H. Hübbers (2006), REG: J. Westhoff (2008) Fls up to 13/rounded truss, tubular funnel-shaped, 50 × 70mm, with 5 very wavy-edged lobes, buds strong purplish pink (62A), opening inside very pale purple (69A), outside strong to light purplish pink (68B-D); with strong orange-yellow (17A), dense, blotch-like spotting on three quarters of the dorsal lobe; intensely scented. Calyx 3mm long, light green. Lvs elliptic, 70 × 40mm, cuneate at apex, with a mucronate tip, cuneate at base, margins flat, hairless. Shrub 1.0 × 0.8m in 10 years. Mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101593).

Etymology: the epithet is a combination of Kranenburg (where it was raised) and the German for fairy, in allusion to the fairy-like flowers)

(r) **‘Kranengold’**

cv. Elepidote rhododendron : ‘Barbara Reuthe’ (s) × ‘New Comet’

H: K.-H. Hübbers (1989), G: K.-H. Hübbers (1995), N: K.-H. Hübbers (2000), REG: J. Westhoff (2008) Fls up to 9/very slightly hanging, loose truss, funnel-shaped, 80 × 110mm, with 7 wavy-edged lobes, buds moderate orange-yellow (164C), opening inside brilliant to pale yellow (11A-C), outside brilliant to light yellow (10A-C); two lines of moderate yellow-green (147B) spots on the dorsal part of the throat; style green, densely hairy. Calyx with ventral lobes 2mm long, dorsal lobes 10mm long, brilliant yellow (8A). Lvs oblong, 100–150 × 45–75mm, mucronate at apex, truncate at base, margins flat, matt green above, hairless. Shrub 1.5 × 1.5m in 10 years. Early to mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101599).

Etymology: epithet a combination of part of the name of the city where it was raised (Kranenburg) and the colour of the flowers. Introduced in Germany in 2006 by Baumschule H. Hachmann, Baumschulen Wüstemeyer and Baumschule Neumann)

(r) **‘Kranenpepper’**

cv. Elepidote rhododendron : ‘Paprika Spiced’ (s) × ‘Märchenland’

H: K.-H. Hübbers (1990), G: K.-H. Hübbers (1996), N: K.-H. Hübbers (2005), REG: J. Westhoff (2008) Fls 16/flattened globe truss, funnel-shaped, 60 × 80mm, with 5 wavy-edged lobes, buds strong purplish

red (63A), opening inside with a brilliant yellow (11A) throat, shading out to pale purplish pink (65D) and then to moderate purplish pink (65A) at the margins, outside deep purplish pink (58D); intensely spotted with moderate reddish orange (173B) on the three dorsal lobes. Calyx 6mm long, yellowish green. Lvs obovate, 80–120 × 30–45mm, mucronate at apex, cuneate at base, margins flat, matt green above, hairless. Shrub 0.6 × 0.7m in 10 years. Mid to late May. Said to be resistant to bud blast.

(Nomenclatural standard: colour print of an electronic image supplied by the registrant (WSY0101604). Etymology: the epithet is a combination of part of the name of the city where it was raised (Kranenburg) and an allusion to the seed parent. The plant was introduced in 2006 by Glendoick Gardens in Scotland and by Baumschule Neumann, Baumschule Ernst Stöckmann and Baumschulen Wüstemeyer in Germany)

(r) **‘Kranenstar’**

cv. Elepidote rhododendron : ‘Lem’s Cameo’ (s) × ‘Deenah’

H: K.-H. Hübbers (1992), G: K.-H. Hübbers (1997), N: K.-H. Hübbers (2000), REG: J. Westhoff (2008) Fls 15/loose, flattened globe truss, funnel-shaped, 50 × 60mm, with 5 wavy-edged lobes, inside moderate purplish pink (62B) at the margins, shading to white in the throat, outside moderate purplish pink (62B), with a strong purplish red (63B) stripe on each lobe; dorsal lobe with a pale greenish yellow (1D) blotch which is covered in strong yellow-green (144C) spots. Calyx 5mm long, greenish, with a reddish rim. Lvs oblong, 80–140 × 30–60mm, mucronate at apex, cuneate at base, margins flat, matt green above, hairless. Shrub 1.0 × 1.1m in 10 years. Early to mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101605).

Etymology: the epithet is derived in part from the name of the city in which it was raised (Kranenburg))

(r) **‘Lavender Satin’**

cv. Elepidote rhododendron : ‘Jonathan Shaw’ (s) × ‘Plum High’

H: J. Barlup (2000), G: J. Barlup (2004), N: J. Barlup (2007), REG: J. Barlup (2008)

Fls 18/ball truss, broadly funnel-shaped, 50 × 80mm, with 5 wavy-edged lobes, moderate purplish red (70A) in bud, opening inside very pale purple (76D) in the centre, shading to very light purple (76B) and growing darker on the lobe tips from light purple (77C) on the ventral lobes to strong purple (77B) on the margins of the dorsal lobes, with a large, heavily spotted, moderate red (183C) dorsal flare extending 40mm from the base; outside very light purple (76B), with strong purple (77B) midveins and margins. Calyx minute. Lvs elliptic, 120 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub with a dense habit, 0.6 × 0.9m in 7 years. Late May.

(a) **'Lee Thomas'**

cv. Evergreen azalea : 'Banka' (s) × 'Target'

H: J.O. Harris (c.1978), G: J.O. Harris (c.1980),

N: J.O. Harris (c.1981), I: J.O. Harris (c.1981),

REG: J.O. Harris (2008)

Fls single, solitary, broadly funnel-shaped, c.35 × 65–125mm, with 6 wavy-edged lobes, moderate purplish pink (62B) in bud, opening light purplish pink (62C) throughout; small, strong reddish orange (42B) spots covering parts of the three dorsal lobes. Calyx c.6mm long, light bluish green (128C). Lvs elliptic, c.40 × c.15mm, broadly acute at apex, cuneate at base, margins slightly downcurved, semi-glossy and dark yellowish green (139A) above, hairs on the lower surface white at first (155C), aging to moderate olive brown (199A). Shrub 0.5 × 0.85m in 10 years. Early May.

(Etymology: named for a member of the former Ralph Pennington chapter of the American Azalea Society. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.767)

(r) **'Lockington's Dawn'**

cv. Elepidote rhododendron : 'Marcus of Lockington'

(s) × 'Australian Sunset'

H: D.J. Dosser (1999), G: D.J. Dosser, N: D.J. Dosser,

REG: D.J. Dosser (2008)

Fls 10/ball truss, broadly funnel-shaped, 70 × 100mm, with 7 wavy-edged lobes, strong red (53C) in bud, opening strong red (53C) on lobes and shading to light yellow (18A) in the throat; outside strong red (53B), shading to strong red (53C); prominent rays of strong red (53C) all around the base of the throat; slightly scented; stigma green. Calyx 25mm long, light yellow (18A). Lvs 125 × 50mm, broadly acute at apex, cuneate at base, margins decurved, hairless. Shrub 1.0 × 0.5m in 9 years. November (Australia). (Nomenclatural standard: colour print provided by the registrant (WSY0101955))

(v) **'Lois Kruppa'**

cv. Lepidote rhododendron : believed to be the same cross that produced 'Santa Lucia' i.e. 'Kurt Herbert Adler' × *leucogigas*

G: Pacific Island Nursery (2007), N: R. Kruppa

(2008), I: Pacific Island Nursery, REG: R. Kruppa

(2008)

Fls 4–5/flat or lax truss, tubular funnel-shaped, 90 × 32mm, with 7 wavy-edged lobes, pink in bud, opening inside moderate purplish pink (62B) at margins, shading to pale purplish pink (56A) towards the throat; outside moderate purplish pink (62B), deepening to deep pink (51C) at the base; unmarked; filaments red; flower colour darkens with age; slightly scented. Lvs elliptic, 115–150 × 38–57mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and moderate olive green (137A) above, moderate yellow-green (146C) below, with tiny dark scales (visible with a x10 lens) on the lower surface which has a yellow-green midvein. Shrub with a tidy habit of growth, 1.0 × 0.9m in 7.5 years. November to January and July to August.

(Similar to 'Santa Lucia' but (when grown together

for comparison) with much deeper-coloured flowers, leaves of a darker green, more vigorous and holding on to more of its leaves. Probably hybridised by the late Pete Sullivan; cuttings/rooted cuttings sent to Pacific Island Nursery from his "church garden" by both Chip Lima and Bill Moynier. Etymology: named for the late wife of the registrant)

(r) **'Lorenzo Delleani'**

cv. Elepidote rhododendron : parentage unknown

G: Parco Burcina, S: E. Accati & N. Furno (2000),

N: E. Accati & N. Furno (2008), REG: E. Accati &

N. Furno (2008)

Fls 14–17/conical truss, funnel-shaped, 40 × 55–65mm, with 5 wavy-edged lobes, buds deep purplish pink (73A), opening deep purplish pink (73A), with a very light/very pale purple (75C-D) centre; relatively inconspicuous spotting of brilliant yellow/vivid orange-yellow (13B/23A) on the dorsal lobe and just extending onto adjacent lobes; stamens 10–12, held within the corolla, anthers moderate yellow (161A), filaments white, with light purplish pink (62C) tips; pistil exceeding the corolla and 15mm longer than the stamens, style pale yellowish pink (36D), stigma vivid reddish orange (43A). Lvs lanceolate, 150–210 × 60–70mm, margins flat, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 25–30mm long. Shrub 5.0m. May. (Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101282). Etymology: named for a painter of the 19th century from Pollone, Biella, Italy)

(r) **'Luigi Squillario'**

cv. Elepidote rhododendron : parentage unknown

G: Parco Burcina, S: E. Accati & N. Furno (2000),

N: E. Accati & N. Furno (2008), REG: E. Accati &

N. Furno (2008)

Fls 14–18/tight hemispherical truss, funnel-shaped, 40–45 × 50–60mm, with 5 wavy-edged lobes, buds strong reddish purple (N78B), opening deep purplish pink (N78C) throughout, with dark red (187B) spots on the dorsal lobe which just extend onto the adjacent lobes; stamens 10, held within the corolla, anthers brownish orange (N167A), filaments deep purplish pink (N78C); pistil equalling or exceeding the corolla, up to 10mm longer than the stamens, style strong yellowish pink (37A), stigma strong red (46A). Lvs oblanceolate, 120–130 × 40–45mm, margins flat, cuneate at base, broadly acute at apex, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 15–20mm long. Shrub 2.0m. May.

(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101284). Etymology: named for the manager of the mutual savings bank of Biella, who funded most of the work that has been carried out on the rhododendrons in the Parco Burcina)

(r) **'Lynn Drake'**

cv. *Elepidote rhododendron* : *yakushmanum* × *tsariense*

H: A.F. George (1972), G: Hydon Nurseries (2000), N: Mrs A.M. George (2008), I: Hydon Nurseries (2008), REG: Hydon Nurseries (2008)

Fls 8/compact, rounded truss, open campanulate, 43 × 50mm, with 5 wavy-edged lobes, light pink in bud, opening inside blush/cream, outside streaked and flushed pink; slight crimson spotting on the dorsal lobe. Calyx very small, greenish cream. Lvs narrowly elliptic, 80 × 25mm, margins decurved, glossy green above, slightly rounded at base, slightly pointed at apex, with a deep snuff brown indumentum below. Shrub 2.5 × 2.5m in 9 years. April.

(Etymology: named for an artist friend, who provided the Georges with a portrait of one of their springer spaniels)

(a) **'Maggie Brown'**

cv. *Deciduous azalea* : 'Cecile' (s) × 'Leonard Frisbie'
H: F.D. Mossman, G: F.D. Mossman,
N: F.D. Mossman (2008), I: Briggs Nursery (2009),
REG: R.W. Cavender (2008)

Fls 10–15/flat truss, tubular funnel-shaped, 45 × 50–55mm, with 5 wavy-edged lobes; varying from single to semi-double or double – the singles have 5 stamens with anthers, the semi-double and double flowers have 1–5 petaloid stamens that vary from being fully formed lobes to extremely deformed and narrow lobes – all forms occur in the same truss but the overall effect is of double flowers; buds deep pink (52C), opening inside variably coloured: 1–3 dorsal lobes strong orange-yellow (17A), the others light pink to deep pink (52C); if fully formed, 1–3 dorsal petaloid stamens are strong orange-yellow (17A), while the remaining petaloids are light to dark pink; with age the yellow fades and is present only at the base and lower half of the dorsal lobe; the outside of the flower has a deep pink (52C) tube and lobes variably coloured: light to deep pink on the ribs and on all or almost all of the lobes; as much as half of a few petals are very pale pink to almost white; strongly scented. Lvs elliptic, 30–65 × 20–30mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 10 years. Mid May to mid June (with usually 4–5 flowers open at any one time over a 7–15 day period).

(Etymology: Maggie Brown is a granddaughter of Frank Mossman)

(r) **'Mahasona'**

cv. *Elepidote rhododendron* : 'Barbara Reuthe' (s) × 'Zelia Plumecocq'

H: K.-H. Hübbers (1990), G: K.-H. Hübbers (1996), N: Mrs L. Hübbers (2004), REG: J. Westhoff (2008)
Fls up to 9 /slightly hanging, loose truss, funnel-shaped, 80 × 110mm, with 7 wavy-edged lobes, inside with a brilliant yellow (9C) throat, shading out to pale greenish yellow (9D) at margins; outside light yellow (10B/C); two lines of strong yellow-green (144A) spots on the dorsal part of the throat. Calyx with ventral lobes 2mm long, dorsal lobes 6mm long,

brilliant greenish yellow (6C). Lvs oblong, to oblong-elliptic or slightly obovate, 110–130 × 40–55mm, mucronate at apex, slightly rounded at base, margins flat, matt green above, hairless. Shrub 1.5 × 1.5m in 10 years. Early to mid May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101608).
Etymology: the epithet in Hindi means "great gold")

(a) **'Marabella'**

cv. *Deciduous azalea* : 'Washington State Centennial' (s) × ('Raimunde' × 'Twilight Sky')

H: K.-H. Hübbers (1994), G: K.-H. Hübbers (1999), N: K.-H. Hübbers (2006), REG: J. Westhoff (2008)

Fls up to 8/slightly flattened globe truss, broadly funnel-shaped, 50 × 90mm, with 5 very wavy-edged lobes, vivid yellow (13A) in bud, opening brilliant yellow (13C) both inside and out, with a vivid orange-yellow (23A) blotch on the dorsal lobe; slightly scented. Calyx 8mm long, light green. Lvs elliptic, 80 × 40mm, cuneate at apex, with a mucronate tip, cuneate at base, margins flat, matt green above, hairless. Shrub 1.7 × 0.7m in 10 years. Mid May.
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101616).
Etymology: named for a town in Trinidad)

(a) **'Margaret Rowell'**

cv. *Evergreen azalea* : 'Mother's Day' (s) × 'Red Slippers'

H: J.O. Harris (c.1978), G: J.O. Harris (c.1980), N: J.O. Harris (c. 1981), I: J.O. Harris (c.1981), REG: J.O. Harris (2008)

Fls hose-in-hose and semi-double, 2/flat truss, broadly funnel-shaped, c.70 × 75mm, with 11–14 flat-edged lobes, opening strong purplish red (60B) throughout, with a deep red (60A) blotch on the two dorsal lobes. Calyx c.6mm long, light yellow-green (142C). Lvs obovate, c.40 × c.15mm, obtuse at apex, cuneate at base, margins slightly upcurved, semi-glossy and deep yellowish green (141B) above, hairless. Shrub 0.35 × 0.5m in 10 years. Late April.

(Etymology: named for a member of the former Ralph Pennington chapter of the American Azalea Society. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.861)

(r) **'Marie-Louise Agius'**

cv. *Elepidote rhododendron* : yellow form of *eclectum* (s) × *Avalanche* Group

H: E. de Rothschild, G: E. de Rothschild, N: E. de Rothschild (2008), REG: E. de Rothschild (2008)

Fls 14–16/lax truss, ventricose-campanulate, 60–70 × 70–80mm, with 5–6 slightly wavy-edged lobes, strong purplish pink (55B) in bud, opening inside white (155D), slightly flushed cream in the throat, outside white (155B), slightly flushed cream on the tube and with a slight purplish pink flushing down the midveins on some flowers; solid dark red (59A) blotch at the base of the dorsal throat (c.20 × 18mm), breaking up apically into a few spots; stamens 15, filaments white, anthers mottled brown; style pale

yellow-green, glandular, stigma large, capitate, pale yellow-green; ovary densely glandular hairy; sweetly scented. Calyx c.3mm long, pale green with a hyaline margin, flushed purplish along the dorsal rim, glandular hairy. Pedicels 30–40mm long, pale yellowish green, heavily flushed red on the dorsal side. Lvs oblanceolate, 165–250 × 53–75mm, margins flat to decurved, cuneate at base, obtuse and mucronate at apex, matt green above, hairless. Shrub 4.0 × 5.0m in c.25 years. Late March.

(Nomenclatural standard: herbarium specimen provided by the registrant (WSY0112002). Etymology: named for a granddaughter of the registrant)

(r) **'Maris'**

cv. *Elepidote rhododendron* : 'Independence Day' (s) × 'Elegant Touch'

H: J. Barlup (1998), G: J. Barlup (2004), N: J. Barlup (2008), REG: J. Barlup (2008)

Fls 18/ball truss, saucer-shaped, 45 × 75mm, with 5 very wavy-edged lobes, strong purplish red (61B) in bud, opening inside and out very pale purple (69A), blending to slightly darker edges of deep purplish pink (70C), with the outsides having strong reddish purple (70B) midveins; spotted flare of moderate purplish red (184C) extending 30mm from the base. Calyx vestigial. Lvs elliptic, 115 × 50mm, broadly elliptic at apex, rounded at base, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.1m in 10 years. Late May.

(Etymology: named for a friend of the registrant's wife)

(r) **'Marnie's Charm'**

cv. *Elepidote rhododendron* : 'Cornubia' (s) × 'Coronation Day'

H: D.J. Dosser (1995), G: D.J. Dosser (2000), N: D.J. Dosser (2008), I: D. Navdella (2008), REG: D.J. Dosser (2008)

Fls 15–20/ball truss, funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, strong purplish red (58B) in bud and throughout on opening; spotted; stamens very short; slightly scented. Lvs oblong to elliptic, 150 × 50mm, acute at apex, truncate at base, margins flat, matt green above, hairless. Shrub 2.0 × 2.0m in 22 years. October to November (Australia).

(Nomenclatural standard: colour print provided by the registrant (WSY0101953). Etymology: named after the introducers's mother Marnie Navdella)

(a) **'Mary Ann Egan'**

cv. Evergreen azalea : 'Amagasa' (s) × 'Grace Freeman No. 2'

H: J.O. Harris (c.1978), G: J.O. Harris (c.1980), N: J.O. Harris (c.1981), I: J.O. Harris (c.1981), REG: J.O. Harris (2008)

Fls single, 2/domed truss, broadly funnel-shaped, c.65 × 80mm, with 5 wavy-edged lobes, strong yellowish pink (35C) in bud, opening inside and out white (155C), all lobes edged with a solid band of moderate reddish orange (35B); inconspicuous central marking of tiny, strong purplish red (60D) speckles. Calyx c.10mm long, brilliant yellow-green (142B). Lvs elliptic, 40 × c.10mm, broadly acute at apex, cuneate to rounded at base, margins slightly upcurved,

semi-glossy and moderate olive green (147A) above, hairless. Shrub with a dense habit, 0.7 × 0.9m in 10 years. Mid April.

(Etymology: named for a member of the former Ralph Pennington chapter of the Azalea Society of America. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.876)

(r) **'Mary's Choice'**

cv. *Elepidote rhododendron* : *fortunei* (s) × 'Caractacus'

G: H.E. Reiley (1978), N: Mrs M.L. Reiley (2006), REG: Mrs M.L. Reiley (2008)

Fls 10/domed truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, strong purplish red (63B) in bud, opening inside moderate purplish red (65A) at margins, shading to light purplish pink (65B) and then to very pale purple (69C) in the centre, with strong purplish red (63B) stripes on the lobes; outside moderate purplish red (65A); strongly scented. Calyx c.3mm long, brilliant yellow-green (154B). Lvs elliptic, 105–115 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (137A) above, hairless. Shrub 2.4 × 4.2m in 30 years. Early May.

(Grown from seed purchased from F.W. Schumacher in 1974)

(a) **'Mattie Barron'**

cv. Evergreen azalea : 'Caroline Dorman' (s) × 'Grace Freeman No. 2'

H: J.O. Harris (c.1978), G: J.O. Harris (c.1980), N: J.O. Harris (c.1981), I: J.O. Harris (c.1981), REG: J.O. Harris (2008)

Fls 1–2/flat truss, double, broadly funnel-shaped, c.40 × 50mm, with 8 very wavy-edged lobes, brilliant yellow-green (149A) in bud, opening white (155D) throughout; unmarked. Calyx c.15mm long, brilliant yellow-green (142B). Lvs elliptic, 25 × c.10mm, broadly acute at apex, cuneate at base, margins slightly upcurved, semi-glossy and moderate yellowish green (139B) above, hairless. Shrub 0.5 × 0.9m in 10 years. Mid April.

(Etymology: named for a member of the former Ralph Pennington chapter of the Azalea Society of America. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.885 in which the flower description is now known to have been in error in describing the flowers as striped)

(v) **'Maurice Kupsch'**

cv. *Lepidote rhododendron* : parentage unknown, but probably a *konori* hybrid

S: H.R. Malone, G: Emu Valley Rhododendron Garden (2002), N: Emu Valley Rhododendron Garden (2007), I: Emu Valley Rhododendron Garden (2007), REG: Emu Valley Rhododendron Garden (2008)
Fls up to 18/lax truss, tubular funnel-shaped, 100 × 75mm, with 7–8 wavy-edged lobes, pale greenish yellow (2D), with slight pink tinges in bud, opening inside pale purplish pink (56C), fading slightly with age, outside pale purplish pink (56C); unmarked;

strongly clove-scented. Calyx 65mm long, deep red (185A), deeply divided. Conspicuous, large, red bud scales also present. Lvs broadly elliptic, 147 × 93mm, margins upcurved, glossy above, obtuse at base, obtuse to broadly acute at apex, scaly. Shrub 1.0 × 1.0m in 12 years. Early autumn (Tasmania). (Nomenclatural standard: colour print provided by the registrant (WSY0101028). Date of the selection unknown to registrant. Etymology: named for a member of the Emu Valley Rhododendron Group who has done a lot of work in their Garden)

(r) **'Maurizio Zumaglini'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 17–18/tight hemispherical truss, funnel-shaped, 40–45 × 64–70mm, with 5 wavy-edged lobes, buds deep to strong purplish red (61A-B), opening deep to moderate purplish pink (N74C-D), with a lighter centre and with rather few dorsal spots of deep to strong purplish red (61A-B); stamens 10, exerted from the corolla, anthers yellowish white (158C), filaments deep purplish pink (N66D); pistil exerted from the corolla and 3–6mm longer than the stamens, style deep purplish pink (N57D), stigma black. Lvs broadly elliptic, 110–111 × 33–34mm, margins flat, broadly acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (147D) below, hairless; petioles 25–29mm long. Shrub 0.5m. May. (Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101288). Etymology: named for a famous physician and botanist of the 19th century, who wrote 'La Flora Pedemontana')

(a) **'Mayaro'**

cv. Deciduous azalea : 'Washington State Centennial' (s) × ('Raimunde' × 'Twilight Sky')
H: K.-H. Hübbers (1994), G: K.-H. Hübbers (1999),
N: K.-H. Hübbers (2006), REG: J. Westhoff (2008)
Fls up to 10/loose, nearly round truss, funnel-shaped, 60 × 110mm, with 5 very wavy-edged lobes, buds vivid reddish orange (42B), opening inside light to pale orange-yellow (23C/D) and vivid reddish orange (34B), outside strong reddish orange (42C) and pale orange-yellow (23D); with a vivid orange-yellow (23A) blotch on the dorsal lobe; slightly scented. Calyx 3mm long, light green. Lvs elliptic, 90 × 30mm, cuneate at apex with a mucronate tip, cuneate at base, margins flat, matt green above, hairless; young foliage tinged red. Shrub 0.8 × 0.7m in 10 years. Mid May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101618). Etymology: named after a seaside resort town in Trinidad)

(r) **'Meela'**

cv. Elepidote rhododendron : 'Amber Touch' (s) × 'Recital'
H: J. Barlup (1996), G: J. Barlup (2000), N: J. Barlup (2008), REG: J. Barlup (2008)
Fls 17/ball truss, funnel-shaped, 40 × 55mm, with 7 wavy-edged lobes, deep pink (50B) in bud, opening

throughout brilliant yellow (10A), fading to light yellow (10C) on the outer edges; unmarked; slightly scented. Calyx 30mm long, brilliant yellow (10A). Lvs elliptic, 80 × 40mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.1m in 8 years. Late May.

(Etymology: named for the daughter of a neighbour of the registrant – her favourite colour is yellow)

(r) **'Mendocino Magic'**

cv. Elepidote rhododendron : parentage unknown
H: L. Charvet (1995), G: Descanso Nurseries (2000),
N: Descanso Nurseries (2007), I: Descanso Nurseries,
REG: Descanso Nurseries (2008)
Fls 16/domed truss, open funnel-shaped, 55 × 75mm, with 5 wavy-edged lobes, deep pink (52B) in bud, opening inside pale purplish pink (55D), shading to strong purplish pink (55B) at margins and with deep red (53A) and moderate red (47A) spotting in the dorsal throat; outside strong purplish pink (55B). Calyx c.1.5mm long, deep purplish pink (55A). Lvs elliptic, 80 × 50mm, broadly acute at apex, rounded at base, margins downcurved, glossy and moderate olive green (137A) above, hairless. Shrub 0.45 × 0.6m in 6 years. Mid April.

(a) **'Midnight Flare'**

cv. Evergreen azalea : 'Red Red' (s) × 'Red Slippers'
H: J.O. Harris (c.1979), G: J.O. Harris (c.1981),
N: J.O. Harris (c.1982), I: J.O. Harris (c.1982),
REG: J.O. Harris (2008)
Fls single, 2–11/domed truss, funnel-shaped, c.55 × 50mm, with 5 flat-edged lobes, deep red (53A) in bud, opening deep red (53A) throughout, with a strong purplish red (63B) blotch composed of large spots on the three dorsal lobes. Calyx 30mm long, strong red (53B). Lvs elliptic, 40 × 20mm, broadly acute at apex, cuneate at base, margins slightly upcurved, semi-glossy and dark bluish green (131A) above, hairless. Shrub with a squat, spreading habit, 1.2 × 1.2m in 10 years. Mid April.

(Etymology: name inspired by the similarity to a railroad flare. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.906)

(v) **'Miss Essie'**

cv. Lepidote rhododendron : 'Neesa' (s) × *konori*
H: G.L.S. Snell (2003), G: K. Hagan (2007),
N: K. Hagan (2008), REG: K. Hagan (2008)
Fls 10/lax truss, tubular funnel-shaped, 80 × 100mm, with 7 wavy-edged lobes, inside light yellow (10B), lightly marked light orange (29B) in the throat, outside brilliant yellow (10A) on the tube, light yellow (10C) on the lobes; unsported; slightly scented. Lvs ovate, 140 × 60mm, acute at apex, cuneate at base, margins flat, glossy green above. Shrub 1.0 × 0.5m in 5 years. Spring to summer. (Nomenclatural standard: colour print provided by the registrant (WSY0101962). Etymology: named for Miss Essie Huxley a long time member of the Australian Rhododendron Society who died in 2008 aged 90)

(a) **'Miss Susie McGhee'**

cv. Evergreen azalea : 'Hershey's Orange' (s) × 'Hershey's Red'

H: J.O. Harris (c.1970), G: J.O. Harris (c.1972),
N: J.O. Harris (c.1972), I: J.O. Harris (c.1972),
REG: J.O. Harris (2008)

Fls hose-in-hose, 2–3/domed truss, funnel-shaped,
c.55 × 40–50mm, with 5+5 wavy-edged lobes,
opening vivid red (57A) throughout, unmarked.
Calyx c.10mm long, strong yellow-green (142A). Lvs
elliptic, 25 × c.6mm, broadly acute at apex, cuneate at
base, margins slightly upcurved, semi-glossy and dark
green (135A) above, hairless. Shrub with a dense
habit, 0.6 × 0.9m in 10 years. Early April.

Syn: 'Miss Suzie'

(Etymology: named for the individual who founded
the first garden club in Atlanta, Georgia, USA.

Revised entry: previously listed as 'Miss Suzie'
(unregistered) in *The International Rhododendron
Register and Checklist*, 2004, p.920)

(a) **'Miss Suzie'** Syn. of 'Miss Susie McGhee'

(r) **'Monbarone'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)

Fls 14/loose, hemispherical truss, funnel-shaped,
35–40 × 50–55mm, with 5 flat or slightly wavy-edged
lobes, buds strong purple (77B), opening very light
purple (76B) at the edges and with a white centre;
conspicuous spotting of strong orange-yellow (163B)
on the dorsal lobe and just extending onto the adjacent
lobes; stamens 10, exserted from the corolla, anthers
white, filaments white; pistil exserted from the corolla
and 10mm longer than the stamens, style strong
purplish pink (73B), stigma dark purple (71A). Lvs
elliptic, 100–145 × 40–50mm, margins flat, broadly
acute at base and apex, matt moderate olive green
(147A) above, moderate yellow-green (146C) below,
hairless; petioles 15–25mm long. Shrub 2.0m. May.
(Nomenclatural standard: colour print of an
electronic image provided by the registrants
(WSY0101287). Etymology: named for a mountain
that overtops the Parco Burcina)

(r) **'Mondnacht'**

cv. Elepidote rhododendron : 'Grünlich' × 'Peucine'
H: K.-H. Hübbers (1993), G: K.-H. Hübbers (1999),
N: K.-H. Hübbers (2007), REG: J. Westoff (2008)

Fls up to 12/flat-topped ball truss, open funnel-
shaped, 55 × 100mm, with 6–7 flat-edged lobes, light
orange-yellow (24C) in bud, opening inside light
greenish yellow (4B/C), outside light to pale greenish
yellow (1C/D); moderate brown/brownish orange
(165A/B) spots towards the base of the dorsal part of
the throat; stigma large, greenish. Calyx 3–10mm
long, strong yellow-green (144A–C). Lvs oblong,
75–120 × 35–70mm, mucronate at apex, truncate at
base, margins slightly wavy-edged, glossy green above,
hairless. Shrub 1.4 × 0.9m in 10 years. Mid May to
early June.

(Nomenclatural standard: colour print of an

electronic image provided by the registrant
(WSY0101626). Etymology: the epithet is the
German for "moon night" in allusion to the intense
fluorescent appearance of the flowers at dusk)

(r) **'Monte Mucrone'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)

Fls 15–16/tight hemispherical truss, funnel-shaped,
40–45 × 60–70mm, with 5 wavy-edged lobes, buds
vivid purplish red (61C), opening deep purplish pink
(68A) throughout, with moderate purplish red and
strong purplish red (59C, 60B) spots on all the lobes
(densest and darkest on the dorsal lobes); stamens 10,
held within the corolla; anthers greyish purplish red
(N77D), filaments white; pistil held within the
corolla, 2–3mm longer than the stamens, style white,
stigma dark red (59A). Lvs lanceolate, 115–120 ×
45–50mm, margins flat, acute at base and apex, matt
moderate olive green (147A–146A) above, moderate
yellow-green (146C) below, hairless; petioles 15–20mm
long. Shrub 2.0m. May.

(Nomenclatural standard: colour print of an
electronic image provided by the registrants
(WSY0101289). Etymology: named for a mountain
that overtops the Parco Burcina)

(r) **'Monviso'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)

Fls 18–22/loose, conical truss, funnel-shaped, 35–40
× 50–60mm, with 5 flat to very slightly wavy-edged
lobes, buds strong purplish red (72B) at apex, vivid
reddish purple (N74A) at base, opening deep purplish
pink (72D) throughout, with strong greenish yellow
(151B) spots over 3–4 lobes but principally on the
dorsal lobe; stamens 10, held within the corolla,
anthers strong reddish purple (72B), filaments
moderate purplish pink (N74D); pistil exserted from
the corolla, 10mm longer than the stamens, style
strong purplish red (58C), stigma deep purplish red
(59B). Lvs elliptic, 135–145 × 40–45mm, surface
slightly convex, margins flat, broadly acute at base and
apex, matt moderate olive green (146A) above,
moderate yellow-green (146D) below, hairless;
petioles 15mm long. Shrub 4.0m. May.

(Nomenclatural standard: colour print of an
electronic image provided by the registrants
(WSY0101291). Etymology: named for a mountain
of the Alps that is visible from the Parco Burcina)

(r) **'Morioka-bijin'**

cv. Elepidote rhododendron : Pilgrim Group (s) ×
(*hyperythrum* × 'The Honourable Jean Marie de
Montague')

H: K. Terai (1979), G: T. Saitō (1990), N: T. Saitō
(2006), REG: T. Saitō (2008), S: T. Saitō (1992)

Fls 6–10/truss, hemispherical truss, funnel-shaped,
40 × 50mm, with 5 slightly wavy-edged lobes, buds
vivid red (45B), opening vivid red (45B) throughout,

unmarked. Calyx 1mm long, red. Lvs lanceolate, 70–100 × 22–33mm, curved at base, acute at apex, hairless. Shrub 0.6 × 0.8m in 28 years (pot grown). Late April.

(Nomenclatural standard: colour slide provided by the registrant (WSY0101455). Etymology: named for the district in which it was raised (Morioka) and “bijin” meaning beautiful woman)

(a) ‘Nancy Lynn’

cv. Evergreen azalea : ‘Mrs Nancy Dippel’ (s) × ‘Florence Waldman’

H: J.F. Klimavicz (1997), G: J.F. Klimavicz (1999), N: J.F. Klimavicz (2007), REG: J.F. Klimavicz (2008) Fls 2/ball truss, saucer-shaped, hose-in-hose and fully double, with no pistil present, 30 × 55mm, with 19 very wavy-edged lobes, buds white (155D) with sectors and flecks of deep purplish pink (N74C), opening inside white (155D), with deep purplish pink (N74C) sectors and flecks, slight spotting of light yellow-green (145C-D) and a pale greenish yellow (1D) throat; outside similarly coloured. No normal calyx present. Lvs elliptic, 45 × 20mm, broadly acute at apex, cuneate at base, margins flat, glossy and strong yellow-green (144A) above, hairless. Shrub with a dense habit, 0.45 × 0.45m in 5 years. Mid May. (Etymology: named for a member of the registrant’s family)

(r) ‘Nicoletta Furno’

cv. Elepidote rhododendron : parentage unknown G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati (2008), REG: E. Accati (2008) Fls 14–18/hemispherical truss, funnel-shaped, 30–35 × 60–65mm, with 5 very wavy-edged lobes, buds strong purplish red (72A), opening light purple (75A), with a lighter centre and with only the dorsal lobe with deep red (53A) spots; stamens 10, exserted from the corolla, anthers pale yellow (11C), filaments very pale purple (76D); pistil exserted from the corolla and the same length as the stamens, style very pale purple (69D), stigma dark purple (79B). Lvs lanceolate, 120–135 × 35–40mm, margins flat, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (147C) below, hairless; petioles 20–25mm long. Shrub 1.0m. May. (Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101293). Etymology: named for the manager of the Natural Reserve of the Parco Burcina)

(r) ‘Nishan’

cv. Elepidote rhododendron : ‘Mondnacht’ (s) × ‘Hachmann’s Kabarett’

H: K.-H. Hübbers (2001), G: K.-H. Hübbers (2006), N: Mrs L. Hübbers (2007), REG: J. Westhoff (2008) Fls up to 20/broad, roundish truss, campanulate/funnel-shaped, 55 × 70mm, with 6 wavy-edged lobes, inside white (155C), outside white (155B), with dark red/deep purplish red (59A/B) spots covering three-quarters of the dorsal throat; stigma greenish. Calyx 2–10mm long, moderate yellow (160A). Lvs oblong-obovate, 80–120 × 50–80mm, mucronate at apex, cuneate at base, margins slightly wavy, glossy

green above, hairless. Shrub 0.6 × 0.7m in 5 years.

Early to mid May.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101622).

Etymology: the epithet in Hindi means “blotch”)

(r) ‘Nordic Smile’

cv. Elepidote rhododendron : *wardii* (s) × (*wightii* × *aureum*)

H: C.A. Lehmann (1985), G: P. Schaarup (1992),

S: P. Schaarup (1992), N: C.A. Lehmann (2007),

I: Hvidbjerg Planteskole (2007), REG: C.A. Lehmann (2008)

Fls 12–14/domed truss, funnel-shaped, 45 × 75mm, with 6 wavy-edged lobes, buds moderate red (N34A), opening light greenish yellow (4B) both inside and out, with pink tinges on the lobes; numerous, moderate red (N34A) spots on the basal two-thirds of the dorsal lobes; stamens 10, filaments 30mm long, white, anthers white; ovary 30mm long; style 30mm long, light green. Calyx absent to 1mm long, very light green. Lvs 50–80 × 30–50mm, cordate at base, obtuse at apex, hairless. Shrub 1.75 × 1.0m in 20 years. Mid April to mid May

(Nomenclatural standard: colour print derived from an electronic image provided by the registrant (WSY0101964). Etymology: named for its geographical origin and for the plant ‘giving a kind of smile’ when it opens its buds)

(r) ‘Nordic Venture’

cv. Elepidote rhododendron : ‘Yaku Sunrise’ (s) × *roxieanum* var. *roxieanum* Oreonastes Group

H: J. Rosenthal (1979), G: S.A. Askjaer (1986),

N: C.A. Lehmann (2008), I: Hvidbjerg Planteskole (2008), REG: C.A. Lehmann (2008)

Fls 21–22/tight domed truss, broadly funnel-shaped, 33 × 47mm, with 5 smooth or slightly wavy-edged, rather broadly rounded and hardly emarginate lobes, buds strong purplish red (58C), opening inside pale purplish pink (56C), outside pale purplish pink (56C), with stripes of strong purplish red (58C); numerous spots on the dorsal lobe of moderate purplish red (65A). Calyx 33mm long, pale purplish pink (56C). Stamens 10, filaments 15–25mm long, white, hairy below, anthers light pink; style pinkish, stigma moderate purplish pink (65A). Lvs narrowly elliptic to narrowly obovate, 50–110 × 11–20mm, margins decurved, narrowly cuneate at base, narrowly acute at apex, with light orange-yellow (25D) hairs below when young, aging to moderate orange (166D). Shrub 1.6 × 1.0m in 20 years. Late April to Mid May. (Nomenclatural standard: colour print of the digital picture supplied by the registrant (WSY0101969). Etymology: named for its geographical origin and to commemorate the collaborative venture to produce it; the cross was made at the registrant’s suggestion with pollen supplied by him)

(r) ‘North Wind’

cv. Elepidote rhododendron : ‘Scintillation’ (s) × ‘Snow Candle’

H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2007), REG: J. Barlup (2008)

Fls 15/ball truss, broadly funnel-shaped, 65 × 90mm, with 5 wavy-edged lobes, light purple (75A) in bud, opening inside very pale purple (69D), with strong reddish purple (70B) veins extending 25mm from the centre on all lobes, and with a tinge of very pale purple (69A) extending along the midveins for 40mm; outside very pale purple (69D), with strong reddish purple (70B) midveins; prominent, deep red (185A), dorsal flare, c.15mm wide and solid for lower 25mm, with concentrated spotting extending another 20mm. Calyx minute. Lvs elliptic, 150 × 65mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.5m in 10 years. Late April.

(r) **'Orange 2002'** Syn. of 'Twilight Fire'

(r) **'Orchid Bouquet'**

cv. *Elepidote rhododendron* : 'Purple Splendour' (s) × 'One Thousand Butterflies'
H: F. Fujioka (1987), G: F. Fujioka (1993),
N: G.H. Kesterson (2008), I: Meerkerk Gardens (2008), REG: F. Fujioka (2008)
Fls 19/ball truss, broadly funnel-shaped, 55 × 100mm, with 5–6 wavy-edged lobes, buds strong purplish red (67A), edged and tipped deep reddish purple (77A), opening inside strong purplish pink (73B), with a c.15mm wide margin and radial lines of strong reddish purple (72B) and distinct deep purplish red (61A) nectaries; outside strong purplish red (71C), with a c.15mm wide margin and radial lines of strong reddish purple (72C); dorsal blotch of light orange-yellow (16C), with brownish orange (165B) spotting. Calyx c.3mm long, moderate olive green (137B). Lvs elliptic, 200 × 65mm, broadly acute at apex, cuneate at base, margins flat, dull and dark bluish green (131A) above, hairless. Shrub 2.1 × 1.2m in 10 years. Early May.

(r) **'Oremo'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 7–10/hemispherical truss, funnel-shaped, 35–40 × 50–55mm, with 5 wavy-edged lobes, buds strong purplish red (67A), opening vivid purplish red to deep purplish pink (67B–C) throughout, with strong greenish yellow (151A/N144A) spots chiefly on the dorsal lobe and on a paler ground apically; stamens 10, exerted from the corolla, anthers moderate purple (79D), filaments deep purplish pink (N57C–D); pistil 10mm longer than the stamens, style deep purplish pink (58D), stigma strong greenish yellow (151B). Lvs usually ovate, 100–110 × 40mm, margins flat, acute at apex, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 15mm long. Shrub 2.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101297). Etymology: named for a stream that runs beside the Parco Burcina, where people used to find gold)

(r) **'Oropa'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 17–18/conical truss, funnel-shaped, 30–35 × 55mm, with 5 flat or slightly wavy-edged lobes, buds strong purplish red/deep purplish pink (64C/N66D) in bud, with a white base, opening white throughout and with discrete spotting of strong greenish yellow (153B–C) on the dorsal lobe alone; stamens 10, held within the corolla; anthers white, filaments white; pistil equalling or exerted from the corolla, exceeding the stamens by 0–10mm, style white, stigma strong purplish red (60B). Lvs lanceolate, 125–140 × 40–45mm, margins flat, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (146D) below, hairless; petioles 20mm long. Shrub 3.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101302). Etymology: named for the location of the Oropa sanctuary, near the Parco Burcina and part of the UNESCO World Heritage site)

(r) **'Our Kate Marise'**

cv. *Elepidote rhododendron* : *macabeanum* × *calophytum*
H: È. de Rothschild, G: E. de Rothschild, N: Lionel de Rothschild, REG: E. de Rothschild (2008)
Fls 18–21/open-topped truss, ventricose-campanulate, 65–70 × c.60–75mm, with 7 slightly wavy-edged lobes, moderate purplish pink (62B) in bud, inside opening white, outside white, very slightly suffused pale cream; blotch of dark red (59A to 187A) at the base of the throat composed of rays on each lobe, rays longest and broadest on the dorsal side; slightly scented; stamens 17, held within the corolla, filaments white, anthers brown; style cream, stigma large, capitate, pale yellow; ovary densely white hairy. Calyx 1mm long, pale yellowish green. Pedicels 40–45mm long, pale yellowish green, flushed with red on the dorsal side. Lvs oblanceolate, 140–230 × 35–70mm, margins flat or very slightly decurved, cuneate at base, acute to obtuse and mucronate at apex, with a single thin layer of dense, cobwebby, very pale fawn hairs (pale yellow, 161C). Shrub 3.0 × 4.0m in c.25 years. March.
(Nomenclatural standard: herbarium specimen provided by the registrant (WSY0112004))

(a) **'Parfait'** Syn. of 'Harris Parfait'

(r) **'Parco Burcina'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 13–16/loose, hemispherical truss, funnel-shaped, 40 × 65mm, with 5 wavy-edged lobes, buds light purple (75B), opening very pale purple (75D) at the margins and with a white centre, with relatively sparse dorsal spotting of strong yellow (N144B), just extending on to the adjacent lobes; stamens 10, held within the corolla, anthers very pale purple (76D),

filaments white; pistil exerted from the corolla, 5–10mm longer than the stamens, style light greenish yellow (4C), stigma dark red (59A). Lvs elliptic, 110 × 45–50mm, margins flat, broadly acute to obtuse and mucronate at apex, matt moderate olive green (147A) above, moderate yellow-green (146D) below; petioles 15–20mm long. Shrub 4.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101301).
Etymology: name for the place where it was selected)

(r) **'Pecos Dawn'**

cv. *Elepidote rhododendron* : 'Amber Touch' (s) × 'Rocky Point'
H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2007), REG: J. Barlup (2008)
Fls 17/ball truss, broadly funnel-shaped, 55 × 90mm, with 7 wavy-edged lobes, opening inside pale greenish yellow (13D), with a deeper shade of light yellow (14D) on the dorsal lobes, tinged strong pink (48C) on the tips, with pale orange-yellow (165D) veining on all lobes, extending from base to tip, and with strong red (53C) nectar pouches extending a quarter of the distance around the base; outside similar but without the 48C tingeing at the tips; heavy strong red (53B) spotting on the three dorsal lobes extending 50mm from base to tip. Calyx 75mm long, pale greenish yellow (13D), with strong red (53B) spotting on the entire calyx. Lvs oblong, 105 × 40mm, broadly acute at apex, rounded at base, margins flat, glossy and moderate olive green (147A) above, hairless. Shrub 1.1 × 1.5m in 12 years. Mid May.
(Etymology: name inspired by the colours in the sky at dawn when seen from a bluff above the Pecos River (New Mexico, USA))

(a) **'Phyllis Jean'**

cv. *Evergreen azalea* : ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'
H: W.L. Clagett (1996), G: W.L. Clagett (2001), N: W. Meyers (2008), I: White's Nursery (2009), REG: W.L. Clagett (2008)
Fls hose-in-hose, 3/flat truss, open funnel-shaped, 32 × 38mm, with 10 wavy-edged lobes, strong red (50A) and white in bud, opening inside strong red (50A) at the margins and shading to almost white at the centre of the lobes, the throat pale yellowish green; outside similar but shading to greenish yellow at the base; large reddish spots on the dorsal lobes. Lvs elliptic, 40 × 20mm, broadly acute at apex, cuneate at base, with flat margins, glossy and moderate olive green (147A) above, with fine white hairs on the lower surface when young. Shrub 1.0 × 0.9m in 11 years. Mid May.
(Etymology: Named for Phyllis Jean Meyers, wife of the William Meyers who was the highest bidder in a fund-raising auction to win the right to name this seedling)

(r) **'Pier Giorgio Frassati'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 20–25/hemispherical truss, funnel-shaped, 32–35

× 45–55mm, with 5 slightly wavy-edged lobes, buds moderate purplish red/strong reddish purple (70A-B), opening with light reddish purple (74C) margins, the centre light and very pale purple (77D, 76C), with dark reddish purple, dorsal spotting; stamens 10, held within the corolla, anthers pale yellowish pink (29D), filaments very pale purple (76C); pistil exerted from the corolla, 10mm longer than the stamens, style strong purplish pink (63C), stigma vivid reddish orange (34A). Lvs broadly lanceolate, 97–119 × 44–57mm, margins flat, acute to obtuse and mucronate at apex, matt moderate olive green (147A) above, moderate yellow-green (146D) below, hairless; petioles 12–15mm long. Shrub 1.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101306).
Etymology: named for a saint from Biella, the province in which the Parco Burcina is situated)

(r) **'Pietro Porcinai'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 21–22/hemispherical truss, funnel-shaped, 38–42 × 54–60mm, with 5 slightly wavy-edged lobes, buds strong reddish purple (N78B), opening moderate purplish pink (N74D), with a very light purple (75C) centre and with deep orange-yellow (163A) and deep greenish yellow (153A) spots on the dorsal lobes (on a paler ground apically); stamens 10, held inside the corolla, anthers moderate purplish pink (186D), filaments very pale purple (73D); pistil exerted from the corolla and 8–10mm longer than the stamens, style strong purplish pink (67D), stigma moderate red (180C). Lvs lanceolate, 116–137 × 41–45mm, slightly twisted, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (146B) below, hairless; petioles 17–20mm long. Shrub 5.0m. June.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101308).
Etymology: named for a 20th century landscape architect practising in Biella and who provided the layout of the car park at the Parco Burcina)

(r) **'Pink Palette'**

cv. *Elepidote rhododendron* : 'Anna' (s) × 'Ken Janek'
H: R.A. Furman (1979), G: R.A. Furman (1985), N: R.A. Furman (2006), REG: R.A. Furman (2008)
Fls 15/domed truss, funnel-shaped, 50 × 75mm, with 5 very wavy-edged lobes, moderate red (47A) in bud, opening inside deep pink (52C) at margins, shading to white (155D) towards the centre, with strong red (46A) dorsal spotting; outside deep purplish pink (54B); long-lasting and of good substance. Lvs elliptic, 100–150 × 40–55mm, obtuse at apex, cuneate at base, margins flat, semi-glossy and dark green above, with a white, hairy indumentum below on mature lvs. Shrub with a dense habit, 2.7 × 2.7m in 25 years, the foliage resistant to yellow leaf spotting. Midseason.

(r) **'Pitacca'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)
Fls 13–18/hemispherical truss, funnel-shaped, 35 ×
60mm, with 6 flat-edged lobes, light purple (76A) in
bud, opening very pale purple (76C) throughout,
with a conspicuous blotch and spotting of dark red
(187B) occupying most of the dorsal lobe and
spreading onto the adjacent lobes; stamens 10,
exserted from the corolla, anthers pale greenish
yellow (1D), filaments very pale mauve (76C); pistil
equalling or exceeding the corolla, the same length as
the stamens, style white, stigma vivid reddish orange
(30A). Lvs lanceolate, 105–110 × 35–40mm, margins
flat, acute at base and apex, matt moderate olive green
(147A) above, moderate yellow-green (146C) below,
hairless; petioles 15mm long. Shrub 2.0m. May.
(Nomenclatural standard: colour print of an
electronic image provided by the registrants
(WSY0101310). Etymology: named after a site in the
Parco Burcina)

(r) **'Plum Wild'**

cv. Elepidote rhododendron : 'Jonathan Shaw' (s) ×
'Plum High'
H: J. Barlup (2000), G: J. Barlup (2004), N: J. Barlup
(2006), REG: J. Barlup (2008)
Fls 19/ball truss, broadly funnel-shaped, 40 × 55mm,
with 5 wavy-edged lobes, dark red (59A) in bud,
opening inside deep purplish red (61A), with a
slightly darker edging of dark red (59A) and a dorsal
flare of dark red (187A) spots extending c.25mm from
the base; outside deep purplish red (61A). Calyx
vestigial. Lvs elliptic, 125 × c.50mm, broadly acute at
apex, rounded at base, margins flat, dull and moderate
olive green (147A) above, hairless. Shrub 1.1 × 1.1m
in 8 years. Late May.

(r) **'Pollone'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)
Fls 16/hemispherical truss, funnel-shaped, 40–45 ×
50–55mm, with 5 wavy-edged lobes, buds deep
purplish pink (68B), opening white, with strong
purplish pink (68B) margins and relatively sparse
spotting of strong yellow-green (144A) on the dorsal
lobe and just extending to the adjacent lobes; stamens
10, held within the corolla, anthers pale yellow
(161D), filaments white; pistil exserted from the
corolla and 15–20mm longer than the stamens, style
white, stigma vivid purplish red (N57A). Lvs
lanceolate, 110–120 × 35–40mm, margins flat, acute
at base and apex, matt moderate olive green (147A-
146A) above, moderate yellow-green (146D) below,
hairless; petioles 20mm long. Shrub 2.0m. May.
(Nomenclatural standard: colour print of an
electronic image provided by the registrants
(WSY0101312). Etymology: named for the town and
municipality where the Parco Burcina is situated)

(r) **'Polynesian Girl'**

cv. Elepidote rhododendron : 'Sweet Mystery' (s) ×
'Polynesian Sunset'
H: C. Weeks (1995), G: C. Weeks (2001),
N: C. Weeks (2008), I: Westgate Gardens Nursery
(2008), REG: C. Weeks (2008)
Fls 9–10/domed truss, funnel-campanulate, 50 ×
55mm, with 5 wavy-edged lobes, strong purplish red
(58B) in bud, opening inside strong purplish red
(63B) at margins, shading paler to brilliant yellow
(13B) at the centre; outside similarly coloured; a few,
vivid yellow (12A), dorsal spots. Lvs elliptic, 115 ×
30mm, acute at apex, cuneate at base, margins flat,
dull and moderate olive green (137A) above, hairless.
Shrub 0.45 × 0.5m in 10 years. Late May.

(a) **'Pride of Lawrenceville'**

cv. Evergreen azalea : 'Bunka' (s) × 'Hinodogiri'
H: J.O. Harris (c.1970), G: J.O. Harris (c.1972),
N: J.O. Harris (c.1972), I: J.O. Harris (c.1972),
REG: J.O. Harris (2008)
Fls single, up to 16/domed truss, open funnel-shaped,
c.55 × 70mm, with 5 wavy-edged lobes, strong
purplish red (63A) in bud, opening inside and out
pale purplish pink (62D), edged with a continuous
band of strong purplish red (63B) and with a diffuse
dorsal blotch of tiny, strong purplish red (63B)
speckles. Calyx c.6mm long, strong yellow-green
(142A). Lvs elliptic, 45 × c.20mm, broadly acute at
apex, cuneate at base, margins slightly upcurved,
semi-glossy and strong yellow-green (143A) above,
hairless. Shrub with a dense habit, 1.1 × 1.5m in 10
years. Late April.
(Etymology: named for the home town of the
registrant. Revised entry: previously listed
(unregistered) in *The International Rhododendron
Register and Checklist*, 2004, p.1099)

(r) **'Provincia di Biella'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)
Fls 15–28/hemispherical truss, funnel-shaped, 45 ×
70mm, with 5 slightly wavy-edged lobes, buds strong
to light purple (77B-D), opening strong to light purple
(77B,C,D), shading to white; with spots of strong
orange-yellow (163B) to strong brown (172A); stamens
10, held within the corolla, anthers light purple (76A),
filaments very light/very pale purple (75C-D); pistil
exserted from the corolla, 5mm longer than the
stamens, style pale purplish pink (62D) to white,
stigma moderate yellowish pink (179D). Lvs lanceolate,
145–200 × 50–60mm, margins slightly wavy, acute
at base and apex, matt moderate olive green (147A)
above, moderate yellow-green (146C) below, hairless;
petioles 12–17mm long. Shrub 1.0m. June.
(Nomenclatural standard: colour print of an electronic
image provided by the registrants (WSY0101314).
Etymology: named for the province in Piedmont,
Italy in which the Parco Burcina is situated)

(r) **'Pyari'**

cv. Elepidote rhododendron : 'Ovation' (s) × 'Rubicon'
H: K.-H. Hübbers (1993), G: K.-H. Hübbers (1998),
N: Mrs L. Hübbers (2006), I: Baumschulen
Wüstemeier (2006), REG: J. Westhoff (2008)
Fls 4–7/lax ball truss, campanulate, 25 × 40mm, with
5 slightly recurved lobes, buds strong red (46A),
opening inside strong red (53B), outside deep red
(53A); unmarked; style cream, nearly twice as long as
the corolla. Calyx 20mm long, deep red (53A). Lvs
elliptic, 60–110 × 25–55mm, mucronate at apex,
cuneate at base, margins flat, glossy green above,
indumentum silver at first, aging to brown. Shrub 0.6
× 0.6m in 10 years. Early to mid May.
(Nomenclatural standard: colour print of an
electronic image provided by the registrant
(WSY0101625). Etymology: the epithet is Hindi for
lovely/love)

(a) **'Rain Fire'**

cv. Evergreen azalea : 'Moonbeam' (s) × 'Lilacinum'
H: J.O. Harris (c.1970), G: J.O. Harris (c.1972),
N: J.O. Harris (c.1972), I: J.O. Harris (c.1972),
REG: J.O. Harris (2008)
Fls single and solitary, broadly funnel-shaped, 45 ×
65mm, with 5 flat-edged lobes, vivid red (45B) in bud,
opening vivid red (45B) throughout; dorsal blotch of
strong purplish red (60B) spots. Calyx c.6mm long,
strong yellowish green (134B). Lvs elliptic, 40 ×
c.15mm, broadly acute at apex, cuneate at base, margins
flat, glossy and deep yellowish green (141B) above,
hairless. Shrub 0.6 × 0.75m in 8 years. Early May.
(Revised entry: previously listed (unregistered) in *The
International Rhododendron Register and Checklist*,
2004, p.1133)

(r) **'Ramella'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)
Fls 16–20/tight, flattened hemispherical truss,
funnel-shaped, 35–45 × 50–60mm, with 5 flat to
slightly wavy-edged lobes, buds vivid purplish red
(N57A), opening vivid purplish red to deep purplish
pink (N57B-C), with a conspicuous blotch of dark to
deep red (59A, 60B) spots on the three dorsal lobes;
stamens 9–10, held within the corolla, anthers
brilliant yellow (13B), filaments moderate to light
purplish pink (65A-B); pistil held within the corolla,
5–10mm longer than the stamens, style strong pink
(48D), stigma vivid reddish orange (34A). Lvs
oblongate, 110–130 × 45–50mm, margins flat,
acute at base, obtuse and mucronate at apex, matt
moderate olive-green (147A–146A) above, moderate
yellow-green (146A) below, hairless; petioles 20mm
long. Shrub 1.0m. May.
(Nomenclatural standard: colour print of an
electronic image provided by the registrants
(WSY0101316). Etymology: a typical family name
found in the Italian province of Biella where the
Parco Burcina is situated)

(a) **'Rivermist'**

cv. Evergreen azalea : 'Sherwoodii' (s) × 'Fedora'
H: J.O. Harris (c.1970), G: J.O. Harris (c.1972),
N: J.O. Harris (c.1972), I: J.O. Harris (c.1973),
REG: J.O. Harris (2008)
Fls hose-in-hose, 3–4/domed truss, broadly
funnel-shaped, c.30 × 40mm, with 5 flat-edged lobes,
light purple (75B) in bud, opening inside and out to a
blend of very light purple (75C) and very pale purple
(75D), with the paler tones more concentrated along
the centres of the lobes, producing a subtle bicolor
effect; tiny, deep purplish pink speckles in a blotch at
the base of the dorsal lobe. Calyx c.20mm long,
moderate yellow-green (138C). Lvs elliptic, 30 ×
c.15mm, broadly acute at apex, cuneate at base,
margins flat, semi-glossy and dark green (131B) above,
hairless. Shrub with an open habit, 1.1 × 0.9m in 8
years. Mid April.

(Revised entry: previously listed (unregistered) in *The
International Rhododendron Register and Checklist*,
2004, p.1165. Can occasionally produce a terminal,
ball-shaped, multiple truss with up to 30 fls)

(r) **'San Barnaba'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)
Fls 12–18/conical truss, funnel-shaped, 35–36 ×
45–55mm, with 5 slightly wavy-edged lobes, buds
strong purplish red (61B), opening deep purplish pink
(N66D), with deep to vivid purplish red (61A,B,C)
spots on the dorsal lobe, spreading onto the adjacent
lobes; stamens 10, held within the corolla, anthers
pale yellow (161D), filaments pale purplish pink
(65D); pistil exerted from the corolla, 12–14mm
longer than the stamens, style light pink (49C),
stigma strong red (46A). Lvs broadly elliptic,
109–126 × 43–50mm, margins flat, narrowly obtuse
at base, narrowly obtuse and mucronate at apex, matt
moderate olive green (147A) above, moderate
yellow-green (147C) below, hairless; petioles
21–27mm long. Shrub 3.0m. May.
(Nomenclatural standard: colour print of an electronic
image provided by the registrants (WSY0101334).
Etymology: named for an ancient church in Pollone,
near where the Parco Burcina is situated)

(r) **'San Pool'**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati &
N. Furno (2008)
Fls 18–20/hemispherical truss, funnel-shaped, 33–39
× 53–63mm, with 5 very wavy-edged lobes, buds
strong purplish red (64B), opening strong purplish
red (71D), with a strong purplish pink (73B) centre
and a very few, inconspicuous, strong purplish red
(61B) spots on the dorsal lobe alone; stamens 10, held
within the corolla, anthers moderate orange (170D),
filaments pale purplish pink (65B); pistil not or just
exserted from the corolla, 7–10mm longer than the
stamens, style light purplish pink (68D), stigma
moderate red (185B). Lvs lanceolate, 115–117 ×

42–44mm, margins flat, with deflexed apices, acute at base and apex, matt moderate yellow-green (147B) above, moderate yellow-green (147C) below, hairless; petioles 10–13mm long. Shrub 1.0m. May. (Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101336). Etymology: named after a location inside the Parco Burcina)

(r) **‘Scheherezade’**

cv. Elepidote rhododendron : *irroratum* ‘Polka Dot’ × *calophytum*
R: E. de Rothschild (pre 1988), N: E. de Rothschild (pre 1988), REG: E. de Rothschild (2008)
Fls up to 18/truss, broadly funnel-shaped, up to 55 × 65mm, with 6 lobes, creamy white (155D), with speckling grading from dark red to moderate purplish red (59A–C) and most dense on the three dorsal lobes, also with a deep purplish red (59B) blotch at the base of the dorsal lobe, extending onto the adjacent lobes; stamens 13, filaments white, anthers fawn (between moderate reddish brown and brownish orange, 166B–C); style pale greenish yellow (1D); ovary green, with a slight red flush. Calyx c.2mm long, pale green-yellow. Lvs oblong-lanceolate, 214 × 53mm, cuneate at base, acute at apex, matt and closest to moderate olive green (147A) above but lighter, finely hairy below (at least on the main vein). Shrub 3.0m or more × 3.0–4.0m in c.20 years. March.
PC 2008

(Nimrod Group. Previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.1230)

(a) **‘Seven Dwarfs’**

cv. Evergreen azalea : ‘Okina-nishiki’ (s) × ‘Grace Freeman No. 2’
H: J.O. Harris (c.1970), G: J.O. Harris (c.1972), N: J.O. Harris (c.1972), I: J.O. Harris (c.1972), REG: J.O. Harris (2008)
Fls single, solitary, broadly funnel-shaped, c.50 × 65mm, with 5 wavy-edged lobes, light purplish pink (62C) in bud, opening inside pale purplish pink (62D), with random markings of strong purplish red (58C) sectors, stripes, and spots of various sizes and shapes, and with a spotted, strong purplish red (63A) blotch on the three dorsal lobes; outside similar, but with silvery overtones. Calyx c.5mm long, brilliant yellow-green (142B). Lvs elliptic, 35 × 15mm, broadly acute at apex, cuneate at base, margins slightly downcurved, semi-glossy and strong yellow-green (144A) above, hairless; some variegated with yellow blotches. Shrub with a dense habit, 0.6 × 1.5m in 9 years. Early May.
(Etymology: named for the several different flower colour patterns exhibited by this plant. Revised entry: previously listed (unregistered) in *The International Rhododendron Register and Checklist*, 2004, p.1244)

(r) **‘S.I.R.’**

cv. Elepidote rhododendron : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000), N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 15–18/tight, conical truss, 40 × 45–50mm, with 5 flat to slightly wavy-edged lobes, buds strong purple (N81B), opening strong purple (N80B), with inconspicuous, small, strong orange-yellow (17A) spots on a paler ground on the dorsal lobe only; stamens 10, held within the corolla, anthers pale yellow (164D), filaments light purple (75A); pistil equalling or exserted from the corolla, 5–10mm longer than the stamens, style deep purplish pink (68A), stigma strong orange-yellow (163B). Lvs elliptic, 90–100 × 35–40mm, margins flat, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (146D) below, hairless; petioles 15mm long. Shrub 4.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101323). Etymology: named for the Società Italiano del Rododendro, which was founded in the Parco Burcina, where its main offices are situated)

(r) **‘Snow Monarch’**

cv. Elepidote rhododendron : ‘Pure Pleasure’ (s) × (*decorum* × *hyperythrum*)
H: R.A. Furman (1997), G: R.A. Furman (2003), N: R.A. Furman (2008), REG: R.A. Furman (2008)
Fls 12/ball truss, broadly funnel-shaped, 75 × 100mm, with 7 wavy-edged lobes and a thick, waxy substance, deep purplish pink (55A) in bud, opening inside white (155B), the throat slightly tinged brilliant yellow-green (150A); outside white (155B); strongly scented. Lvs elliptic, 50–115 × 25–40mm, broadly acute at apex, cuneate at base, margins wavy, glossy and dark green above, hairs on lower surface white when young, aging to light tan. Shrub with an exceptionally dense habit, the foliage hugging the ground, 1.5 × 1.8m in 22 years. Early to mid May.

(r) **‘Sonnenwende’**

cv. Elepidote rhododendron : ‘Golden Splendour’ (s) × (un-named seedling no.192 ‘Edward Dunn’ × *auriculatum*)
H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2007), N: K.-H. Hübbers (2008), I: Baumschulen Wüstemeyer, REG: J. Westhoff (2008)
Fls up to 10/broad, flat, loose truss, open funnel-shaped, 100 × 120mm, with 5–7 very wavy-edged lobes, inside light greenish yellow/pale yellow (8C/D), outside light greenish yellow (8B/C); vivid yellow (12A) to vivid orange-yellow (23A) spots on half of the dorsal lobe (spots dense at the margin and grading into a blotch in the throat); delicately scented. Calyx 50mm long, cut into 5 irregular lobes, coloured like the inside of the corolla, the outside with a strong red (46A) rim. New growth with numerous, long, narrow, red bud scales. Lvs oblong, 200 × 70mm, cuneate and mucronate at apex, cuneate at base, margins decurved, matt green above, hairless. Shrub 1.5 × 1.0m in 5 years. June.
(Nomenclatural standard: colour print of an electronic

image provided by the registrant (WSY0101637).
Etymology: the epithet is German for the solstice
– the plant flowers nearly at the summer solstice)

(r) **'Sophie Elizabeth'**

cv. *Elepidote rhododendron* : 'Miss B.L. Jones' (s) × 'Petra's Debut'
H: D.J. Dosser (2000), G: D.J. Dosser (2006),
N: D.J. Dosser (2007), I: B.S. Coates (2007),
REG: D.J. Dosser (2008)
Fls 20/conical truss, funnel-shaped, 40 × 88mm, with 5 very wavy-edged lobes, buds strong purplish red (63C), opening inside light purple (77D) and turning white with age, outside light purple (77D); slight rays of pale greenish yellow (2D) spotting on the dorsal lobe and with a yellow suffusion at the base of the corolla; slightly scented. Lvs lanceolate, 180 × 64mm, margins slightly wavy, margins decurved, glossy above, truncate at base, acute at apex, hairless. Shrub 1.25 × 1.0m in 7 years. November (Australia).
(Nomenclatural standard: colour print provided by the registrant (WSY0101029). Etymology: named for a granddaughter of the introducer)

(r) **'Sunset Dreams'**

cv. *Elepidote rhododendron* : 'Amber Touch' (s) × 'Rio'
H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2006), REG: J. Barlup (2008)
Fls 14/domed truss, open funnel-shaped, 60 × 85mm, with 5 wavy-edged lobes, deep pink (51B) in bud, opening inside light pink (49C), with a deeper shade of light pink (50D) on the tips of the lobes and a flare of light yellowish pink (26D) extending 45mm upwards from the base on the three dorsal lobes; outside light pink (50D), with deep pink (50B) midribs extending to the edge of the corolla and a blend of light yellowish pink (26D) and light pink (50D) extending 40mm from the base on the dorsal lobes between the midribs; fine, strong red (53D) spotting on the dorsal lobes extending 40mm from the base. Calyx minute. Lvs elliptic, 140 × 45mm, broadly acute at apex, cuneate at base, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.7 × 1.0m in 10 years. Late April.

(r) **'Sun Song'**

cv. *Elepidote rhododendron* : 'Invitation' (s) × 'Cupcake'
H: J. Barlup (1999), G: J. Barlup (2003), N: J. Barlup (2007), REG: J. Barlup (2008)
Fls 15/domed truss, broadly funnel-shaped, 45 × 60mm, with 5 wavy-edged lobes, deep yellowish pink (43C) in bud, opening inside light greenish yellow (4C), fading to pale yellow-green (4D) at the edge of the corolla and with a slight tinge of deep pink (47D) on the tips of the three dorsal lobes; outside light greenish yellow (4C), fading to pale yellow-green (4D) at the edge of the corolla; a few strong red (53D) spots on the dorsal lobe extending 35mm from the base. Calyx 35mm long, light greenish yellow (4C), edged deep pink (47D). Lvs elliptic, 110 × 45mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above,

hairless. Shrub 0.6 × 0.9m in 7 years. Late April.

(r) **'Sun Stream'**

cv. *Elepidote rhododendron* : 'Nancy Evans' (s) × 'Lem's Tangerine'
H: J. Barlup (2000), G: J. Barlup (2004), N: J. Barlup (2007), REG: J. Barlup (2008)
Fls 17/domed truss, open funnel-shaped, 65 × 80mm, with 5 wavy-edged lobes, vivid red (45A) in bud, opening inside and out pale orange-yellow (16D) on the outer edges, shading to a slightly darker light orange-yellow (16C) in the interior, with light yellowish pink (29C) veining on all the lobes; strong red (47B) nectar pouches extending c.10mm from the base. Calyx 25mm long, pale orange-yellow (16D), with strong red (47B) edging. Lvs elliptic, 105 × 55mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.1 × 1.1m in 8 years. Mid May.

(a) **'Tatum's Deep Pink'**

cv. *Deciduous azalea* : wild-collected selection from *occidentale*
S: T. Tatum (1974), N: T. Tatum (1974),
REG: R.W. Cavender (2008)
Fls 10–15/flat truss, tubular funnel-shaped, 50–65 × 55–75mm, with 5 wavy-edged lobes, buds deep pink (52B), in full sun opening inside white, with some light purplish pink (55C) on just the midrib or up to the entire inside of one to five lobes; in shade the light purplish pink is less intense and may appear only on the midrib of one or two lobes; outside very variable, from mostly white with deep purplish pink (55A) on all the midribs to deep purplish pink (55A) covering all or a portion of one to five lobes; in full sun the dorsal lobe is light purplish pink (55C), with spots extending to the lobe margins, in shade the dorsal lobe has much less light purplish pink and has strong orange-yellow (17A) on the lower half; strongly scented. Calyx 3–4mm long, vivid reddish orange (42A). Pedicels also vivid reddish orange (42A). Lvs elliptic, 50–75 × 15–25mm, acute at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above, hairless. Shrub 2.1 × 0.9m in 20 years. Mid May to mid June (with 5–6 fls open at any one time over a 7–15 day period).
(Collected by Tom Tatum (as TT-74-no.1) from Stagecoach Hill azalea reserve, south of Orick, California, USA in 1974)

(a) **'Teeny Weeny'**

cv. *Evergreen azalea* : ('Elsie Lee' × 'Dorsett') (s) × 'Helen Curtis'
H: W.L. Clagett (1996), G: W.L. Clagett (2000),
N: W.L. Clagett (2008), I: White's Nursery (2009),
REG: W.L. Clagett (2008)
Fls hose-in-hose and semi-double, 3/flat truss, broadly funnel-shaped, 30 × 38mm, with 16 wavy-edged lobes, buds strong purplish pink (68B) at tips, shading to white at the base, opening inside nearly white, lightly tinged light purplish pink (68D) mainly at the margins and with a very narrow edge of strong purplish pink (68B) on many lobes; outside similar, but the pink colour is more prominent at the base;

with a pale greenish yellow tinge and spotting in the throat. Calyx c.5mm long, pale green. Lvs elliptic, 30 × 12mm, broadly acute at the apex, cuneate at the base, with flat margins, glossy and moderate olive green (146A) above, hairless. Shrub 0.38 × 0.38m in 11 years. Early May.

(r) **'Terai-shiro'**

cv. Elepidote rhododendron : *degronianum* subsp. *degronianum*, white-flowered × *degronianum* subsp. *degronianum*, white-flowered
H: K. Terai, G: K. Terai (2002), N: K. Terai (2007), REG: K. Terai (2008)
Fls 13–18/hemispherical truss, open funnel-shaped, 70 × 70mm, with 5 smooth or slightly wavy-edged lobes, buds pure white and opening pure white inside and out, with inconspicuous, yellow, dorsal markings. Lvs narrowly oblanceolate, 80 × 25mm, margins recurved, broadly cuneate at base, acute at apex, with a pale fawn indumentum below. Shrub 0.4 × 0.6m in 8 years. Late April.
(Nomenclatural standard: colour print provided by the registrant (WSY0101031). First grafted by the registrant in 1999)

(a) **'Thaissa'**

cv. Evergreen azalea : 'Carol Kittel' (s) × 'Good Times'
H: J.F. Klimavicz (1999), G: J.F. Klimavicz (2001), N: J.F. Klimavicz (2007), REG: J.F. Klimavicz (2008)
Fls 3/ball truss, saucer-shaped, hose-in-hose and semi-double, 25 × 40mm, with 17 wavy-edged lobes, strong to light purple (N80B-C) in bud, opening inside white, shading to light purple (N80D) on the lobe edges; outside strong to light purple (N80B-C); unmarked. Calyx c.4mm long, moderate yellow-green (138B). Lvs obovate, 50 × 20mm, obtuse at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above, hairless. Shrub with a dense habit, 0.45 × 0.45m in 5 years. Mid May.
(Etymology: named for a member of the registrant's family)

(r) **'Tiger's Nest'**

cv. Elepidote rhododendron : selection from *arboreum* subsp. *arboreum*
G: High Beeches Gardens Conservation Trust, N: High Beeches Gardens Conservation Trust (2008), REG: High Beeches Gardens Conservation Trust (2008)
Fls c.14/hemispherical truss, campanulate, 43–47 × c.43–46mm, with 5 slightly wavy-edged lobes, strong red (53C) throughout, with blackish spots on the dorsal side of the throat and on the base of the three dorsal lobes, reduced, paler spotting on and below the other lobes and a small black blotch at the base of the dorsal throat; filaments white, flushed pink right at base, hairless, anthers blackish; style white, suffused purplish pink in the basal half and near the apex, hairless, stigma small, pale purplish pink; ovary densely white hairy, very pale green. Calyx c.1.0mm long, rim-like, densely white hairy, very shallowly lobed. Pedicels densely hairy, tinged pink/red. Lvs elliptic to narrowly elliptic, up to 120 × 33mm,

margins flat to very slightly decurved, acute to acuminate at apex, narrowly cuneate to cuneate at base, with a pale fawn, plastered indumentum below, darkening slightly with age. Shrub c.3.0 × 1.5m in c.20 years. Late April.

(Nomenclatural standard: herbarium specimen provided by the registrant (WSY0112005).

Etymology: named for the Taksang Monastery, Bhutan (known locally as the Tiger's Nest) – the seed from which this plant was raised was collected by Anne and Edward Boscawen on the path up to the monastery at an altitude of 10,000ft (1,600ft above Paro, Bhutan) in April 1987, as AEB162)

(a) **'Toco'**

cv. Deciduous azalea : *occidentale* (Smith & Mossman no. 189) (s) × un-named seedling from 'Gibraltar'
H: K.-H. Hübbers (1988), G: K.-H. Hübbers (1993), N: K.-H. Hübbers (2004), I: Bioplant in Vitro, REG: J. Westhoff (2008)
Fls up to 16/loose, roundish truss, open funnel-shaped, 50 × 80mm, with 5 wavy-edged, somewhat reflexed lobes, inside strong purplish red to strong red (54A–53D), outside strong/deep pink (51A/B); vivid orange (28A) blotch on the dorsal lobe; slightly scented; stamens and style long exerted, prominent. Calyx 2–4mm long, light green. Lvs elliptic, 80 × 37mm, cuneate at apex with a mucronate tip, cuneate at base, margins flat, matt green above, hairless; with good autumn colour. Shrub 1.8 × 0.8m in 10 years. June.

(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101643).
Etymology: named for a city in Trinidad)

(r) **'Tropical Sunser'**

cv. Elepidote rhododendron : 'Apricot Nectar' (s) × 'Lem's Cameo'
H: C. Weeks (1995), G: C. Weeks (2002), N: C. Weeks (2003), I: Westgate Gardens Nursery (2003), REG: C. Weeks (2008)
Fls 8–9/domed truss, funnel-shaped to open funnel-shaped, 65 × 105mm, with 7 wavy-edged lobes, strong purplish red (63A) in bud, opening inside strong purplish red (58C), outside similarly coloured; with a blotch of fine, dark red (59A) spots centred around the filaments on the three dorsal lobes. Calyx 25mm long, strong purplish red (63A), with a dark red (59A) blotch. Lvs elliptic, 125–150 × 70mm, acuminate at apex, cuneate at base, margins wavy and slightly downcurved, dull and moderate olive green (147A) above, hairless. Shrub 1.4 × 1.4m in 12 years, Late May.

(r) **'Twilight Fire'**

cv. Elepidote rhododendron : 'Maricela' (s) × ('Lem's Cameo' × (('Lem's Cameo' × 'Ken Janeck') × 'Brandt's Tropicana') × ('Whopper' × 'Brandt's Tropicana'))
H: J. Barlup (2000), G: J. Barlup (2004), N: J. Barlup (2007), REG: J. Barlup (2008)
Fls 12/domed truss, 55 × 90mm, with 7 very wavy-edged lobes, deep red (53A) in bud, opening inside strong reddish orange (42B), with darker rays

of strong red (46A) extending along the midrib from the base to the outer edge on each lobe; outside similarly coloured; basal markings of deep red (185A), c.10mm wide on the dorsal lobe and c.3mm wide on other lobes; dorsal flare c. 40mm long, consisting of moderate red (185B) spots. Calyx irregular, up to 45mm long, strong reddish orange (42B), with a deep red (185A) base. Lvs elliptic, 125 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and dark yellowish green (139A) above, hairless. Shrub 0.9 × 0.9m in 6 years. Mid May.
Syn: 'Orange 2002'

(r) **'Unforgettable'**

cv. *Elepidote rhododendron* : 'Anna' (s) × 'Nestucca'
H: R.A. Furman (1979), G: R.A. Furman (1984),
N: R.A. Furman (2008), REG: R.A. Furman (2008)
Fls 20/domed truss, open funnel-shaped, c.65 × 90mm, with 5 very wavy-edged lobes, deep pink (52B) in bud, opening inside pale pink (49D), edged strong purplish pink (55B), and with a basal blotch of moderate red (37A); outside deep purplish pink (55A); bracts bright red. Lvs elliptic, 75–175 × 25–50mm, acute at apex, cuneate at base, margins flat, glossy and dark green above, hairy below, the hairs white when young, aging to tan; young foliage copper-coloured. Shrub 1.2 × 1.8m in 17 years, with red colour persistent in the stems. Midseason.

(r) **'Valfenera'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 11–13/conical truss, funnel-shaped, 30–35 × 50–55mm, with 5 wavy-edged lobes, vivid purplish red (N57A) in bud, opening vivid purplish red to deep purplish pink (N57B-C), with deep red to strong purplish red (60A-B) dorsal lobe spotting (just extending to the adjacent lobes); stamens 10, held within the corolla, anthers moderate red to dark yellowish pink (181C-D), filaments deep purplish pink (N66D); pistil equalling or exserted from the corolla, 10mm longer than the stamens, style deep purplish pink (61D), the stigma dark red (59A). Lvs broadly elliptic, 90–95 × 50mm, margins flat, obtuse at base, obtuse and mucronate at apex, matt moderate olive green (146A) above, moderate yellow-green (146D) below, hairless; petioles 20mm long. Shrub 1.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101340). Etymology: named for one of the farmsteads within the Parco Burcina)

(r) **'Valle Elvo'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 16–20/hemispherical truss, funnel-shaped, 50–55 × 50–55mm, with 5 smooth-edged lobes, strong purplish red (67A) in bud, opening with very pale purple (73D) centres and vivid purplish red (67B)

margins, dorsal lobe with strong greenish yellow to vivid yellow (151A-B, 9A) spotting, just extending to the adjacent lobes; stamens 10, held within the corolla, anthers dark purplish pink (186C), filaments strong purplish pink (68B); pistil 2–5mm longer than the stamens but still 10mm shorter than the corolla, style light purplish pink (55C), stigma deep pink (185D). Lvs lanceolate to slightly oblanceolate, 120–130 × 50mm, margins flat, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 15–20mm long. Shrub 4.0m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101342). Etymology: named for the valley in which the Parco Burcina is situated)

(r) **'Vandorba'**

cv. *Elepidote rhododendron* : parentage unknown
G: Parco Burcina, S: E. Accati & N. Furno (2000),
N: E. Accati & N. Furno (2008), REG: E. Accati & N. Furno (2008)
Fls 12–13/hemispherical truss, funnel-shaped, 40 × 60–65mm, with 5 slightly wavy-edged lobes, deep purplish pink (67C) in bud, opening light purple, with a very pale purple centre (75A-D) and with dorsal spotting of brilliant yellow-green (149A), with a few vivid reddish orange (34B) spots at the margins of the spotted area; stamens 10, held within the corolla, anthers deep purplish pink (N66D), filaments deep purplish pink (66C) with lighter shades; pistil equalling or exserted from the corolla, 5–10mm longer than the stamens, style white, stigma vivid red (45C). Lvs lanceolate, 140–165 × 50–55mm, margins flat, acute at base and apex, matt moderate olive green (147A) above, moderate yellow-green (146C) below, hairless; petioles 20mm long. Shrub 2.5m. May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101344). Etymology: named for the stream that runs along the border of the Parco Burcina)

(r) **'Violet Dreams'**

cv. *Elepidote rhododendron* : 'Plum High' (s) × 'Jonathan Shaw'
H: J. Barlup (1997), G: J. Barlup (2001), N: J. Barlup (2007), REG: J. Barlup (2008)
Fls 27/ball truss, broadly funnel-shaped, 45 × 65mm, with 5 wavy-edged lobes, deep purplish red (71A) in bud, opening inside very pale purple (76D) extending from base, edged with strong purple (77B); outside strong purple (77B), with deep purplish red (77A) on the midribs; two deep purplish red (77A) streaks, c.15mm long, at the base of the dorsal lobe, becoming a flare of dark purple (79A) spots extending c.30mm from the base. Calyx vestigial. Lvs elliptic, 155 × 65mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.1 × 1.4m in 11 years. Late May.

(r) **'William Lewis'**

cv. Elepidote rhododendron : 'Lem's Tangerine' (s) × 'Frank Galsworthy'
H: E.F. Newcomb (1999), G: C. Blankenship (2004), S: C. Blankenship, N: C. Blankenship, I: Chuck's Nursery and Hammond Nursery (2010), REG: C. Blankenship (2008)
Fls 20/ball to conical truss, open funnel-shaped, 55 × 70mm, with 5 wavy-edged lobes, buds strong red (53B), opening inside strong red (53D), with the centre of each lobe from the throat to nearly the edge pale purplish pink (56D), with a moderate orange (167B) dorsal flare, speckled strong red (46A); outside strong red (53C). Calyx c.1.5mm long, light yellow-green (145C). Lvs 90–140 × 20–40mm, broadly acute at apex, cordate at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub with a dense habit, 1.0×1.6m in c.10 years. Mid May. (Etymology: named for the grandfather of the registrant)

(a) **'Xi Lin Men'**

cv. Evergreen azalea : sport from 'Pulchrum'
H: C. Zhang (1999), G: C. Zhang (2000), N: C. Zhang, J. Zhang & W. Tian (2006), I: Stock Co. of Bioscience & Technology of Yunnan Green Land (2007), REG: C. Zhang, J. Zhang & W. Tian (2008)
Fls 1–3/open truss, broadly funnel-shaped, 55–60 × 60–70mm, 5-lobed, pinkish red in bud and opening pinkish red throughout, with deep red dorsal spotting. Calyx 5mm long, green. Lvs narrowly oblong, 30–40 × 15–20mm, pointed at apex, broadly cuneate at base, margins flat, hairy (the hairs pale white). Shrub 0.6–0.8 × 1.0m in 20 years, with purple young shoots. April to May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101885))

(a) **'Xue Mei Ren'**

cv. Evergreen azalea : sport from 'Huang Yue'
H: C. Zhang (1998), G: C. Zhang (1999), N: C. Zhang, J. Zhang, Q. Qiao, W. Tian, Z. Wu & B. Sun (2006), REG: C. Zhang, J. Zhang, Q. Qiao, W. Tian, Z. Wu & B. Sun (2008)
Fls solitary, plum blossom shape (a loose double with petaloid stamens and style), 40 × 50–60mm, with 10 or more lobes (the outer whorl very deeply divided), pale green in bud, opening white throughout, unmarked; slightly scented. Calyx 2mm long, pale green. Lvs 20–30 × 10–15mm, shortly acuminate at apex, cuneate at base, hairy (hairs white when young, aging to pale yellow). Shrub with a low spreading habit, 0.3–0.5 × 1.0m in 10 years. Flowers twice, once in April to May and again in November to December (in Kunming, Yunnan, China).
(Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0101886))

(a) **'Zi Yan'**

cv. Evergreen azalea : sport from *simsii*
H: C. Zhang (1985), G: C. Zhang (1988), N: C. Zhang & L. Gao (2006), REG: C. Zhang & L. Gao (2008)
Fls hose-in-hose, 2–3/truss, 30–40 × 30–50mm, with 5 flat-edged lobes, pale purple inside and out, with amaranth (dark reddish purple) markings on the dorsal lobes. Calyx 30–40mm long, pale purple, with irregularly laciniate margins. Lvs obovate to oblanceolate, 40–50 × 15–20mm, shortly acuminate at apex, broadly cuneate at base, margins flat, with brown hairs. Shrub 0.6–0.8 × 1.0m in 20 years. Mid March to May.
(Nomenclatural standard: colour print of an electronic image provided by the registrants (WSY0101890))

Names and addresses

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

* = deceased

Accati, Prof. E.
Università degli Studi di Torino, Grugliasco, Italy

Askjaer, S.A.
Spøttrup, Denmark

Balbo, A.
Leini, Italy

Barlup, J.
Bellevue, Washington, USA

Barratt, J.
Hadsen, Tasmania, Australia

Baumschule Ernst Stöckmann
Bad Zwischenahn-Rostrup, Germany

Baumschule Neumann
Berlin, Germany

Baumschulen Wüstemeyer
Schermbek, Germany

Berg, C.
Silverdale, Washington, USA

*Berg, W.E.
Port Ludlow, Washington, USA

Bioplant in Vitro
Grand-Halleux, Belgium

Blankenship, C.
Arlington, Washington, USA

Blue Mountain Nurseries
Tapanui, West Otago, New Zealand

Bordais, Mme C.
Scaer, France

Briggs Nursery
Olympia, Washington, USA

Burncoose Nurseries
Redruth, Cornwall, England

*Campbell, B.W.
Dunedin, New Zealand

Cavender, R.W. & Mrs K.
Red's Rhodes, Sherwood, Oregon, USA

*Charvet, L.W.
Fort Bragg, California, USA

Chuck's Nursery
Arlington, Washington, USA

Clagett, W.L.
Derwood, Maryland, USA

Coates, B.S.
Wilmot, Tasmania, Australia

Colville, Mrs F.E.
Launceston, Cornwall, England

Cowles, J.C.
Stow, Massachusetts, USA

Descanso Nurseries
Fort Bragg, California, USA

Dosser, D.J.
Lower Wilmot, Tasmania, Australia

Dover Nursery
Sandy, Oregon, USA

Dunedin Rhododendron Group
Dunedin, New Zealand

Elder, J.
Oamaru, New Zealand

Emery, A.
Briggs Nursery, Porter, Washington, USA

Emu Valley Rhododendron Garden
Burnie, Tasmania, USA

Feller, B.
Old Field, New York, USA

Feng, B.
Kunming Institute of Botany, Yunnan, China

Forest Lane Nursery
Fort Bragg, California, USA

Fujioka, F.
Freeland, Washington, USA

Furman, R.A.
Brewster, Massachusetts, USA

Furno, N.
Parco Burcina, Biella, Italy

Gao, L.
Kunming Institute of Botany, Yunnan, China

Garbutt, T.A.
Oamaru, North Otago, New Zealand

George, A.F. & Mrs A.M.
Hydon Nurseries, Godalming, Surrey, England

Glendoick Gardens
Glencarse, Perthshire, Scotland

Grall, R.
Plougar, France

Griswold, D.A.
Kirkland, Washington, USA

Griswold Nursery
Kirkland, Washington, USA

- Gunn, Mrs M.
Chiddingfold, Surrey, England
- Hagan, G. & Mrs K.
Kingston Beach, Tasmania, Australia
- Hammond Nursery
Arlington, Washington, USA
- Harris, J.O.
Lawrenceville, Georgia, USA
- High Beeches Gardens Conservation Trust
Handcross, Sussex, England
- Huang, Y.
Kunming Institute of Botany, Yunnan, China
- Hübbers, K.-H.
Kranenburg, Germany
- Hvidbjerg Planteskole
Hvidbjerg, Denmark
- Hydon Nurseries
Godalming, Surrey, USA
- Jones, K.
Fort Bragg, California, USA
- Kesterson, G.H.
Freeland, Washington, USA
- Klimavicz, J.F.
Vienna, Virginia, USA
- Kruppa, R.
Volcano, Hawaii, USA
- Lehmann, C.A.
Klampenborg, Denmark
- Love, C.
Fort Bragg, California, USA
- Malone, H.R.
Lapoinya, Tasmania, Australia
- Masuda, Y.
Gamou-district, Shiga-prefecture, Japan
- Meerkerk Gardens
Greenbank, Washington, USA
- Meyer, D.
Briggs Nursery, Elma, Washington, USA
- Meyers, W.
Baltimore, Maryland, USA
- Mitchell, R.A.
Volcano, Hawaii, USA
- *Moffat, J.A.R.
Launceston, Cornwall, England
- Mossman, F.D.
Vancouver, Washington, USA
- Moyles, E.W.
Oakland, California, USA
- Navdella, D.
Launceston, Tasmania, Australia
- Nelson's Nursery
Sedro Woolley, Washington, USA
- Newcomb, L.L. & Mrs E.F.
Snohomish, Washington, USA
- *Norman, Lady
Chiddingfold, Surrey, England
- Murcott, R.
East Norwich, New York, USA
- Pacific Island Nursery
Kea'au, Hawaii, USA
- Parco Burcina
(Natural Reserve of Parco Burcina), Pollone,
Biella, Italy
- Potente, R.J.
Muttontown, New York, USA
- Qiao, Q.
Kunming Institute of Botany, Yunnan, China
- Reading, B.
Snohomish, Washington, USA
- Reiley, *H.E. & Mrs M.L.
Woodsboro, Maryland, USA
- RHS Garden, Rosemoor
Rosemoor, Devon, England
- *Rosenthal, J.
Long Island, New York, USA
- Rothschild, *E. de & *L. de
Exbury Gardens, Hampshire, England
- Saint-Jalm, J.F.
Guemene sur Scorff, France
- Saitō, T.
Morioka-city, Iwate-prefecture, Japan
- Schaarup, P.
Silkeborg, Denmark
- Singing Tree Gardens,
McKinlayville, California, USA
- Snell, G.L.S.
Highfields, Queensland, Australia
- Société Bretonne du Rhododendron
Scaer, France
- Stock Company of Bioscience and
Technology of Yunnan Green Land
Kunming, Yunnan, China
- Sun, B.
Kunming Institute of Botany, Yunnan, China
- Terai, K.
Tonami-city, Toyama-prefecture, Japan
- Tian, W.
Kunming Institute of Botany, Yunnan, China
- Tregunna, P.
Formerly head gardener, Caerhays Castle,
Gorran, Cornwall, England

Wallace, D.S.
 Singing Tree Gardens, McKinlayville,
 California, USA

Weeks, C.
 Westgate Gardens Nursery, Eureka,
 California, USA

Westgate Gardens Nursery
 Eureka, California, USA

Westoff, J.
 Bremen, Germany

White's Nursery
 Germanstown, Maryland, USA

Williams, C.H.
 Caerhays Castle, Gorran, Cornwall, England

Williams, F.J.
 Caerhays Castle, Gorran, Cornwall, England

*Williams, J.C.
 Caerhays Castle, Gorran, Cornwall, England

Winberg, J.D.
 Fall City, Washington, USA

Wu, Z.
 Kunming Institute of Botany, Yunnan, China

Yamamoto, Y.
 Gamou-district, Shiga-prefecture, Japan

Zhang, C.
 Kunming Institute of Botany, Yunnan, China

Zhang, J.
 Kunming Institute of Botany, Yunnan, China

Corrections to the International Rhododendron Register and Checklist

The following list indicates further significant additions and corrections that need to be made to the printed text of the *Register and Checklist* published in 2004 or to the four subsequent Supplements.

The Registrar is grateful to all those who have brought such errors to his notice and welcomes knowledge of further amendments that may need to be made.


Minor changes are not noted here but such changes have been made to the database and will be available in any future electronic version.

The database will also be taking into account additional information that comes to light relating to any individual entry, as well as including further entries to the Checklist.

In the following list the page number refers to the page in the *Register and Checklist* (or one of its Supplements) on which the entry concerned is to be found.

- | | |
|------------------|--|
| 'Carlo Contini' | (Third Supplement, p.5).
The WSY number should read WSY0070956 not WSY 0090956 |
| 'Helen Martin' | (First Supplement, p.17–18).
The WSY number should read WSY0041899 not WSY0041900 |
| 'Mendocino Gold' | (Third Supplement, p.16).
The hybridiser should read Charvet not Charbet |
| 'Rajpur' | (p.1133).
The WSY number should read WSY0004518 not WSY0019220 |
| 'Toyama 21' | (Second Supplement, p.42).
The WSY number should read WSY0070566 not WSY0070567 |

ISBN 978-1-907057-00-7



9 781907 057007 >