



Royal  
Horticultural  
Society

# The International Rhododendron Register & Checklist (2004)

*Sixth Supplement*



© 2011 The Royal Horticultural Society  
80 Vincent Square, London SW1P 2PE, United Kingdom  
[www.rhs.org.uk](http://www.rhs.org.uk)

International Rhododendron Registrar:  
Dr A C Leslie

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781907057236

Printed and bound in the UK by Page Bros, Norwich

*Cover: Rhododendron 'Mrs Charles E. Pearson', painted by Alfred John Wise*  
Image courtesy of RHS Lindley Library

**The International  
Rhododendron Register and Checklist (2004)**  
*Sixth Supplement*

Introduction	page 2
Names registered, January 2009 – December 2010	page 3
Names and addresses	page 55
Corrections	page 58

# Introduction

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first five Supplements included names registered from 1 January 2003 to 31 December 2008. This sixth Supplement includes all those registered in 2009 and 2010. A further list of corrections is also included here.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7–10 of the Introduction. It may be helpful here to repeat that the following abbreviations are employed:

(a)	azalea
(az)	azaleodendron
cv.	cultivar
G	grown to first flower by
gp	Group
H	hybridised by
I	introduced by
N	named by
(r)	rhododendron
R	raised by
REG	registered by
S	selected by
(s)	seed parent
(v)	vireya

Where possible, colour references are given to the *RHS Colour Chart* (1966, 1986, 1995 & 2001) with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS website ([www.rhs.org.uk/plants/plant-science/plant-registration-forms/rhodoform](http://www.rhs.org.uk/plants/plant-science/plant-registration-forms/rhodoform)) or from one of the national or regional registrars as follows:

<b>North America</b>	Mrs J.W. Murray, 21 Squire Terrace, Colts Neck, New Jersey 07722-1021, USA
<b>Australia</b>	Mr K. Gillanders, Woodbank, 2040 Huon Road, Longley, Tasmania 7150, Australia
<b>New Zealand</b>	Mr B.J. Coker, 42 Massey Crescent, St Albans, Christchurch 8014, New Zealand
<b>Japan</b>	Dr K. Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken, Japan 768-0021

**Dr A.C. Leslie**

**International Rhododendron Registrar, 109 York Street, Cambridge CB1 2PZ, England (Email: [alanleslie@rhs.org.uk](mailto:alanleslie@rhs.org.uk))**

## Names registered January 2009–December 2010

### (a) 'Adyn'

cv. Deciduous azalea: parentage unknown – a chance seedling

H: D.J. Dosser (1990), G: D.J. Dosser, N: D.J. Dosser (2010), REG: D.J. Dosser (2010)

Fls single, 15–20/ball truss, tubular funnel-shaped, 50 × 70mm, with 5 wavy-edged lobes, buds vivid reddish orange (41A), opening inside and out vivid yellow (15B), dorsal lobe with vivid yellow (17C) spots; slightly scented. Lvs broadly lanceolate, 100 × 40mm, acute at apex, cuneate at base, matt green above, margins flat, hairless. Shrub 1.5 × 0.75m in 20 years. October (Australia).

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the nephew of the registrant

### (a) 'Al Holm'

cv. Evergreen azalea: parentage unknown

H: D.A. Griswold (1999), G: Griswold Nursery (2003), N: Griswold Nursery (2010), REG: Griswold Nursery (2010)

Fls solitary, 40 × c.15mm, elongated tubular funnel-shaped, with 5 wavy-edged lobes, moderate red (160A) in bud, opening strong red (39A) throughout, with speckles of deep yellowish pink (43C) on the dorsal lobes. Lvs oblong, 25 × c.15mm, obtuse at apex, cuneate at base, margins upcurved, dull and moderate olive green (137B) above, hairless. Shrub 1.4 × 1.2m in 11 years. Mid May.

Etymology: named for the father-in-law of Orine Holm, part owner of Griswold Nursery

### (a) 'Allyson Ruth'

cv. Evergreen azalea: 'Myu-no-tsuki' (s) × 'Red Slippers'

H: J.H. Coleman (1992), G: J.R. Hindle (1997), N: R. Holbrook (2008), REG: J.R. Hindle (2009)

Fls 2/truss, open funnel-shaped, 50 × 70mm, with 5 flat-edged lobes, white tinged light purplish pink (65B) in bud, opening inside white, with a broad margin tinged pale purplish pink (56A–C) and marked with four rows of light yellow-green (145C)

spots on the dorsal lobe which fan upwards from the base; outside light purplish pink (65B) at the base, fading to almost white at middle of the lobes and slightly blushed pale purplish pink (56C) at the margins. Calyx c.3mm long, moderate yellow-green (139C). Lvs elliptic, 45 × 15mm, broadly elliptic at apex, cuneate to rounded at base, margins flat, moderate olive green (137A) above, with silvery hairs below when young; loses much of its foliage in winter. Shrub with an open habit, 1.2 × 1.1m in 10 years. Mid April.

Etymology: named for the grand-daughter of Ruth Holbrook

### (r) 'Amber Peach'

cv. Elepidote rhododendron: 'Apricot Fantasy' (s) × 'Hill's Low Red'

H: J. Barlup (1994), G: J. Barlup (1999), N: J. Barlup (2010), REG: J. Barlup (2010)

Fls 13/domed truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, moderate red (47A) in bud, opening throughout light orange-yellow (22D), blending to strong pink (50C) at the margins; vivid red (46B) nectar pouches extend c.6mm from the base on all lobes. Calyx length erratic, up to 25mm long, light orange-yellow (22D), with vivid red (46B) flares extending c.6mm from base on all lobes. Lvs elliptic, 110 × 45mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.1m in 10 years. Early May.

### (r) 'Amber Sun'

cv. Elepidote rhododendron: 'Hotei' × 'Brandt's Tropicana' (s) × 'Hill's Low Red'

H: J. Barlup (1990), G: J. Barlup (1995), N: J. Barlup (2009), REG: J. Barlup (2009)

Fls 11/domed truss, broadly funnel-shaped, 65 × 100mm, with 5 wavy-edged lobes, vivid red (46B) in bud, opening inside pale yellow (20C), with brilliant yellow (20A) on the dorsal lobes which blends with shades of pink to become strong pink (48D) near the margins; outside similarly coloured, but with strong pink (48D) midribs; nectaries deep red (53A), extending c.10mm from the base; anthers very dark; stigma bright green. Calyx c.40mm long, deep

yellowish pink (41D), with a deep red (53A) spot at the base. Lvs elliptic, broadly acute at the apex, rounded at the base, margins upcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 10 years. Early May.

(r) **'Anne Laure'**

cv. Elepidote rhododendron: 'Loderi Helen' × unknown

H: J.-F. Saint-Jalm (1994), G: J.-F. Saint-Jalm (2000), N: J.-F. Saint-Jalm (2000), REG: J.-F. Saint-Jalm (2009)

Fls 12–14/low-domed truss, broadly funnel-shaped, 66 × 110mm, with 7–8 slightly wavy-edged lobes, buds deep purplish pink (N57C), opening inside and out white (155C), with the lobe margins very pale purple (75D), with no markings; stamens 16, filaments white; style strong yellow-green (145A), stigma strong yellow-green (144A); scented. Calyx 2–3mm long, strong greenish yellow (141C), edged moderate red (182A). Lvs elliptic, 105–125 × 38–55mm, margins flat, matt and moderate olive green (137A) above, broadly acute at apex, rounded at base, hairless. Shrub 1.4 × 2.0m in 14 years. Mid April to early May.

Nomenclatural standard: colour print provided by the registrant (WSY0112414). Etymology: named for one of the registrant's daughters

(r) **'Aperitif'**

cv. Elepidote rhododendron: pink-flowered form of an unknown *souliei* hybrid (s) × 'Point Defiance'

H: K.-H. Hübbers (1990), G: K.-H. Hübbers (1998), N: K.-H. Hübbers (2009), REG: J. Westhoff (2009)

Fls c.10/rounded, but very loose truss, funnel-shaped, 50 × 70mm, with 7 wavy-edged lobes, strong red (53C) in bud, opening inside strong pink (54C), with a yellowish band down the middle of each lobe, outside deep purplish pink/strong pink (54B/C), the dorsal throat heavily spotted with dark red (187B). Calyx 2mm long, light green. Lvs oblong, 110 × 40mm, margins flat, wedge-shaped and minutely pointed at apex, truncate to slightly rounded at base, matt green above, hairless. Shrub 2.5 × 2.5m in 20 years. Late April.

(v) **'B.C. Davidson'**

cv. Lepidote rhododendron: 'Island Sunset' (s) × 'Gardenia Odyssey'

H: B. Clancy, G: G. Roelink, N: K. Hagan,

REG: K. Hagan (2009)

Fls 12/rounded truss, tubular funnel-shaped, 50 × 65mm, with 5 flat-edged lobes, inside ground colour vivid reddish orange (33B), the throat vivid orange (28B), with a flare of vivid reddish orange (34A) forming a lobed ring around the centre of the flower, otherwise unmarked, outside vivid reddish orange (41A). Lvs broadly lanceolate, 130 × 50mm, abruptly acute at apex, cuneate at base, glossy green above, margins upcurved, without indumentum. Shrub 0.6 × 0.6m in 4–5 years. Spring/summer. Nomenclatural standard: colour print provided by the registrant (WSY0112038). Etymology: named for Barry Davidson a prominent grower of vireya rhododendrons in Hobart, Tasmania

(a) **'Barbara Tozzi'**

cv. Evergreen azalea: 'Mrs Nancy Dippel' (s) × 'Florence Waldman'

H: J.F. Klimavicz (2000), G: J.F. Klimavicz (2002), N: J.F. Klimavicz (2010), REG: J.F. Klimavicz (2010)

Fls varying from hose-in-hose to semi-double and fully double, 2/truss, saucer-shaped, 50 × 65mm, with 18 very wavy-edged lobes, very pale purple (69A) in bud, opening inside very pale purple (69A), fading to white in the centre, with the base strong to light yellow-green (145A–D), with flecks and a few stripes of deep to strong purplish pink (67C–D); outside very pale purple (69A). Calyx c.30mm long, light yellow-green (145D). Lvs elliptic, 115 × 50mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above, hairless. Shrub with a dense habit, 0.5 × 0.5m in 5 years. Mid May.

Etymology: named for a neighbour of the registrant

(r) **'Beau Vallon'**

cv. Elepidote rhododendron: 'Halopeanum' × 'September Song'

G: J.-F. Saint-Jalm (2003), N: J.-F. Saint-Jalm (2010), REG: J.-F. Saint-Jalm (2010), H: J.-F. Saint-Jalm (1997)

Fls 10–13/domed truss, open-funnel-shaped, 70 ×

90mm, with 5 slightly wavy-edged lobes, strong purplish red (58B) in bud, opening inside pale yellow (8D) in the centre, with broad strong purplish pink (63C) margins, outside strong purplish pink (63C), with strong purplish red (63B) on the dorsal lobe and on half of each adjacent lobe; solid blotch at the base of throat deep purplish red (59B) and with numerous spots of dark greyish purple (N186B) on the dorsal lobe and half of each adjacent lobe; stamens 14, filaments hairy below; style yellow, stigma green; ovary glandular. Calyx petaloid, conspicuous after the corolla has dropped, 8–16mm long, white (N155D), with conspicuous dark purplish red (59B) markings. Lvs elliptic, 140–170 × 50–58mm, margins flat, broadly acute at apex, oblique at base, moderate olive green (137A), hairless. Shrub 2.1 × 1.9m in 12 years. Late April to mid May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the village in which the registrant lives

(r) **‘Ben Moi Bessie’**

cv. *Elepidote rhododendron*: ‘Lemon Lodge’ × *campylocarpum*  
H: I.F. Gordon (pre 2007), G: I.F. Gordon, N: I.F. Gordon, REG: S. Wagstaff (2010)  
Fls 5–7/rounded truss, open funnel-campanulate, 40–50 × 100mm, with 7 wavy-edged lobes, dark reddish orange (172B) in bud, opening inside yellow (light yellow-green (2C) at the base, shading to pale greenish yellow (2D) at the margins), outside yellow (pale greenish yellow (2D) at the base, shading to pale yellow-green (4D) at the margins, with a dorsal flare (c.10 × 10mm) on the dorsal lobe, initially deep red (185A) and fading to strong purplish red (53A); anthers light yellow-green (145D); stigma strong yellow-green (144B); not scented. Calyx 3–4mm long, brilliant greenish yellow (151D). Lvs broadly elliptic, 95–140 × 48–70mm, margins upcurved, apiculate at apex, cordate at base, matt green above, hairless. Shrub 1.0 × 0.9m in 7 years. Mid to late October (New Zealand).

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the raiser’s wife and the name of their garden (Ben Moi)

(r) **‘Beth Beveridge’**

cv. *Elepidote rhododendron*: unknown (pale pink)  
× unknown (white)  
H: G. White (2003), G: G. White (2010), N: G. White (2010), REG: G. White (2010)  
Fls 14/ open-topped truss, funnel-shaped, 45 × 35mm, with 5 slightly wavy-edged lobes, deep red in bud, opening pale cerise inside, with darker dorsal spotting, outside deeper cerise, paling towards the base on both sides. Calyx short, yellowish green, red-rimmed. Lvs elliptic, 135 × 40mm, obtuse to broadly acute at apex, cuneate at base, glossy green above, margins upcurved, sparsely hairy. Shrub 0.5 × 0.5m in 6 years.

Etymology: named for the owner of the parent plants for whom the registrant has gardened for 10 years

(v) **‘Bill Moynier’**

cv. *Lepidote rhododendron*: ‘Gardenia Odyssey’ × *aurigeranum* (s) × *macgregoriae* × *aurigeranum*  
H: W.A. Moynier (1997), G: S. Bertelmann & R. Marques (2003), N: Pacific Island Nursery (2009), I: Pacific Island Nursery (2007), REG: R.F. Kruppa (2009)  
Fls c.12/flat truss, tubular funnel-shaped, 65 × 60mm, with 5 wavy-edged lobes, reddish purple in bud, opening inside moderate yellowish pink (37B) and strong yellowish pink (43D) on the lobes, blending into a light greenish yellow (4C) tube, the colours darkening with age; outside is similar except that the yellowish pink is limited to the outer edges of the lobes; stigma and style bright green; filaments very light yellow; anthers brown on one side, cream on the other. Lvs elliptic, 135 × 65mm, acuminate at apex, cuneate at base, margins upcurved, semi-glossy and dark yellowish green (139A) above, hairless. Shrub 1.2 × 1.5m in 8 years.

Etymology: named for the hybridiser, a noted vireya breeder

(a) **‘Bobbi Gail’**

cv. Evergreen azalea: a double pink florists’ azalea (s) × ‘Treasure’ (Glenn Dale)  
H: B.Y. Morrison (1947), G: B.Y. Morrison, N: W.C. Miller (2008), I: W.C. Miller (2008), REG: W.C. Miller (2009)

Fls single (very occasionally a stamen may be petaloid), 2/flat truss, openly to broadly funnel-shaped, 58 × 60–70mm wide, with 5 broadly overlapping and slightly wavy-edged lobes, white with occasional markings of strong purplish red (71B) in bud, opening inside white with a faint purplish tinge, fading to white, with occasional stripes and sectors of strong purple (77B); dorsal blotch of discrete spots, varying with age from moderate orange-yellow (164B) to moderate purplish pink (186D); outside white with a very pale purplish tinge; self-coloured flowers of strong purple (77B) are not uncommon; stamens usually 7, filaments white, anthers light brown. Calyx normal, lobes 7–9mm long, moderate yellow-green (146D), with a strong yellow-green (144A) margin. Lvs elliptic, 50 × 24mm, acute at apex, cuneate at base, margins flat, dull and slightly more yellow than moderate yellow-green (137C) above, with hairs on the midrib, veins and margin (hairs white below, tan above, all aging to tan). Shrub with a dense habit, 0.84 × 1.37m in 10 years. April to May. Published in *The Azalean* 30(1): 6–7 (2008), with several colour illustrations. Etymology: named for Mrs James B. McCeney (née Bobbi Gail French). She and her husband are members of the Brookside Chapter of the American Rhododendron Society and friends of the registrant

(r) **‘Bon-hime’**

cv. Elepidote rhododendron: ((*yakushmanum* × ‘Sunrise’) × *degronianum* subsp. *degronianum*) × *yakushmanum* (s) × *forrestii* var. *tumescens*  
H: H. Murata (1990), S: H. Murata (1995), G: H. Murata (2002), N: H. Murata (2005), REG: H. Murata (2009)  
Fls (4–)6/truss, open campanulate, 70 × 90mm, with (4–)5(–6) slightly wavy-edged lobes, red in bud, opening inside pink, outside deep pink, unmarked. Calyx 1mm long, yellow-green. Lvs broadly elliptic, 50 × 25mm, rounded at base and apex margins flat, glossy green above, hairless. Dwarf shrub, 23 × 50cm in 18 years (without use of bonsai techniques). Mid April. Nomenclatural standard: colour slide provided by the registrant (WSY0112294)

(r) **‘Boomer’**

cv. Elepidote rhododendron: *hyperythrum* (s) × ‘Sham’s Ruby’  
H: J.T. Thornton (1990), G: L.O. Miller (1995), N: L.O. Miller, REG: L.O. Miller (2009)  
Fls 12/conical truss, broadly funnel-shaped, 40 × 50mm, with 5 wavy-edged lobes, red in bud, opening inside pink, the dorsal lobe with maroon stippling, outside opening pink. Lvs oblong, 115 × 50mm, broadly acute at apex, rounded at base, margins downcurved, glossy and dark green above, hairless. Dense shrub, 2.1 × 2.1m in 15 years. Mid April. Etymology: the name refers back to the late 19th century organised land grant programme in Oklahoma State, when potential claimants would line up behind a starting line waiting for a gun to signal a dash for a favoured plot. The “Boomers” waited for the gun. The “Sooners” jumped the gun

(r) **‘Bovee’s White Water’**

cv. Elepidote rhododendron: *yakushmanum* ‘Koichiro Wada’ (s) × *wardii*  
H: R.M. Bovee (1976), G: R.M. Bovee (pre 1979), N: M. & D. Saner (2009), I: Bovees Nursery, REG: M. & D. Saner (2009)  
Fls 8–10/ball truss, open funnel-shaped, 75 × 50mm, with 5 very wavy-edged lobes, light pink in bud, opening white throughout, with yellow spotting in the throat. Lvs oblanceolate, 75–100 × 25–40mm, acute at apex, oblique at base, margins flat, semi-glossy and dark green above, hairless; petioles red. Shrub 1.2 × 1.5m in 15 years. Mid May. Syn: ‘White Water’  
Originally named ‘White Water’ and introduced by Bovees Nursery under this epithet, but re-named by the Saners to avoid confusion with the earlier ‘Whitewater’

(r) **‘Braxton’**

cv. Elepidote rhododendron: *hyperythrum* (s) × ‘Scintillation’  
H: J.T. Thornton (1993), G: L.O. Miller (1995), N: L.O. Miller (1996), REG: L.O. Miller (2009)  
Fls 11/truss, broadly funnel-shaped, c.40 × 75mm, with 5 wavy-edged lobes, pink in bud, opening inside pink and fading to white, with maroon

stippling on the dorsal lobe; outside light pink, fading to white. Lvs oblong, 100 × 50mm, broadly acute at apex, rounded at base, margins downcurved, glossy and dark green above, hairless. Dense shrub, 1.5 × 2.1m in 14 years. Mid April.

Etymology: named for the grandson of the registrant

(v) **'Brazen Beauty'**

cv. Lepidote rhododendron: 'Gardenia Odyssey' (s) × *laetum*

H: B.E. Oldham (1993), G: B.E. Oldham (2000), N: B.E. Oldham (2001), REG: B.E. Oldham (2009)  
Fls 5–7/full, shallow-domed truss, tubular funnel-shaped, 60 × 90mm, with 5 broadly overlapping, wavy-edged lobes, buds brilliant yellow-green (149C), with strong purplish red (61B) bracts, opening inside brilliant orange-yellow (23B) blending into light orange-yellow (22A), outside brilliant yellow (21C) on the lobes, strong purplish red (58B) towards the top of the tube, blending to brilliant yellow (10A) at the base; marked with strong purplish red (58B) at the lobe bases both inside and out; not scented. Calyx less than 1mm long, green. Lvs broadly elliptic to elliptic, 130 × 60–65mm, margins flat, apiculate at apex, attenuate at base, glossy and dark olive green above, scaly (the scales light yellow-green, 145B, becoming more sparse with age and giving the young leaves a greyish tinge). Dense, fastigiate shrub, 1.9 × 1.0m in 15 years. Frequent repeat-flowering.  
Nomenclatural standard: colour print provided by the registrant (WSY0112741). Etymology: named after comments about "its brazen brilliance"

(r) **'Bruce Briggs'**

cv. Elepidote rhododendron: 'Seaview Sunset' (s) × *peudochrysanthum* × 'Etta Burrows'  
H: F. Fujioka (1998), G: F. Fujioka (2004), N: Briggs Nursery (2009), I: Briggs Nursery (2009), REG: F. Fujioka (2009)  
Fls 26/flat truss, funnel-campanulate, 50 × 40mm, with 5 wavy-edged lobes, vivid reddish orange (40B, edged 40A) in bud, opening inside vivid reddish orange (40A) at margin, changing to moderate pink (35D) from margin to throat, with a small, strong red (46A) blotch in the dorsal throat; outside vivid

reddish orange (40A) at margin, gradually fading to pale yellowish pink (36D) at the base. Calyx 25mm long, pale yellowish pink (36D), with vivid red (44A) stripes. Lvs elliptic, 165 × 60mm, broadly acute at apex, rounded at base, margins downcurved, dull and dark yellowish green (136B) above, with very fine hairs below (hairs pale yellow (160D) at first, aging to deep orange-yellow (163A)). Shrub 0.9 × 0.8m in 10 years. Late April.

Etymology: named for the founder of Briggs Nursery (USA)

(v) **'Bruce Winter'**

cv. Lepidote rhododendron: (*macgregoriae* × *aurigeranum*) × 'Doctor Hermann Sleumer' (s) × *konori* × *christianae*

H: R.A. Mitchell (2002), G: R.A. Mitchell, N: R.A. Mitchell (2010), REG: R.A. Mitchell (2010)  
Fls 6–7/flat truss, tubular funnel-shaped, 65 × c.65mm, with 5–7 very wavy-edged lobes, lobes open inside and out strong red (51A), shading to pink and then yellow-green down the throat and tube, with brownish yellow spots (c.1.8mm wide) at lobe margins inside and out; anthers very dark brown, filaments very light yellow; stigma light green; style very light yellow; slightly scented. Calyx absent. Lvs elliptic, 140 × 55mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (137B) above, without visible indumentum. Shrub 0.9 × 0.6m in 7 years. January.  
Etymology: named for a good friend of the registrant and a charter member of the ARS Hawaiian Chapter

(r) **'Burgundy Lace'**

cv. Elepidote rhododendron: 'Anna' × 'Frank Galsworthy' (s) × (*yakushimanum* × 'Burgundy') × 'Purple Splendour'  
H: F. Fujioka (1993), G: F. Fujioka (2009), N: F. Fujioka (2009), REG: F. Fujioka (2009)  
Fls 25/ball truss, broadly funnel-shaped, c.70 × 100mm, with 5 very wavy-edged lobes, buds strong purplish red (63A) at base for c.15mm, then changing to dark red (59A); opening inside vivid purplish red (61C) from the throat to 6mm from the edge, which is dark red (59A) except for the dorsal

lobe which is edged pale purplish pink (62D); outside strong purplish red (63A) at base and changing to dark red (59A) after 15mm; large, rather dense spotted blotch of moderate orange (167B) and dark yellowish green (189A) on the dorsal lobe. Lvs elliptic, 150 × 60mm, broadly acute at apex, rounded at base, margins slightly downcurved, semi-glossy and dark bluish green (131A) above, hairless. Shrub with a dense habit, 1.2 × 1.5m in 9 years. Late May.

(r) ‘**Caerhays Crossbill**’

cv. Lepidote rhododendron: *spinuliferum* (s) × *lutescens*

H: J.C. Williams (1920s), G: J.C. Williams,

N: C.H. Williams & J. Parsons (2009),

REG: C.H. Williams & J. Parsons (2009)

Fls up to 15/truss, funnel-shaped, up to 28mm long, 6-lobed, light yellow (10C/160B) to pale greenish yellow (1D), tinged a bit more pink than strong to moderate yellowish pink (34A/B); stamens 10, filaments near strong red (53C), shading to cream at both ends, anthers moderate to light reddish brown (177A/B); style exerted, coloured as the filaments. Calyx c.1mm long, green, pubescent. Pedicels c.12mm long, green, tinged near deep yellowish pink (47C). Calyx, pedicels and ovary white hairy and with green/yellow scales. Lvs elliptic, up to 87mm long, acuminate at apex, green, tinged near to dark red (187A), hairless above, with a few hairs on the veins and midrib below and with brown scales.

Vigorous shrub, 3.0–3.8 × 1.9–2.4m in 10–15 years. Late March.

PC 2009

Nomenclatural standard: herbarium specimen of the PC plant (WSY0112838)

(r) ‘**Candycorn**’

cv. Elepidote rhododendron: ‘Nancy Evans’ × ‘Orange Marmalade’ (s) × ‘Orange Marmalade’ × ‘Lem’s Cameo’

H: D.S. Wallace (2001), G: D.S. Wallace (2006),

N: D.S. Wallace (2008), I: Singing Tree Gardens and Nursery (2010), REG: D.S. Wallace (2010)

Fls 15/domed truss, funnel-campanulate, 50 × 65mm, with 5 wavy-edged lobes, vivid red (45B) in

bud, opening inside brilliant yellow (15C), shading to deep yellowish pink (44D) at the edges, with strong red (46A) nectar pouches; outside brilliant yellow (15C) towards the base, shading to deep yellowish pink (44D) towards the edges; unmarked. Calyx 20mm long, light greenish yellow (8B). Lvs elliptic, 100 × 45mm, broadly acute at apex, cuneate at base, margins flat, glossy and strong green (132B) above, hairless. Shrub 0.6 × 0.8m in 10 years. Mid May.

(r) ‘**Carine de Beau Vallon**’

cv. Elepidote rhododendron: ‘Nancy Evans’ × *fortunei*

H: J.-F. Saint-Jalm (1999), G: J.-F. Saint-Jalm

(2006), N: J.-F. Saint-Jalm (2010), REG: J.-F. Saint-

Jalm (2010)

Fls 11–13/ball truss, open funnel-shaped, 60–65 × 85–90mm, with 7 wavy-edged lobes, deep purplish pink (64D) in bud, opening inside light yellow (10C), with broad margins of strong purplish pink (73B), outside light yellow (10C) at base, with deep purplish pink (54B) margins; solid blotch of strong red (46A) at the base of the throat; stamens 14, filaments hairy at the base; ovary glandular. Calyx 2–15mm long, strong red (46A) and pale greenish yellow (10D). Lvs elliptic, 140–180 × 60–75mm, margins decurved, broadly acute at apex, rounded at base, moderate yellowish green (139B) above, hairless. Shrub 1.8 × 1.0m in 11 years. Late April to early May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the second daughter of the registrant and the village in which they live

(r) ‘**CeeCee**’

cv. Elepidote rhododendron: ‘Vulcan’ (s) × *hyperythrum*

H: S.B. Southerland (1997), G: S.B. Southerland

(2000), N: S.B. Southerland (2000),

REG: S.B. Southerland (2010)

Fls up to 11/domed truss, open funnel-shaped, 45 × 55mm, with 5 flat-edged lobes, close to deep purplish pink (61D) throughout, unmarked. Lvs elliptic, 140 × 50mm, broadly acute at apex,

rounded at base, margins flat, dull and dark green above, hairless. Shrub with a dense habit, 1.5 × 1.8m in 12 years; heat and insect tolerant. Early April. Etymology: nickname of the registrant's wife – her maiden name being Carolyn Cook

(v) '**Charlotte Rouse**'

cv Lepidote rhododendron: *loranthiflorum* (s) × *zoelleri*

H: J.L. Rouse (1980), G: J.L. Rouse, N: A. Rouse (2006), REG: A. Rouse (2006)

Fls 5–7/open truss, tubular-campanulate, 50 × 40mm, with 5 very narrowly wavy-edged lobes, buds light yellow-green (2C), opening inside light pink (49C) on lobes and with a pale yellow-green (4D) tube, outside light pink (49C) towards rim of lobes, otherwise pale yellow-green (4D), unmarked. Calyx less than 1mm long. Lvs broadly elliptic to oblanceolate, 45–65 × 20–30mm, margins wavy, glossy green above, cuneate at base, acute at apex, scaly. Shrub 2.0 × 1.0m in 13 years. July to August (Australia).

Nomenclatural standard: colour print provided by the registrant (WSY0096140)

Registered in 2006 but accidentally omitted from the Register supplements until now

(r) '**Chemainus**'

cv. Elepidote rhododendron: 'Noyo Brave' (s) × ('C.I.S.' × *rex*) × 'Hansel'

H: F. Fujioka (1985), G: F. Fujioka (1992),

N: F. Fujioka (2010), REG: F. Fujioka (2010)

Fls 18/domed truss, funnel-campanulate, 50 × 75mm, with 5 wavy-edged lobes, vivid red (52A) in bud, opening inside deep purplish pink (55A), with deep red (53A) spots on the dorsal lobe becoming solid near the throat; outside deep pink (52C); not fading with age. Calyx 25mm long, strong pink (52D). Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, glossy and dark bluish green (131A) above, with a slight silvery tomentum above, hairy all over below (hairs moderate orange-yellow (164B), at first, aging to moderate orange-yellow (165C)). Dense shrub, 1.2 × 1.4m in 10 years. Early. Etymology: named for a small town on the east coast of Vancouver Island, British Columbia, Canada

(r) '**Cheryl Inger**'

cv. Elepidote rhododendron: 'Lockington Pride' × 'Loderi Titan'

H: D.J. Dosser (2000), G: D.J. Dosser,

N: D.J. Dosser (2010), REG: D.J. Dosser (2010)

Fls 14/ball truss, funnel-shaped, 65 × 50mm, with 7 wavy-edged lobes, often with unevenly reflexed margins and tips, buds white (158D), opening white (158D) throughout, dorsal lobes with a ray and spots of moderate red (186B), the marking extending around the base of the throat; scented. Lvs oblanceolate, 130 × 50mm, rounded at apex, cuneate to narrowly rounded at base, margins flat, matt green above, hairless. Shrub 1.5 × 1.0m in 10 years. October (Australia).

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the sister of a friend of the registrant

(r) '**Cheryl Renee Jonsson**'

cv. Elepidote rhododendron: parentage unknown

H: D.A. Griswold (1969), G: Griswold Nursery

(1985), N: Griswold Nursery (2010),

REG: Griswold Nursery (2010)

Fls 14/conical truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, light purplish pink (62C) in bud, opening pure white throughout (close to 155C), unmarked. Lvs oblong, 75 × 40mm, broadly acute at apex, cuneate at base, margins downcurved, dull and dark yellowish green (139D) above, hairless. Shrub with an open habit, 1.5 × 0.9m in 41 years. Mid May.

Etymology: named for the wife of the local Pastor, a good friend of the Griswold Nursery

(r) '**Cinq Jours de Bonheur**'

cv. Elepidote rhododendron: parentage unknown

H: L. Le Duigou (1997), G: A. Le Duigou (2007),

N: L. & A. Le Duigou (2009), REG: L. & A. Le Duigou (2009)

Fls 20/rounded truss, open funnel-shaped, 45 × 65mm, with 5 slightly wavy-edged lobes, vivid purplish red (61C) in bud, opening white (155D), with a tinge of pink towards the edge of the lobes and a conspicuous dorsal blotch of dark purplish red (N79A) spots; stamens 9; style white (155D). Calyx

2mm long, strong yellow-green (145A). Lvs elliptic, 125 × 45mm, margins decurved, acute at base and apex. Slow-growing, compact and rounded shrub, 0.75 × 0.8m in 12 years. May.

Nomenclatural standard: colour print provided by the registrant (WSY0112738). Etymology: named to reflect the fact that cut trusses last for at least 5 days in a vase

(r) **'Copperhill'**

cv. *Elepidote* rhododendron: 'Lem's Cameo' (s) × 'Deerah'

H: K.-H. Hübbbers (1992), G: K.-H. Hübbbers (2000), N: K.-H. Hübbbers (2005), I: Baumschule Neumann & Stöckmann Baumschulen (2008), REG: J. Westhoff (2009)

Fls up to 14/ rounded but slightly loose truss (flat above), open campanulate, 40 × 50mm, with 5 wavy-edged lobes, buds moderate red (47A), opening inside brilliant yellow (13C) in the throat and shading to light yellowish pink (36A) at the end of the lobes, outside deep pink (51B), with parts of the dorsal lobes light orange-yellow (16C); spots on the dorsal part of the throat deep red (53A); slightly scented. Calyx up to 3mm long, reddish. Lvs ovate, 80 × 35mm, slightly rugose, cuneate with a very small point at the apex, rounded at base, margins flat, glossy above, hairless; new foliage copper-red. Small, slow-growing shrub, 0.9 × 0.8m in 10 years. April.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0104990). Etymology: derived from the appearance of the plant and the place in which it grows in the garden where it was raised

(r) **'Cordula Reich'**

cv. *Lepidote* rhododendron: *ambiguum* × *cinnabarinum* Concatenans Group × *cinnabarinum* Roylei Group

H: W. Reich (1992), G: W. Reich (2000), N: W. Reich (2004), REG: W. Reich (2010)

Fls up to 12/open-topped truss, tubular-campanulate, 45–50 × 38–43mm, with 5 flat-edged lobes, buds pinkish lilac to strong red or deep pink (50A–B), opening inside light orange (28C), outside

light orange (28C) at margin, shading to moderate reddish orange/strong yellowish pink (35B–C) and to strong red/deep pink (50A–B) on the tube; darker reddish veins both inside and out. Calyx 2–3mm long, greenish. Pedicels greenish. Lvs elliptic, 44–74 × 22–30mm, broadly acute at apex, cuneate at base, margins flat, glossy bluish green above and with prominent veins, with a scaly indumentum. Shrub 1.6 × 1.2m in 15 years. Mid to late May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the daughter of the registrant

(r) **'Cowles Pinwheel'**

cv. *Elepidote* rhododendron: parentage unknown G: Heritage Museums and Gardens (1970s), S: Sandwich Club (2009), N: Sandwich Club (2009), REG: Sandwich Club (2009)

Fls 15/domed truss, open funnel-shaped, c.75 × 90mm, with 5 wavy-edged lobes, strong purplish red (63A) in bud, opening inside pale yellow-green (155A) in the centre, with strong to vivid purplish red (61B–C) margins and a small, strong greenish yellow (151B–C) dorsal flare; outside strong purplish red (63A). Lvs oblong, 125 × 50mm, broadly acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 8 years. Late May.

(r) **'Dathocarmapie'**

cv. *Lepidote* rhododendron: *cinnabarinum* (s) × unknown

H: Mme M.T. Bleuzen (2000), G: Mme M.T. Bleuzen (2010), N: Mme M.T. Bleuzen (2010), REG: Mme M.T. Bleuzen (2010)

Fls 5/lax truss, funnel-campanulate, 45 × 60mm, with 5 smooth-edged lobes, brownish orange (164A) in bud, opening strong reddish orange (31B) inside, strong reddish orange (31A) outside, unmarked; stamens orange-pink; style orange-red. Calyx 3mm long, light yellowish green (135D). Lvs oblong, 60 × 30mm, rounded at base, broadly acute at apex, margins flat, glossy green above, with a scaly light olive grey (197B) indumentum below, which

ages to light olive brown (199B). Compact shrub, 0.9 × 0.8m in 10 years. Mid to late April.  
Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: the epithet comprises parts of the names of the registrant's five grandchildren Damien, Thomas, Carla, Malo and Pierre

(r) **'Dear Barbara'**

cv. Elepidote rhododendron: 'Vulcan's Flame' × *yakushimanun* 'Koichiro Wada'  
H: T. Huisman (1986), G: T. Huisman (1991),  
N: T. Huisman (2009), I: T. Schröder (2010),  
REG: T. Huisman (2009)  
Fls 10–12/ball truss, campanulate to funnel-campanulate, 65–80 × 60–70mm, with 5 wavy-edged lobes, buds vivid purplish red (N57A to N57B), opening inside strong purplish red (58B), the margins very light pink to almost white, outside vivid purplish pink (N66B) at the base with pale purplish pink (65D) margins; unmarked; after a week the flowers age to champagne white with paler margins. Calyx very short. Lvs lanceolate, 70–110 × 20–30mm, acute at apex, cuneate at base, convex but with margins flat, dull green above, hairless. Shrub 1.0 × 1.2m in 10 years. May.  
Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0108029). Known as seedling no. Z 240.  
Etymology: named for a former director of the American Rhododendron Society, Barbara Hall

(r) **'Deenah'**

cv. Elepidote rhododendron: 'Graf Lennart' (s) × 'Blewbury'  
H: K.-H. Hübbbers (1987), G: K.-H. Hübbbers (1992), N: Mrs L. Hübbbers (1999),  
REG: J. Westhoff (2009)  
Fls 7–10/uneven, flat, loose truss, campanulate, 30 × 30mm, with 5 wavy-edged lobes, buds moderate red (47A), opening inside pale yellow-green (4D), with deep red (53A) spots on two-thirds of the dorsal throat (showing through to the outside), outside light greenish yellow (4B), with a deep pink (48A) stripe on each lobe. Calyx 2–3mm long, greenish. Lvs ovate, 50–90 × 30–40mm, mucronate

at apex, rounded at base, margins decurved, glossy above, hairless. Shrub 1.0 × 1.0m in 15 years. Early to mid April.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0104989). Etymology: named for a niece of the raiser's wife

(r) **'Deerah'**

cv. Elepidote rhododendron: 'Graf Lennart' (s) × 'Blewbury'  
H: K.-H. Hübbbers (1987), G: K.-H. Hübbbers (1992), N: Mrs L. Hübbbers (1999),  
REG: J. Westhoff (2009)  
Fls c.12/rounded truss (flattened on top), open campanulate, 35 × 40mm, with 5 wavy-edged lobes, buds deep yellowish pink (39B), opening pale yellow-green (4D) both inside and out, with deep red (53A) spots on two-thirds of the dorsal throat which merge into two bands. Calyx minute, light green. Lvs ovate, 50 × 25mm, mucronate at apex, rounded at base, margins flat, glossy above, hairless. Shrub 1.0 × 0.8m in 15 years. Early to mid April.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0104088). Etymology: named for a niece of the raiser's wife

(r) **'Diana Marguerite'**

cv. Elepidote rhododendron: 'Consolini's Windmill' × 'Unique Marmalade' (s) × 'Off-shoot' × ('Phipps Yellow' × 'Viennese Waltz')  
H: G. Woodard (2002), G: G. Woodard (2004),  
N: T. Weekes (2009), I: G. Woodard,  
REG: G. Woodard (2009)  
Fls 22/conical truss, broadly funnel-shaped, 100 × 100mm, with 6–7 wavy-edged lobes, light pink (49C) in bud, opening inside vivid yellow (17B), with subtle picotee edges of dark pinkish red at first, fading to pink; outside light yellowish pink (38D). Lvs elliptic, 125 × 60mm, broadly acute at apex, oblique at base, margins downcurved, semi-glossy and dark green above, hairless. Shrub 1.2 × 0.9m in 7 years. Mid May.

(r) **'Donna Mae Eagle'**

cv. Elepidote rhododendron: 'Lem's Goal' × 'Goldsworth Orange' (s) × 'Whitney's Orange'  
H: D.A. Griswold (1969), G: Griswold Nursery (1982),  
N: Griswold Nursery (2010), REG: Griswold  
Nursery (2010)

Fls 10/lax truss, funnel-campanulate, c.50 ×  
c.50mm, with 5 wavy-edged lobes, vivid reddish  
orange (43B) in bud, opening inside pale yellow  
(18C) in the centre, tinged pink, with vivid reddish  
orange (34B) dorsal spots and occasional randomly  
distributed spots of the same colour on the other  
lobes; outside light yellowish pink (38C). Calyx  
6mm long, vivid reddish orange (43B). Lvs  
lanceolate, 100 × 50mm, broadly acute apex,  
cuneate at base, margins downcurved, dull and  
moderate yellow-green (147B) above, hairless. Shrub  
with an open habit, 1.7 × 1.8m in 41 years. Mid May.  
Etymology: named for a local landscape designer  
and friend of the Griswold Nursery

(r) **'Double Dots'**

cv. Elepidote rhododendron: 'Whitney's Tiger Lily'  
× 'Graf Lennart' × *calophytum*  
H: T. Huisman (1995), G: T. Huisman (2000),  
N: T. Huisman (2009), I: T. Schröder (2010),  
REG: T. Huisman (2009)

Fls 5–8/slightly loose ball truss, open funnel-  
shaped, 60–75 × 65–80mm, with 5 wavy-edged  
lobes, buds light purple (75B), opening inside light  
greenish yellow (4C) in the throat, shading to light  
pink or white at the margins, outside very pale  
purple (75D) at the base, becoming light pink to  
almost white at the margins; strong purplish red  
(67A) spots on all the lobes, especially on the dorsal  
lobe and these show through as vivid purplish red  
(67B) spots on the outside of the corolla. Calyx  
30–40mm long, petaloid, also spotted all round as  
in the corolla. Lvs lanceolate, 70–110 × 20–30mm,  
cuneate to rounded at base, broadly acute at apex,  
margins flat, matt green above, hairless. Shrub 1.5 ×  
1.2m in 10 years. Mid May.

Nomenclatural standard: colour print provided by  
the raiser (WSY0108030). Grown as seedling no. Z  
512. Etymology: the name derives from the extensively  
spotted corolla and similarly spotted, petaloid calyx

(a) **'Double Talk'**

cv. Deciduous azalea: 'Homebush' (s) × unknown  
H: J. Neal (1970s?), G: J. Neal (1978), N: J. Neal  
(1980), REG: Mrs M.L. Reiley (2010)

Fls double, 12/domed truss, narrowly funnel-shaped,  
50 × 75mm, with 11–12 very wavy-edged lobes,  
vivid reddish orange (34B), with vivid yellow (14A)  
midribs in bud, opening inside light orange-yellow  
(16B), suffused strong reddish orange (34C); outside  
similar to inside, with the addition of vivid yellow (14A)  
midribs and a green base; unmarked; stigma bright  
green; stamens exerted 25mm beyond the corolla.  
Calyx c.3mm long, pale green. Lvs elliptic, 50–65 ×  
25mm, acute at apex, cuneate at base, margins  
downcurved, semi-glossy and moderate yellow-green  
(137C) with bronze overtones above, with colourless  
hairs above and below. Shrub with an open habit,  
2.1–2.4 × 0.9–1.2m in 15 years. Early April.

(r) **'Duane Johnson'**

cv. Elepidote rhododendron: 'Butter Brickle' (s) ×  
'September Song'  
H: J. Winberg (1990), G: J. Winberg (1994),  
N: E. Sather (2007), I: Hazelwood Gardens (1998),  
REG: J. Barlup (2009)

Fls 13/flat truss, broadly funnel-shaped, 50 ×  
75mm, with 5 wavy-edged lobes, moderate red  
(181A) in bud, opening inside vivid yellow (16A)  
and changing to strong yellowish pink (37A) on the  
edges of the corolla; outside opens strong reddish  
orange (34C) and changes to strong yellowish pink  
(37A) on the edges; deep red (53A) nectaries  
extending c.10mm from the base. Calyx c.15mm  
long, vivid yellow (16A), with two deep red (53A)  
flares c.5mm long. Lvs elliptic, 125 × 45mm, broadly  
acute at apex, rounded at base, margins downcurved,  
dull and moderate olive green (147A) above, hairless.  
Shrub 2.3 × 3.1m in 18 years. Mid May.

(v) **'Easter Pastel'**

cv. Lepidote rhododendron: *aurigeranum* (s) ×  
'Pink Delight'

H: P. Sullivan (pre 2000), G: S. Bertelmann &  
R. Marques (2003), N: S. Bertelmann & R. Marques  
(2009), I: Pacific Island Nursery (2009),  
REG: R.F. Kruppa (2009)

Fls 10–14/flat truss, tubular funnel-shaped, 60 × 60mm, with 5 flat-edged lobes, opening inside with a wide edge of pale pink (49A) and pale purplish pink (56A), blending into a very light yellow that brightens into the pale yellow (20C) throat and tube; outside similar, with multiple shades of pink; anthers brown, filaments light yellow; stigma green, style light green. Lvs elliptic, 140 × 50mm, acute at apex, cuneate at base, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub with an open, upright habit, 0.7 × 0.4m in 5 years. Spring and autumn.

(r) **‘Ed Reiley’s Delight’**

cv. Elepidote rhododendron: ‘Helen Everitt’ (s) × ‘Harvest Moon’  
G: H.E. Reiley (1960s), N: Mrs M.L. Reiley (2010), REG: Mrs M.L. Reiley (2010)  
Fls 7–9/domed truss, broadly funnel-shaped, 65 × 90mm, with 6 wavy-edged lobes, greenish white (155C) in bud, opening yellowish white (155D), with a basal blotch of light yellow-green and a few reddish brown streaks c.10–15mm long; outside yellowish white (155D); stigma strong yellow-green (144A), style strong yellow-green (145A); ovary strong yellowish green (141C); strongly scented. Calyx c.3–6mm long, strong yellow-green (144B). Lvs oblong, 90–100 × 45mm, obtuse at apex, rounded at base, margins flat, semi-glossy and moderate olive green (137A) above, the veins strong yellow-green (143C), hairless. Shrub 1.8 × 2.4m in 40 years. April to May.  
Hybridiser uncertain: possibly A.M. Shammarello or J.B. Gable

(r) **‘Ed’s Fall Frolic’**

cv. Elepidote rhododendron: *yakushimanum* × ‘Mary Belle’ (s) × ‘Crest’ × ‘Evening Glow’  
H: H.E. Reiley (2002), G: H.E. Reiley (2005), N: Mrs M.L. Reiley (2009), REG: Mrs M.L. Reiley (2009)  
Fls 10/domed truss, funnel-campanulate, 40 × 60mm, with 6 wavy-edged lobes, moderate reddish orange (35B) in bud, opening inside light greenish yellow (3D) with deep orange-yellow (163A) freckles on the three dorsal lobes; outside light greenish yellow (3D) with moderate pink (35D) highlights

that fade to light greenish yellow (3D). Calyx c.30mm long, light greenish yellow (3C). Lvs oblong, 95–125 × 40–50mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (137A) above, hairless. Shrub 0.8 × 1.2m in 7 years. October to November.

(r) **‘Elizabeth Mathews’**

cv. Elepidote rhododendron: ‘Beauty of Lockington’ (s) × ‘Tortoiseshell Wonder’ × ‘Ross Maud’  
H: D.J. Dosser, G: D.J. Dosser (2009), N: D.J. Dosser (2010), REG: D.J. Dosser (2010)  
Fls 12/ball truss, funnel-shaped, 50 × 78mm, with 6 very wavy-edged lobes, buds strong purplish red (63B), opening inside white, outside pale purplish pink (65D) and white, vivid purplish red (57B) spots on the three dorsal lobes. Calyx 14mm long, strong purplish red (63B). Lvs lanceolate, 150 × 38mm, with strongly reflexed and irregularly wavy margins, broadly acute at apex, truncate at base, matt green above, hairless. Shrub 1.0 × 0.75m. November (Australia).  
Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a friend of the registrant

(r) **‘Ericho’**

cv. Elepidote rhododendron: ‘Barbara Reuthe’ (s) × ‘New Comet’  
H: K.-H. Hübbers (1989), G: K.-H. Hübbers (1996), N: K.-H. Hübbers (2004), REG: J. Westhoff (2009)  
Fls 10/very loose truss, broadly funnel-shaped, 60 × 80mm, with 7 wavy-edged lobes, buds dark red (187B) at apex, greenish yellow at the base, opening bicoloured: inside brilliant yellow (8A) ground, mottled deep to strong red (53A–C), especially on the dorsal lobes which have red patches, outside strong to deep pink (51 A–C), with a yellow base and veining. Calyx 15mm long, coloured like the outside of the corolla. Lvs oblong, 120 × 60mm, rounded and with a slight point at apex, slightly rounded at base, margins flat, matt above, hairless. Shrub 1.5 × 1.5m in 10 years. Late April.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0104987). Etymology: named after a brother of the raiser (Erich Hübbers)

(r) **'Evening Embers'**

cv. Elepidote rhododendron: 'Plum High' (s) × 'Olin O. Dobbs'

H: J. Barlup (1996), G: J. Barlup (2000), N: J. Barlup (2009), REG: J. Barlup (2010)

Fls 18/ball truss, broadly funnel-shaped, 50 × 70mm, with 6 wavy-edged lobes, dark red (59A) in bud, opening inside and out strong purplish red (60B), shading to darker edges of deep red (60A) and with spotted dark greyish reddish brown (200A) flares on two dorsal lobes which start from c.10mm from the base and extend c.25mm. Lvs elliptic, 120 × 50mm, broadly acute at apex, rounded at base, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 10 years. Late May.

(r) **'Faro'**

cv. Elepidote rhododendron: 'Manglesii' (s) × unknown

H: A. Le Duigou (1997), G: A. Le Duigou (2007), N: L. & A. Le Duigou (2010), REG: L. & A. Le Duigou (2010)

Fls 19/conical truss, broadly funnel-shaped, 65 × 100mm, with 5–6 flat-edged lobes, buds strong reddish purple (72C), opening inside very pale purple (69D), with the margins very light to light purple (75C–75A), outside very pale purple (76C), with five light purple (75A) stripes; heavy and conspicuous spotting of deep greenish yellow (153A) on the dorsal lobes; stamens 9; style moderate purplish pink (65A). Lvs oblanceolate, 190 × 56mm, broadly acute at apex, cuneate at base, margins flat, matt green above. Shrub 2.2 × 3.0m in 13 years. May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: the epithet is the Breton word meaning proud

(r) **'Forest Sprite'**

cv. Lepidote rhododendron: *augustinii* (s) × *keiskei* var. *ozawae* 'Yaku Fairy'

H: B.N. Starling (2002), G: B.N. Starling (2007), N: B.N. Starling, REG: B.N. Starling (2009)

Fls up to 7/shoot tip (from a terminal and sometimes also lateral buds), funnel-shaped, up to 25 × 40mm,

with 5–6 flat-edged lobes, buds deep purplish pink (73A), opening inside light purple (75A), fading to white in the throat, with brown flecking on the dorsal lobes, outside light purple (75A), the tips of some lobes white; stamens up to 12, filaments white (with a hint of pink), anthers brown; style white, slightly tinted green at base, stigma brown/green. Calyx 1–2mm long, green. Pedicels up to 25mm long, green, flushed rusty red, sparsely brown scaly. Lvs elliptic to ovate, up to 52 × 25mm, margins flat, more or less glossy and close to moderate olive green (147A) above, acute to shortly acuminate at apex, slightly rounded at base, with pale green scales below which age to brown; petioles c.10mm long, yellow-green, with brown scales. Shrub with a spreading habit, 0.6 × 0.75m in 7 years. Late April to early May.

PC 2009

Nomenclatural standard: herbarium specimen of the award plant (WSY0112427). Etymology: the registrant lives within the Dartmoor National Park (Dartmoor Forest) and has often given fantasy names to his plants (one parent being 'Yaku Fairy')

(r) **'Gaelle des Vignes'**

cv. Elepidote rhododendron: Aladdin Group (s) × Polar Bear Group

H: J.-F. Petton (2001), G: L. Petton (2006), N: J.-F. Petton (2006), REG: J.-F. Petton (2009)

Fls 6–10/lax truss, tubular funnel-shaped, 70 × 70mm, with 7 wavy-edged lobes, buds deep purplish pink (70C), opening inside white, flushed pink at the base of the tube and with inconspicuous pink spots on one lobe, outside white, with a pink base (recorded as light purple, 75B); stamens 14–18, with up to 2 being petaloid in some flowers, filaments white; style hairy, stigma pale greenish yellow; slightly scented. Calyx negligible. Lvs oblong-lanceolate, 180 × 50mm, cuneate to almost truncate at base, acute and apiculate at apex, wavy-edged, hairless. Shrub 1.05 × 1.05m in 8 years. June.

Nomenclatural standard: colour print provided by the registrant (WSY0112747). Etymology: named for Gaelle, the granddaughter of the registrant who lives south of Nantes in an area where she is surrounded by vineyards of Muscadet vines

(r) **'Gerstenkorn'**

cv. Elepidote rhododendron: 'Whitney's Tiger Lily' × ('Cunningham's White' × 'Sappho') × *yakushimanum* 'Koichiro Wada'

H: T. Huisman (1995), G: T. Huisman (1999),

N: T. Huisman (2009), I: T. Schröder (2010),

REG: T. Huisman (2009)

Fls 7–10/ball truss, funnel-shaped, 50–60 × 55–65mm, with 6 wavy-edged lobes, each lobe slightly bilobed, buds vivid reddish orange (33A), opening inside light pinkish white with a touch of cream (155D), some lobes with a flare of pinkish white (N155C); outside opening at first pinkish white (N155C) with strong purplish pink (62A) veins, when fully open coloured like the inside; blotches and spots vivid reddish orange (34A) to moderate red (N34A), spread over all lobes (although mostly on the three dorsal lobes), the blotches breaking up into spots apically: the blotches match cross the lobes (i.e. the left half of one lobe with the right half of the next lobe). Calyx 1–3mm long, pinkish white, with some hairs. Lvs elliptic, 65–105 × 25–40mm, acute at apex, cuneate at base, glossy green above, flat, hairless. Shrub 0.9 × 1.2m in 10 years. Early to mid May.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0108031). Known as seedling no. Z 01–98. Etymology: named for a nurseryman from near Westerstede in Germany

(r) **'Gertrud Welz'**

cv. Elepidote rhododendron: *yakushimanum* (s) × *forrestii* Repens Group

H: H.-J. Danneil (1978), G: H.-J. Danneil (1985),

N: H.-J. Danneil, REG: H.-J. Danneil (2009)

Fls 3–4/loose, open truss, subtended by large pale bracts, funnel-campanulate, 40 × 30mm, with 5 slightly wavy-edged lobes, red in bud, opening blood red throughout, with inconspicuous spots which are only a little darker than the ground colour. Calyx short and very pale. Lvs elliptic to obovate, 30–40(–60) × 10–15mm, obtuse and mucronate at apex, cuneate at base, margins flat, dark glossy green above, hairless. Shrub 0.4 × 0.8m in 30 years. Early April.

A plant exhibiting most of the characters of its pollen parent. Nomenclatural standard: colour print provided by the registrant (WSY0112393)

(r) **'Girlspower'**

cv. Elepidote rhododendron: 'Fantastica' (s) × 'Double Eagle'

H: H. Hesse (1999), G: H. Hesse (2005),

N: H. Hesse, REG: H. Hesse (2010)

Fls 6–10/ loose, rounded truss, broadly funnel-shaped, 35–47 × 40–70mm, with 5 flat-edged lobes, fire-red in bud, opening bright red throughout, with a strong red-brown marking at the centre of the dorsal lobe and dark red spots on the apical part of the dorsal lobe. Calyx 1–3mm long, pale green. Lvs elliptic, 55–105 × 22–50mm, rounded at apex, rounded/partly narrowed at base, margins flat, matt green above, hairless. Very compact shrub, 0.75 × 0.8m in 10 years. Mid May.

(r) **'Gletschereis'**

cv. Elepidote rhododendron: 'Queen Souriya' (s) × 'Blanka'

H: K.-H. Hübbers (1988), G: K.-H. Hübbers

(1994), N: K.-H. Hübbers (2009), REG: J. Westhoff (2009)

Fls 16/roundish, flattened, loose truss, funnel-shaped, 60 × 90mm, with 7 wavy-edged lobes, buds moderate purplish red to strong reddish purple (70A–B), opening inside nearly white and blushed red-purple, outside white and blushed red-purple on the dorsal part, with dense, greyish red (184A) spots, like a blotch, deep in the dorsal part of the throat, which coalesce into two bands. Calyx 3mm long, translucent pink. Lvs ovate, 90 × 60mm, wedge-shaped with a minute point at apex, truncate to slightly rounded at base, margins flat to slightly upcurved, slightly glossy above, hairless. Shrub 1.8 × 1.8m in 20 years. Early May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0104984). Etymology: the epithet means glacial ice

(r) **'Golden Mist'**

cv. Elepidote rhododendron: 'Percy Wiseman' (s) × 'Windwalker'

H: J. Barlup (2003), G: J. Barlup (2007), N: J. Barlup (2010), REG: J. Barlup (2010)

Fls 18/ball truss, saucer-shaped, 45 × 75mm, with 5 wavy-edged lobes, deep pink (48A) in bud, opening inside light yellow (10C), shading to pale greenish yellow (10D) at margins of all lobes, outside pale greenish yellow (10D), with a strong greenish yellow (153B) flare on the dorsal lobe extending c.30mm from the base. Calyx c.10mm long, light yellow (10C). Lvs elliptic, 140 × 55mm, broadly acute at apex, rounded at base, margins flat, glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.8m in 6 years. Mid May.

(r) **'Goldstückchen'**

cv. Elepidote rhododendron: 'Marzipan' (s) × 'Jalisco Eclipse'

H: K.-H. Hübbers (1990), G: K.-H. Hübbers (1996), N: K.-H. Hübbers (2009), REG: J. Westhoff (2009)

Fls 8/rounded, flat-topped truss, funnel-shaped, 35 × 50mm, with 5 wavy-edged lobes, inside brilliant to pale greenish yellow (9C–D), outside brilliant to light yellow (10A–B), with a deep red (53A) blotch in the dorsal throat, breaking up into a few spots; the colour non-fading in the sun. Calyx 2–3mm long, green, with a red rim. Lvs broadly elliptic, 75 × 40mm, obtuse and minutely apiculate at apex, truncate at base, margins flat, matt above, hairless. Shrub 1.0 × 1.0m in 10 years. Late May to early June. Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0104993). Etymology: named after a type of wine, the word meaning a little piece of gold

(r) **'Gordon's Trophy'**

cv. Elepidote rhododendron: *yakushimanum* (s) × un-named Lem hybrid

H: G. Heyduck (1972), G: G. Heyduck (1977), N: C. Smith (1982), I: D. Smart, REG: D. Smart (2010)

Fls 14/truss, funnel-campanulate, 50 × 65m, with 5 wavy-edged lobes, opening inside pale orange-yellow (19D), with the dorsal lobes pale yellow (19C) and

all blending to pale purplish pink (56B)

approximately 5mm from the edge; outside pale orange-yellow (19D), with pale purplish pink (56B) midribs; moderate orange-yellow (164C) spotting on the dorsal lobe, extending c.40mm from the base; flower colour fades to creamy white, with pink highlights. Calyx irregular, up to c.30mm long, pale orange-yellow (19D). Lvs elliptic, c.105mm × 25mm, broadly acute at apex, cuneate at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairy below (hairs greyish yellow-green (148D) at first, aging to greyish yellow-green (195B); petioles hairy. Shrub 0.9 × 0.9m in 10 years. Late April.

(r) **'Grace Sweeney'**

cv. Elepidote rhododendron: 'Scintillation' (s) × unknown

H: D.A. Griswold (1969), G: Griswold Nursery (1980), N: Griswold Nursery (2010), REG: Griswold Nursery (2010)

Fls 14/conical truss, broadly funnel-shaped, 50 × 65mm, with 5 very wavy-edged lobes, deep purplish pink (73A) in bud, opening inside pale purplish pink (62D) at the margins, shading paler towards the centre and with sparse, slightly darker pink spots on the dorsal lobes; outside strong purplish pink (55B). Lvs oblong, 90 × 45mm, broadly acute at apex, cuneate at base, margins downcurved, dull and moderate olive green (137A) above, hairless. Shrub with a dense habit, 1.5 × 1.4m in 41 years. Early June.

Etymology: named for the mother of Greg and Tony Sweeney, the former a member of the Olympic Chapter of the American Rhododendron Society

(r) **'Graeme Eaton'**

cv. Elepidote rhododendron: 'Countess of Lockington' (s) × 'Australian Sunset'

H: D.J. Dosser (2000), G: D.J. Dosser, N: D.J. Dosser, REG: D.J. Dosser (2009)

Fls 8–10/ball truss, funnel-shaped, 40 × 100mm, with 5 wavy-edged lobes, buds strong red (50A), opening inside light yellowish pink (29C) outside deep pink (50B); not marked; slightly scented. Lvs oblong-elliptic, 100 × 40mm, broadly acute at apex,

cuneate at base, margins flat, matt green above, hairless. Shrub 1.5 × 1.0m in 9 years. November (Australia).

Etymology: named for a past Australian *Rhododendron* cultivar registrar

(v) **'Graham Snell'**

cv. *Lepidote* rhododendron: 'Gold Bullion' (s) × 'Lemon Spice'

H: G.L.S. Snell, G: K. Hagan (2008), N: K. Hagan (2010), REG: K. Hagan (2010)

Fls 7/loose truss, tubular funnel-shaped, 60 × 80mm, with 5 flat-edged lobes, inside brilliant yellow (11A), outside vivid yellow (13A) with the base flushed brilliant yellow-green (150A), unmarked. Lvs elliptic, 95 × 45mm, acute at apex, cuneate at base, margins flat, with no visible indumentum. Shrub 1.0 × 0.45m. Early spring. Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the well-known breeder of *vireya* rhododendrons

(r) **'Gretzelhöschen'**

cv. *Elepidote* rhododendron: 'Gretzel' (s) × unknown

H: K.-H. Hübbers (1984), G: K.-H. Hübbers (1993), N: K.-H. Hübbers (2009), REG: J. Westhoff (2009)

Fls c.6/flat truss, funnel-shaped, 60 × 80mm, with 5 wavy-edged lobes, strong red (53C) in bud, opening inside strong to light purplish pink (55B–C), outside deep to strong purplish pink (55A–B), with dark red (187C) spots on the dorsal lobe; slightly scented. Calyx 35mm long, coloured like the corolla, with the same spotting. Lvs lanceolate, 80 × 50mm, cuneate at the apex, truncate at the base, margins slightly decurved, glossy above, with a few, light brown, woolly hairs on the mature foliage. Small, slow-growing shrub, 1.1 × 1.1m in 20 years. Late April to early May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0104982). Etymology: based on the name of the seed parent (a variation on the female name Gretel) and a word play on the petaloid (i.e. hose-in-hose type) calyx and the German meaning of the word "Höschen" (small trousers/underpants for women)

(r) **'Gwendolyn Wright'**

cv. *Elepidote* rhododendron: 'Ring of Fire' (s) × 'Courtenay Queen'

H: H.R. Wright (2001), G: H.R. Wright (2006), N: H.R. Wright (2009), REG: H.R. Wright (2009)

Fls 10/domed truss, saucer-shaped, 40 × 90mm, with 6–7 wavy-edged lobes, dark red to deep purplish pink in bud, opening inside with a yellowish green centre and paler greenish yellow arms extending along the middle of each lobe to the light to moderate pink borders surrounding each lobe; a network of pale veins is visible within most of the corolla. Calyx c.5mm long. Lvs oblong, 120–150 × 40–60mm, semi-glossy and mid green above, hairless. Shrub 1.0 × 0.75m in 4 years. Mid May. Etymology: named for the wife of the registrant

(v) **'Gypsy Bell'**

cv. *Lepidote* rhododendron: *pauciflorum* (s) × *womersleyi*

H: A. Rouse (1998), G: A. Rouse (2005), N: A. Rouse (2009), REG: A. Rouse (2009)

Fls pendent, 2–3/truss, slightly curved tubular-campanulate, tube 15–20mm long, lobes 6–7mm long, corolla 20mm wide, with 5 flat-edged lobes, strong red (45D) inside and out, unmarked. Lvs elliptic to obovate or rounded, 12–20 × 6–12mm, margins flat to slightly decurved, glossy above, obtuse to rounded at apex, broadly tapered to rounded at base, scaly. Shrub 0.3 × 0.6m in 11 years, forming a wiry, multi-branched, spreading bush. October (i.e. spring in Australia) and occasionally in autumn.

Nomenclatural standard: colour print provided by the registrant (WSY0112398)

(r) **'Hannah Grace'**

cv. *Elepidote* rhododendron: 'Tom Everett' (s) × 'Fashion Plate'

H: T. Ahern (1995), G: T. Ahern (2004), N: T. Ahern (2009), REG: T. Ahern (2009)

Fls 10/ball truss, broadly funnel-shaped, c.60 × 100mm, with 7 flat-edged lobes, buds light pink, edged yellow, opening inside deep yellow in the centre and shading to light yellow halfway along the lobes, the edges tinged with light pink; outside light

yellow, with pink edges; unmarked; moderately scented. Lvs obovate, 140 × 50mm, obtuse at apex, cuneate at base, margins flat, dull and mid green above, hairless. Shrub 1.4 × 1.2m in 13 years. Mid May.

Etymology: named for one of the registrant's grandchildren

(v) **'Hawaii Sunrise'**

cv. Lepidote rhododendron: (*macgregoriae* × *laetum*) × *saxifragoides*, F<sub>2</sub> (s) × (*macgregoriae* × *laetum*) × *saxifragoides*, F<sub>2</sub>

H: E.W. Moyles (1999), G: R. Marques & S. Bertelmann (2003), N: R. Marques & S. Bertelmann (2008), I: Pacific Island Nursery (2008), REG: R.F. Kruppa (2009)

Fls 6–9/flat truss, tubular funnel-shaped, 45 × c.50mm, with 5 flat-edged lobes, moderate purplish red (64A) in bud, opening inside brilliant greenish yellow (4A) at the base of the tube, blending into pale purplish pink (56B) before reaching the margins; outside light greenish yellow (3C) at the base of the tube, and changing gradually to orange-yellow before reaching the margins; unmarked; stigma and style light green; filaments yellow, anthers yellowish brown. Lvs elliptic, 85–90 × 20–35mm, either broadly acute or acuminate at apex, cuneate at base, margins upcurved, glossy and moderate olive green (137B) above, with no visible indumentum. Shrub 0.5–0.8 × 0.45–0.5m in 5 years. Flowering three or four times a year between May/June and November/December.

(v) **'Heart of Pineapple'**

cv. Lepidote rhododendron: (*macgregoriae* × *laetum*) × *saxifragoides*, F<sub>2</sub> (s) × (*macgregoriae* × *laetum*) × *saxifragoides*, F<sub>2</sub>

H: E.W. Moyles (1999), G: R. Marques & S. Bertelmann (2004), N: R. Marques & S. Bertelmann (2009), I: Pacific Island Nursery (2008), REG: R.F. Kruppa (2009)

Fls 4–5/flat truss, tubular funnel-shaped, 50 × c.20–30mm, with 5 flat-edged lobes, strong yellow-green (145A) in bud, opening inside and out light greenish yellow (8B) at the base, fading to much lighter pale greenish yellow (9D) at the

margins, with a barely discernible pink tinge on the lobes; a fine green line extending from the base to halfway up the lobes; stigma and style green; filaments yellow, anthers brownish yellow. Lvs elliptic, 40–45 × c.20mm, acuminate at apex, cuneate at base, margins upcurved, glossy and moderate olive green (137B) above, with no visible indumentum. Shrub with a dense habit, 0.5–0.8 × 0.45–0.5m in 5 years. Flowering two to four times a year between May/June and November/December.

(r) **'Heol Breizh'**

cv. Elepidote rhododendron: 'Karen Triplet' (s) × 'Nelda Peach'

H: L. Le Duigou (2003), G: A. Le Duigou (2009), N: L. & A. Le Duigou (2010), REG: L. & A. Le Duigou (2010)

Fls 15–17/domed truss, broadly funnel-shaped, 65 × 100mm, with 5–7 flat-edged lobes, buds moderate red (47A), opening inside a mixture of pale greenish yellow (10D), light yellow (11B) and light greenish yellow (8B), flushed pale purplish pink around the margins, with basal nectar pouches moderate purplish red (58A); outside light greenish yellow (8B), with deep pink (48B) margins; inconspicuous deep greenish yellow (153A) spots on the dorsal lobe; stamens 12–14; style pale yellow-green (149D), stigma strong yellow-green (144A). Calyx partially petaloid, 15–40mm long, light greenish yellow (8B), with a strong red (53B) basal blotch, margins tinged pink. Lvs oblong, 140 × 45mm, rounded at base, cuspidate at apex, margins flat to decurved, matt green and bullate above, hairless; petioles moderate red (47A). Compact shrub 0.7 × 0.5m in 6 years.

May to June.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: the epithet is Breton for "Sun of Brittany" and refers to the flower colour and the area in which it was raised

(r) **'Hesse's Biggi'**

cv. Elepidote rhododendron: 'Mrs J.G. Millais' (s) × Elizabeth de Rothschild Group

H: H. Hesse (1996), G: H. Hesse (2003), N: H. Hesse, REG: H. Hesse (2010)

Fls 15–17/ slightly loose, rounded truss, open

funnel-shaped, 44–65 × 70–100mm, with 5–7 wavy-edged and reflexed lobes, creamy white in bud, opening creamy white throughout, with red and partially brownish yellow markings on the dorsal lobe. Calyx 1–2mm long, green or reddish. Lvs elliptic, 50–120 × 26–50mm, acute at apex, rounded at base, matt green above, margins flat, hairless. Compact shrub, 1.2 × 1.3m in 14 years. Early May.

Etymology: Biggi is the nickname of the registrant's wife, Brigitte

(r) **'Hesse's Nerisha'**

cv. Elepidote rhododendron: 'Peste's Fire Light' (s) × 'Hachmann's Kabarett'  
H: H. Hesse (1999), G: H. Hesse (2007),  
N: H. Hesse (2009), REG: H. Hesse (2009)  
Fls 10–16/full, spherical truss, open funnel-shaped, 40–53 × 45–75mm, with 5 wavy-edged lobes, inside white, with a striking dark red blotch on the three dorsal lobes, the other lobes with many red dots, outside whitish pink. Calyx 3–5mm long. Lvs elliptic, 53–125 × 20–50mm, margins flat, acute/partly rounded at apex, rounded at base, matt green above, hairless. Shrub 0.7 × 0.4m in 10 years. Mid to late May..

(v) **'Highland Bonsai'**

cv. Lepidote rhododendron: *luraluense* (s) × *womersleyi*  
H: A. Rouse (2001), G: A. Rouse (2005),  
N: A. Rouse (2009), REG: A. Rouse (2009)  
Fls pendent, 2–3/truss, tubular-campanulate, tube 20mm long, lobes 12mm long, corolla 30mm wide, with 5 slightly and narrowly wavy-edged lobes, strong yellowish pink (43D) inside and out. Lvs elliptic, 15–20 × 7–10mm, margins slightly decurved, glossy green above, obtuse to acute at apex, tapered at base, scaly. Shrub 0.5 × 0.6m in 8 years, forming a wiry, multi-branched plant. October (Australia).  
Nomenclatural standard: colour print provided by the registrant (WSY0112402). Etymology: one of the parents is from the highlands of Papua New Guinea, and the plant has potential as a bonsai subject

(r) **'Himmelberg'**

cv. Elepidote rhododendron: *keiskei* 'Yaku Fairy' (s) × *glaucophyllum*, white-flowered  
H: W. Reich (1999), G: W. Reich (2004),  
N: W. Reich (2009), I: Glendoick Gardens (2010),  
REG: W. Reich (2010)  
Fls 5–6/loose truss, open funnel-shaped, 34–36 × 48–50mm, with 5 very wavy-edged lobes, buds pale yellow to yellowish white (158B–C), opening throughout pale yellow to yellowish white (158B–C) and fading to white after two days, un-marked. Calyx 6–7mm long, brownish orange (165B) to light olive brown (199B). Lvs ovate to lanceolate, 7–67 × 5–24mm, pointed at apex, cuneate at base, margins flat to slightly decurved, matt to semi-glossy and moderate olive green (137A–B) above, with a scaly indumentum. Shrub 0.3 × 0.5m in 7–8 years. Early May.  
Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for one of the hills near where the registrant lives in Alfeld, Germany

(r) **'Holden's Solar Flair'**

cv. Elepidote rhododendron: un-named yellow-flowered D.G. Leach hybrid (no. 67-LL-6) × 'Casanova' (s) × (((('Powell Glass' × (*fortunei* subsp. *discolor* × *Fabia* Group))) × ('Russell Harmon' × 'Goldsworth Orange')) × (('Russell Harmon' × 'Goldsworth Orange') × 'Calcutta'))  
H: D.G. Leach (1973), G: D.G. Leach (1982),  
N: S.L. Krebbs (2008), I: Briggs Nursery (2010),  
REG: S.L. Krebbs (2009)  
Fls 12–14/incomplete domed truss, broadly funnel-shaped, 45 × 55mm, with 5 flat-edged lobes, of waxy texture, moderate pink (35D) in bud, opening inside pale yellow (8D), with a strong flare consisting of a deep red (53A) blotch surmounted with spots of the same colour on the dorsal lobe; outside pale yellow (8D); bud scales red. Lvs elliptic, 110 × 40mm, broadly acute at apex, rounded at base, margins flat, glossy and moderate olive green (147A) above, hairless. Shrub 2.1 × 2.4m in 26 years. Midseason.  
The spelling of "Flair" as such, not "Flare", is deliberate

(r) **'Holden's Spring Herald'**

cv. Elepidote rhododendron: 'Catalga' (s) × *rex* subsp. *fictolacteum*  
H: D.G. Leach (1975), G: D.G. Leach (1988),  
N: S.L. Krebs (2009), I: Briggs Nursery (2012),  
REG: S.L. Krebs (2009)  
Fls 13–14/domed truss, funnel-shaped, 50 × 65mm, with 5 flat-edged lobes, light purple (75B) in bud, opening inside very pale purple (69D) and fading to white, with a deep purplish red (61A) blotch and a spotted flare in the throat and on the dorsal lobe; outside very pale purple (59C), fading to white. Lvs elliptic, 190 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (137A) above, with moderate orange-yellow (164C) hairs below. Shrub 5.5 × 3.7m in 34 years. Late April.

(r) **'Honey Creek'**

cv. Elepidote rhododendron: *hyperythrum* (s) × 'Princess Elizabeth'  
H: J.T. Thornton (1993), G: L.O. Miller (2003),  
N: L.O. Miller (2003), I: Briggs Nursery,  
REG: L.O. Miller (2009)  
Fls 11/ball truss, broadly funnel-shaped, 40 × 50mm, with 5 wavy-edged lobes, red in bud, opening inside pink, shading lighter in the throat, the dorsal lobe with maroon stippling, outside opening a deeper shade of pink. Lvs oblong, 125 × 50mm, acute at apex, rounded at base, margins downcurved, dark green above, hairless. Dense shrub, 1.8 × 2.4m in 12 years. Mid to late April. Etymology: named for an arm of the Grand Lake in Oklahoma State, USA, which is near the property of the registrant where this rhododendron is grown

(r) **'Honeydrops'**

cv. Lepidote rhododendron: 'Laurin' (s) × *cinnabarinum* 'Nepal'  
H: W. Reich (1994), G: W. Reich (2003),  
N: W. Reich (2005), REG: W. Reich (2010)  
Fls 7–8/open truss, pendulous, tubular-campanulate, 37–38 × 30–32mm, with 5 flat-edged lobes, buds strong pink (48C) to moderate purplish pink (66D), opening inside light orange-yellow (19B), outside light orange-yellow (19B) at the

margins shading to strong pink (48C) and moderate purplish pink (66D) at the base, aging to moderate to light purplish pink (65A–B); unmarked. Calyx 4.5mm long, yellowish green. Pedicels yellowish green. Lvs elliptic, 25–57 × 16–25mm, broadly acute at apex, cuneate at base, margins flat, glossy green above, with a scaly indumentum. Shrub 0.75–0.8 × 0.55m in 12 years, remarkably dwarf for a *cinnabarinum* hybrid. Mid to late May. Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: suggested by the partially golden colour of the hanging flowers

(r) **'Horai-asahi'**

cv. Elepidote rhododendron: wild origin selection from *makinoi*  
S: K. Ohta (2007), N: K. Ohta (2010),  
REG: K. Ohta (2010)  
Fls 6–10/half ball truss, broadly funnel-campanulate, 40 × 60mm, with 5 more or less smooth-edged lobes, deep pink in bud, opening pinkish white both inside and out, with fine, red, dorsal spotting. Calyx 3–4mm long, light brown. Lvs linear, 120 × 15mm, margins recurved, acuminate at base and apex, glossy green above, with a white indumentum below when young, turning brown with age and easily rubbed off. Shrub 2.5 × 2.0m in c.80 years. Late April. Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: "Horai" is the name of the Japanese mountain on which this plant was found, "asahi" means sunrise

(r) **'Ice Prism'**

cv. Elepidote rhododendron: *bureauvii* × 'Ken Janeck' (s) × 'One Thousand Butterflies'  
H: J. Barlup (1998), G: J. Barlup (2003), N: J. Barlup (2009), I: Dover Nursery (2009), REG: J. Barlup (2009)  
Fls 16/ball truss, broadly funnel-shaped, 45 × 75mm, with 7 very wavy-edged lobes, strong purplish red (60C) in bud, opening inside white (155D), lightly tinged a creamier shade of yellowish white (158C) on the dorsal lobes, outside white (155D); with a strong red (53C), speckled dorsal flare extending 35mm over-all from the base and

appearing to be almost solid colour within c.20mm of the centre. Calyx 25mm long, white (155D), with a strong red (53C) flare covering most of each lobe. Lvs elliptic, 150 × 75mm, broadly acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.2 × 1.5m in 10 years. Mid April.

(r) **'Jack Morgan'**

cv. *Elepidote rhododendron*: 'Pink Prelude' (s) × 'Cameo'

H: Mrs E.M. Watson (1992), G: Mrs E.M. Watson, N: Mrs E.M. Watson (2007), REG: Mrs E.M. Watson (2009)

Fls 17/ball truss, funnel-campanulate, 75 × 100mm, with 7 very wavy-edged lobes, moderate pink in bud, opening inside and out light pink, fading to nearly white; dark red brush-like markings in the throat below the dorsal lobes; heavy flower substance. Lvs elliptic, 170 × 70mm, acuminate at apex, oblique at base, margins flat, dull and moderate yellow-green (146B–C) above, hairless. Shrub with a dense habit, 1.6 × 1.8m in 15 years. March to April.

Etymology: named for a friend of the registrant who has been helping to tend her garden

(v) **'Jean Curtis'**

cv. *Lepidote rhododendron*: 'Anacapa' (s) × 'Clare Elsie'

H: W.A. Moynier (2000), G: S. Bertelmann and R. Marques (2004), N: S. Bertelmann & R. Marques (2009), I: Pacific Island Nursery (2009), REG: R.F. Kruppa (2010)

Fls 9–12/domed truss, elongated tubular funnel-shaped, 90 × 90–100mm, with 5–6 flat-edged lobes, dark reddish purple in bud, opening deep purplish pink (58D), edged strong purplish red (58B), with the tube becoming lighter to pale purplish pink (56C) and finally yellowish pink at the base; outside similar except that the base becomes yellowish green; unmarked; style red to very dark pink, stigma dark red; filaments pink, merging into cream, anthers dark red. Lvs elliptic, 165 × 70mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and moderate yellow-green (137A) above, strong to moderate yellow-green

(144A–146B) below, with no visible indumentum. Dense, upright shrub, 0.9 × 0.6m in 6–8 years. May to June and January to February.

Etymology: named for the former president of the Panaewa Zoo and Vireya Gardens Association in Hilo, Hawaii

(r) **'Joachim Reich'**

cv. *Lepidote rhododendron*: 'Gletschernacht' (s) × 'April Rose'

H: W. Reich (1995), G: W. Reich (2001), N: W. Reich (2007), REG: W. Reich (2010)

Fls double, with no normal stamens, up to 7/open truss, open campanulate, 22–29 × 30–36mm, light to very pale purple (84C–D) to very light to very pale purple (85C–D), base of throat and style strong to light purplish pink (73B–C). Calyx 2mm long. Lvs elliptic to broadly elliptic, 15–30 × 8–15mm, rounded (to rarely pointed) at apex, broadly cuneate at base, margins flat, glossy green above; some leaves deciduous, becoming wine-red to reddish brown. Shrub 0.6 × 0.4m in 10 years. April.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the deceased younger brother of the registrant

(r) **'Joe Becales'**

cv. *Elepidote rhododendron*: 'Janet Blair' (s) × 'Goldsworth Orange'

H: J. Becales, G: J. Becales, N: D. Mac Mullan (2009), REG: D. Mac Mullan (2009)

Fls 9/domed truss, funnel-campanulate, 60 × 75mm, with 5 wavy-edged lobes, moderate pink tinged yellowish in bud, opening inside light yellow/orange at the ovary and shading through peach pink to very pale pink at the margins, with orange speckles in the throat and extending along the dorsal lobe about halfway to the margin; outside medium pinkish yellow, with light pink and cream at the margins. Calyx c.6mm long, light to medium yellow. Lvs elliptic, 90 × 40mm, broadly acute at apex, oblique at base, margins flat, semi-glossy and mid green above, hairless. Shrub 1.8 × 1.2m in 10 years. Late May to early June.

Etymology: named in memory of a friend who had a fine garden that included many of his own hybrid rhododendrons

(r) **'John Addison Shiel'**

cv. Elepidote rhododendron: 'Crest' (s) × 'Odee Wright'

H: D.A. Griswold (1969), G: Griswold Nursery (1980), N: Griswold Nursery (2010), I: Griswold Nursery (2010), REG: Griswold Nursery (2010)  
Fls 13/lax truss, open funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, vivid reddish orange (33A) in bud, opening inside light yellowish pink (29C), with randomly distributed pale pink markings on the lobes; outside light yellow (11B); anthers dark reddish brown; stigma bright green; ovary bright green. Lvs elliptic, 75 × 40mm, broadly acute at apex, cuneate at base, margins flat, dull and moderate yellowish green (137D) above, hairless. Shrub 1.8 × 2.1m in 41 years. Mid May.

Etymology: named for the son-in-law of Orine Holm, part owner of the Griswold Nursery

(r) **'Jonathan Coker'**

cv. Elepidote rhododendron: 'Lemon Lodge' (s) × unknown

H: B.J. & H. Coker (c.1994), G: B.J. & H. Coker (c.1999), N: B.J. & H. Coker (2010), REG: B.J. & H. Coker (2010)  
Fls 15/upright, rounded truss, open funnel-shaped, 45 × 45mm, with 7 wavy-edged lobes, buds brilliant yellow-green (154C), opening inside light yellow-green (154D) with brilliant yellow-green (154B) in the throat, outside light yellow-green (154D); very light dorsal spotting of brilliant yellow-green (154B). Lvs elliptic, 135 × 55mm, cuspidate at apex, rounded at base, margins flat, matt green above, hairless. Shrub 2.5 × 1.2m in 16 years. Midseason. Nomenclatural standard: colour print provided by the registrants (WSY). Etymology: named for the son of the registrants

(r) **'Jordan's Arabesque'**

cv. Elepidote rhododendron: parentage unknown  
G: J. van de Sande (2000), N: J. van de Sande, REG: J. van de Sande (2009)  
Fls 13/conical truss, open funnel-shaped, 55 × 90mm, with 6 wavy-edged lobes, deep purplish pink (66C) in bud, opening inside off-white, tinged very pale purplish pink and shading to a broad margin of

moderate purplish pink (68C), the centre pale yellow-green (4D), outside light purplish pink (68D); unmarked. Calyx c.1.5mm long, strong yellowish green (141C). Lvs oblong, 130 × 65mm, obtuse at apex, oblique at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.4 × 1.4m in 10 years. Mid May.

Selected by the registrant as an unbloomed seedling from a group donated at a meeting of hybridisers. Etymology: named for the registrant's daughter, Jordan, who is a dancer (and hence also the reference to the classic dance position, an arabesque)

(r) **'Julie N. Firth'**

cv. Elepidote rhododendron: 'Scintillation' (s) × 'Lem's Cameo'

H: D.A. Griswold (1969), G: Griswold Nursery (1980), N: Griswold Nursery (2010), REG: Griswold Nursery (2010)  
Fls 6/conical truss, open funnel-shaped, c.65 × c.75mm, with 5 wavy-edged lobes, strong purplish pink (62A) in bud, opening pale greenish yellow (2D) throughout, tinged pink and with moderate reddish orange (35B) spotting on the three dorsal lobes; outside strong purplish pink (62A) at the margins. Calyx c.6mm long, light yellowish pink (19B). Lvs oblong, 125 × 50mm, obtuse at apex, cuneate at base, dull and moderate yellow-green (148A) above, hairless. Shrub 1.2 × 2.4m in 41 years. Mid May.

Etymology: named for a long-time friend of the Griswold Nursery

(r) **'June Gardiner'**

cv. Elepidote rhododendron: ('Lem's Cameo' × double form of *griffithianum*) seedling no.1 (s) × ('Lem's Cameo' × double form of *griffithianum*) seedling no.10

H: Mrs S.E. Davies (1991), G: Mrs S.E. Davies (1996), N: Mrs S.E. Davies (2007), REG: Mrs S.E. Davies (2010)

Fls semi-double, 10–13/loose rounded truss, broadly campanulate, 40–50 × 80–95mm, the outer corolla with 6–7 wavy-edged lobes, with 1–5 additional lobes internally, buds light yellowish pink (36B)

with a strong purplish red (54B) dorsal stripe, opening inside white with a stripe of pale pink (49D) down the central one third of each lobe which fades with age to almost white, outside white with a similar stripe but of deep purplish pink (54B) and fading to moderate purplish pink (54D); otherwise unmarked; the extra inner corolla lobes coloured as the main corolla and the overall effect is of a striped corolla at first aging to almost white; stamens 11–19; stigma vivid red (45A); scented. Calyx 5mm long, brilliant greenish yellow (151D). Lvs oblong, 85–120 × 30–47mm, margins wavy, cuneate at apex and base, hairless. Shrub 1.7 × 1.5m in 19 years. October (New Zealand).

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the registrant's mother

(r) '**June Wedding**'

cv. Elepidote rhododendron: parentage unknown – seed collected from RHS Garden, Wisley, UK  
G: C.L. Justice (1985), N: C.L. Justice (2009),  
REG: D. Justice (2010)

Fls 5–7/domed truss, broadly funnel-shaped, 70 × 90mm, with 7 wavy-edged lobes, greenish white in bud, opening white inside and out, with a central blotch of a yellowish green colour (close to brilliant yellow-green, 142B), marked in the quadrant between 10 and 2 o'clock with an orange-like colour between approximately strong orange-yellow (24B) and vivid reddish orange (34B). Calyx 1mm long, green. Lvs oblong, 200 × 70mm, broadly acute at apex, minutely cordate at base, margins downcurved, semi-glossy and mid green above, hairless. Shrub with an open habit, 5 × 3m in 45 years. Late June.

Said to resemble *decorum* subsp. *cordatum*, but with longer petioles

(r) '**June Yellow**'

cv. Lepidote rhododendron: *lepidostylum* × *keiskei*  
'Yaku Fairy'

H: W. Reich (1988), G: W. Reich (1993),

N: K.N.E. Cox (c.2002), I: Glendoick Gardens (2003), REG: W. Reich (2010)

Fls 4–5/truss, open tubular to open funnel-shaped,

with 5 wavy-edged lobes, greenish yellow in bud, opening light yellow-green (2C) throughout. Lvs broadly elliptic, 20–40 × 14–24mm, pointed at apex, margins flat to upcurved, glossy green above, with a scaly indumentum; petioles very short. Shrub with a dense habit, 30–35 × 60–70cm in 12 years. Late May to early June.

Nomenclatural standard: colour print provided by registrant (WSY). Etymology: name suggested by flower colour and flowering time

(r) '**Kaitlyn's Delight**'

cv. Elepidote rhododendron: parentage unknown

S: Corneliussen's Nursery, G: Corneliussen's Nursery, N: Corneliussen's Nursery (2009),

I: Corneliussen's Nursery (2009),

REG: Corneliussen's Nursery (2009)

Fls 9/flat truss, campanulate, 45 × 65mm, with 5 wavy-edged lobes, strong red (46A) in bud, opening inside moderate yellowish pink (38B) with a light orange-yellow (22C) centre which fades to light orange-yellow (22D), prominent vivid red (45B) dorsal spotting and many fewer more subdued spots on the remaining lobes; outside deep yellowish pink (39B), fading to strong yellowish pink (35C); stigma bright green. Calyx 40mm long, inside light orange-yellow (22C), with vivid red (45A) dorsal spotting, outside deep yellowish pink (39B), with vivid greenish yellow stripes: the large calyx gives the appearance of a double flower. Lvs elliptic, 90 × 40mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and dark bluish green (131A) above, hairless. Shrub 0.6 × 0.9m in 7 years. Mid May.

Etymology: named for the Corneliussen granddaughter who has wavy red hair and freckles.

(r) '**Kari's Surprise**'

cv. Elepidote rhododendron: 'Fantastica' (s) × 'Snow Candle'

H: J. Barlup (1997), G: K.R. Grov (2009),

N: P.M. Jørgensen (2010), REG: P.M. Jørgensen (2010)

Fls c.10/flat truss, open funnel-shaped, 50–60 × 50mm, with 5 wavy-edged outer lobes and with petaloid stamens, reddish in bud, opening pinkish

rose throughout; unmarked. Calyx 2mm long, green. Lvs lanceolate, 100 × 30mm, tapering at apex, pointed at base, margins decurved, glossy green above, with a brown indumentum below when young, aging to beige. Shrub 0.5 × 0.5m in 10 years. Late May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the grower, who was greatly surprised when it first flowered

(r) **'Kayla Joy'**

cv. Elepidote rhododendron: 'Tom Everett' (s) × 'Fashion Plate'

H: T. Ahern (1995), G: T. Ahern (2009), N: T. Ahern (2009), REG: T. Ahern (2009)  
Fls 11/ball truss, broadly funnel-shaped, c.60 × 100mm, with 7 flat-edged lobes, mauve in bud, opening inside light yellow, outside light yellow, tinged pink along the midveins; throat mid green, with light green spots on the dorsal lobe extending from the base to about halfway to the margin; strongly scented. Lvs elliptic, 150 × 45mm, broadly acute at apex, rounded at base, margins flat, hairless. Shrub 1.5 × 1.2m in 13 years. Mid May.

Etymology: named for one of the registrant's grandchildren

(r) **'Kera'**

cv. Elepidote rhododendron: 'Maverick' (s) × 'Tia'  
H: J. Barlup (2001), G: J. Barlup (2005), N: J. Barlup (2008), REG: J. Barlup (2009)  
Fls 11/ball truss, broadly funnel-shaped, 45 × 70mm, with 5 wavy-edged lobes, strong red (53B) in bud, opening inside white (155D), the dorsal lobe tinged slightly yellowish white (158D) and with all lobes edged light purplish pink (55C) extending inward c.5–6mm; outside similarly coloured; a central basal blotch of moderate red (185B) extends upwards c.15 × c.20mm on dorsal lobes, with more spotting extending another c.6mm; streaks of the same colour extend on all lobes for c.15mm. Lvs elliptic, 70 × 45mm, broadly acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, hairless. Shrub with a dense habit, 0.5 × 0.8m in 8 years. Early May.

(v) **'Kilauea Kid'**

cv. Lepidote rhododendron: (*macgregoriae* × *laetum*) × *saxifragoides*, F<sub>2</sub> (s) × (*macgregoriae* × *laetum*) × *saxifragoides*, F<sub>2</sub>

H: E.W. Moyles (1999), G: R. Marques & S. Bertelmann (2003), N: R. Marques & S. Bertelmann (2008), I: Pacific Island Nursery (2008), REG: R.F. Kruppa (2009)

Fls 3–5/flat truss, tubular funnel-shaped, 50 × 65mm, with 5 flat-edged lobes, strong purplish red (60B) in bud, opening inside vivid reddish orange (32A) at margins, shading to strong yellowish pink and then paler in the throat; outside similarly coloured; unmarked; filaments orange, anthers dark orange; style yellowish green, stigma orange. Lvs elliptic, 55 × c.25mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and dark green (132A) above, with no visible indumentum. Shrub 0.5 × 0.4m in 5 years. Flowering intermittently, for up to three months continually, between May and November/December.

(v) **'Kimo'**

cv. Lepidote rhododendron: *konori* (Jake's form) (s) × *konori* var. *phaeopleplum* × *zoelleri*

H: R.A. Mitchell (2002), G: R.A. Mitchell (2008), N: R.A. Mitchell (2010), REG: R.A. Mitchell (2010)

Fls 8–9/flat truss, elongated tubular funnel-shaped, 80 × 85mm, with 5–7 very wavy-edged lobes, opening inside and out pale yellow-green (157C) to greenish white (157D), with the outside of the tube a barely detectable very light yellow; brownish spots (c.0.8mm wide) at lobe margins; anthers light tan, filaments white; stigma light green, style very light pink; strongly scented. Calyx absent. Lvs of two shapes, either elliptic, 110 × 55mm, acuminate at apex and cuneate at base, or obovate, 110 × 60mm, broadly acute at apex, rounded at base, both semi-glossy and dark yellowish green (139A) above, with no visible indumentum. Shrub with a dense habit, 0.8 × 0.6m in 7 years. January.

Etymology: named for the registrant's son, Jim. Kimo is the Hawaiian equivalent of this name

(r) **'Kirsten Gilbert'**

cv. Elepidote rhododendron: 'Roza Stevenson' (s) × unknown

H: R.J. Gilbert (c.1999), G: R.J. Gilbert, N: R.J. Gilbert (2009), REG: R.J. Gilbert (2009)  
Fls 8–10/open-topped truss, funnel-campanulate, 55 × 80mm, with 5 wavy-edged lobes, buds light yellow (15D), with deep apricot stripes, opening light greenish yellow/pale yellow-green (8C/4D) throughout (the apricot stripes fading as the flower opens), with a small, dark rosy red blotch at the base of the throat; style and filaments pale yellow, stigma green; not scented. Calyx 2mm long, rosy. Lvs narrowly ovate to elliptic, 110 × 40mm, margins decurved, broadly acute or obtuse at apex, narrowly truncate at base, mucronate, hairless. Shrub with a dense, bushy habit, 1.2 × 1.2m in c.10 years. April. Nomenclatural standard: colour print provided by the registrant (WSY0112296). Etymology: named for the registrant's wife

(a) **'Kodai Orange'**

cv. Evergreen azalea: *simsii*, open-pollinated  
G: M.S. Viraraghavan (2008), N: M.S. Viraraghavan (2010), REG: M.S. Viraraghavan (2010)  
Fls 2/truss, broadly funnel-campanulate, 40 × 40mm, with a 5-lobed outer whorl and 3–4 petaloid stamens, lobes wavy-edged, orange-red in bud, opening orange-red throughout, with darker pinkish orange-red dorsal spotting. Calyx 4mm long, greenish brown. Lvs elliptic, 65 × 15mm, acute at both base and apex, margins flat, sparsely hairy. Shrub 1.0 × 1.0m in 10 years. April to May. Etymology: "Kodai" is a shortened form of Kodaikanal, a hill station in the Palni Hills in southern India where the registrant lives and where this plant was raised. Grown from seed collected by Mr Thinakom in 1998 by the Phu River, near Loei, Thailand at 1250m

(a) **'Kodai Ruffles'**

cv. Evergreen azalea: *simsii*, open-pollinated  
G: M.S. Viraraghavan (2008), N: M.S. Viraraghavan (2010), REG: M.S. Viraraghavan (2010)  
Fls double, 2/truss, open funnel-shaped, 30 × 40mm, with a 5-lobed outer whorl and 6–7 large,

petaloid stamens, all lobes very wavy-edged, dark reddish pink in bud and throughout on opening, unmarked. Calyx 15mm long, light green. Lvs elliptic, 60 × 30mm, broadly cuneate at base, narrowly obtuse at apex, glossy green above, apparently slightly toothed, very sparsely hairy. Shrub 1.5 × 0.75m in 10 years. April to May. Etymology: Kodai is an abbreviated form of Kodaikanal, the hill station in the Palni Hills in southern India where the registrant lives and where this plant was raised. Grown from seed collected by Mr Thinakom in 1998 by the Phu River, near Loei, Thailand at 1250m

(r) **'Korrigan'**

cv. Elepidote rhododendron: 'Nancy Evans' (s) × 'Hyperion' (Waterer)  
H: L. Le Duigou (2000), G: A. Le Duigou (2005), N: A. Le Duigou (2009), REG: L. Le Duigou (2009)  
Fls 17/rounded truss, funnel-shaped, 55 × 80mm, with 5 flat to slightly wavy-edged lobes, buds light yellow (18A), with a slight reddish flush around the tips, opening inside pale yellow (18C), with extensive vivid reddish orange (34A) spotting on the dorsal lobe and half of each adjacent lobe; stamens 12, filaments and style pale yellow. Calyx petaloid, 20–35mm long, with reflexed lobes, pale yellow (18C), with conspicuous vivid reddish orange (34A) spotting. Lvs elliptic, 105 × 35mm, margins flat, apiculate at apex, acute at base, hairless. Shrub 0.75 × 0.8m in 8 years. May. Nomenclatural standard: colour print provided by the registrant (WSY0112403). Etymology: Korrigan is a figure in Breton mythology

(r) **'Kranenbells'**

cv. Elepidote rhododendron: 'Barbara Reuthe' (s) × 'New Comet'  
H: K.-H. Hübbers (1989), G: K.-H. Hübbers (1996), N: K.-H. Hübbers (2004), REG: J. Westhoff (2009)  
Fls c.9/very loose truss (the flowers mostly pendent), funnel-shaped, 70 × 90mm, with 7 wavy-edged lobes, inside light yellow (12C), shading to light yellowish pink (38D) at the margins, outside strong yellowish pink (31C), becoming yellow; scented. Calyx 15mm long, yellow, with a small red rim. Lvs

oblong, 120 × 60mm, cuneate with a point at the apex, truncate at the base, margins flat, matt above, hairless. Shrub 1.5 × 1.5m in 10 years. Late April. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112995). Etymology: named for the flower shape and the city in which it was raised (Kranenburg)

(a) **'Kranenelfe'**

cv. Deciduous azalea: 'Anna Pavlova' (s) × 'Silver Slippers'  
H: K.-H. Hübbers (1991), G: K.-H. Hübbers (1997), N: K.-H. Hübbers (2009), REG: J. Westhoff (2009)  
Fls double, c.8/dense, rounded truss, tubular funnel-shaped, 45 × 50mm, with c.10 wavy-edged lobes, striped pink and green in bud, opening inside light greenish yellow (8C), outside white, blushed yellow and pink; scattered vivid yellow (9A) spots; intensely scented. Calyx 5mm long, light green. Lvs elliptic, 70 × 40mm, pointed at apex, tapered at base, margins slightly upcurved, matt above, with a few hairs on the midrib when mature. Shrub 1.0 × 1.1m in 15 years. Early May.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112997). Etymology: named for the city of origin (Kranenburg) and the elfin flowers

(r) **'Kranenglut'**

cv. Elepidote rhododendron: 'Edward Dunn' × *auriculatum* (s) × 'Good News' × 'Erato'  
H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2008), N: K.-H. Hübbers (2010), REG: J. Westhoff (2010)  
Fls 10–12/lax, domed truss, tubular funnel-shaped, 75 × 80mm, with 5 wavy-edged lobes, strong red (53D) in bud, opening inside deep purplish pink (55A), outside strong purplish red (58B), with a dorsal blotch (in the throat and on the lobes) made up of dense spotting of vivid red (45A). Calyx 5mm long, dark red. Lvs oblong, 200 × 50mm, broadly acute at apex, truncate-oblique at base, margins flat to slightly upcurved, slightly glossy green above, hairless; new growth intense coppery red, with a few woolly hairs on the upper surface. Shrub 2.1 × 1.2m in 10 years. Late June.

Etymology: a combination of the city of origin (Kranenburg) and the glowing colour of the flower in sunlight ("glut" means "glow" in German)

(r) **'Kranenrosette'**

cv. Elepidote rhododendron: 'Deerah' (s) × 'Viscy'  
H: K.-H. Hübbers (1993), G: K.-H. Hübbers (2007), N: K.-H. Hübbers (2008), REG: J. Westhoff (2009)  
Fls 10/ rounded, flat-topped truss, funnel-shaped, 40 × 40mm, with 5 wavy-edged lobes, inside white (155C), outside white (155A), with small, deep pink (185D) spots on two-thirds of the dorsal throat. Calyx 3mm long, yellowish green. Lvs lanceolate, in dense, superposed whorls of different sizes, 30 × 10mm and 70 × 20mm (the smallest nearer the apex), cuneate at apex, rounded at base, margins decurved, glossy above, hairless. Shrub 0.7 × 0.5m in 15 years. April.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112994). Etymology: refers to the leaf arrangement and the city in which it was raised (Kranenburg)

(r) **'Kranensonne'**

cv. Elepidote rhododendron: 'Admiral Vanessa' (s) × 'Deerah'  
H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2010), N: K.-H. Hübbers (2010), REG: J. Westhoff (2010)  
Fls c.16/loose conical truss, funnel-shaped, 65 × 90mm, with 6–7 wavy-edged lobes (sometimes with a slit in one of the dorsal lobes), buds strong purplish red (54A), opening inside light yellowish pink (27B) with light purplish pink (55C) on the middle of the lobes, outside moderate pink (49B) with strong purplish pink (55B) on the midveins; very conspicuous, broad blotch of dark red to deep purplish red (187A–D) spots in the throat and on the three dorsal lobes. Calyx 9–20mm long, coloured on the outside like the corolla, on the inside like the blotch. Lvs oblanceolate, 120 × 45mm, broadly acute at apex, oblique at base, margins flat to slightly decurved, glossy green above, hairless. Shrub 1.3 × 0.65m in 9 years. Mid to late June.  
Etymology: named for the city of origin (Kranenburg) and the appearance of the flower like a setting sun ("sonne" means "sun" in German)

(r) **'Lady of the Isles'**

cv. Elepidote rhododendron: Repose Group × Elizabeth de Rothschild Group  
H: E. de Rothschild, G: E. de Rothschild, N: E. de Rothschild, REG: Leopold de Rothschild (2010)  
Fls 9–11/truss, tubular funnel-shaped, 80 × 110mm, with 5–6 wavy-edged lobes, buds pink, opening inside moderate purplish pink (62B), with deep purplish pink (61D) around the edges, outside light purplish pink (73C), shading to a nearly white base; with green spotting on the dorsal lobe; filaments white, anthers brown; style creamy yellow, stigma reddish brown. Calyx 40–45mm long, yellow-green. Lvs oblong, 200 × 75mm, acute at apex, cuneate at base, margins flat, hairless. Shrub 3.0 × 3.0m in 20 years. May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named in honour of the Duchess of Cornwall

(r) **'Laura Shiel'**

cv. Elepidote rhododendron: parentage unknown  
H: D.A. Griswold (1969), G: Griswold Nursery (1980), N: Griswold Nursery (2010), REG: Griswold Nursery (2010)  
Fls 17/ball truss, saucer-shaped to broadly funnel-shaped, 50 × 50mm, with 5 flat-edged lobes, vivid purplish red (61C) in bud, opening strong purplish pink (68B) inside and out, with lighter pink areas on all lobes; unmarked. Lvs oblong, 75 × 40mm, broadly acute at apex, cuneate at base, margins downcurved, dull and dark bluish green (133A) above, hairless. Shrub with a dense habit, 1.4 × 1.2m in 41 years. Mid May.  
Etymology: named for the daughter of Orine Holm, part owner of Griswold Nursery

(v) **'Lava Fountain'**

cv. Lepidote rhododendron: 'Sunny' × *rubineiflorum*, dwarf form  
H: T. Lelliott (c.1999), G: R. Marques & S. Bertelmann (pre 2004), N: R. Marques & S. Bertelmann (2009), I: Pacific Island Nursery (2004), REG: R.F. Kruppa (2009)  
Fls 3–5/flat truss, tubular funnel-shaped, the face of the flower flattening and becoming convex with age,

45 × 45–50mm, with 6 flat-edged lobes, moderate red (183C) in bud, opening inside and out deep yellowish pink (43C) to vivid reddish orange (44C), the tube somewhat lighter than 44C; unmarked; filaments white, anthers very dark orange; style very light green, stigma reddish orange. Lvs of two sizes and shapes: some elliptic, 25 × 10mm, acute at apex, cuneate at base, others ovate, 35 × 15mm, broadly acute at apex, rounded at base, both with slightly upcurved margins, semi-glossy and dark bluish green (131A) above, with no visible indumentum. Shrub 0.4 × 0.45m in 5 years. Flowering three to four times a year for extended periods between May and November/December.

(r) **'Lavender Cushion'**

cv. Lepidote rhododendron: parentage unknown  
H: W.F. Winkler (1980s), G: W.F. Winkler or Cummins Garden (1980s), N: J. van de Sande (2009), I: W.F. Winkler or Cummins Garden (1980s), REG: J. van de Sande (2009)  
Fls 7/ball truss, saucer-shaped, 25 × 35mm, with 5 wavy-edged lobes, moderate purplish pink (66D) in bud, opening inside very light purple (76B), with tiny, darker purple speckles on the dorsal lobe; outside strong reddish purple (78B), fading to light purple (76B) at calyx. Calyx c.3mm long, light yellow-green (140D). Lvs oblong, 75 × 15mm, acuminate and twisted at apex, cuneate at base, margins downcurved, semi-glossy and moderate olive green (137A) above, hairless. Shrub with a dense habit, forming a neat, rounded mound, 0.55 × 0.9m in 20 years. Early May.  
Syn: 'Winkler D-67'  
Published in RareFind Nursery cat. 2009, p.24.  
Etymology: the epithet is descriptive of the plant

(r) **'Laxmi'**

cv. Elepidote rhododendron: 'Golden Splendour' (s) × 'Edward Dunn' × *auriculatum*  
H: K.-H. Hübbes (2000), G: K.-H. Hübbes (2009), N: K.-H. Hübbes (2010), REG: J. Westhoff (2010)  
Fls 8–9/flat truss, narrowly funnel-shaped, 120 × 80mm, with 6–7 wavy-edged lobes, buds strong pink (49A), opening inside a blend of pastel colours, mainly pale yellowish pink (27D), tinged with pink

on the lobes, in the throat brilliant yellow (10A), outside a blend of pastel colours (pale yellowish pink (27D), pale orange-yellow (16D) and deep pink (47D)), with greenish at the base and darker veins from the base to the margin; with a deep red (185A) blotch on the upper third of the throat. Calyx 3mm long, greenish, with white hairs. Lvs oblong, 150 × 70mm, obtuse and mucronate at apex, cordate at base, margins decurved, matt green above, with whitish hairs on the midrib when young. Shrub 1.4 × 1.3m in 10 years. Mid June.

Etymology: named for the wife of the hybridiser

(r) **'Lily and Flora'**

cv. Elepidote rhododendron: 'Roza Stevenson' × unknown

H: R.J. Gilbert (c.1999), G: R.J. Gilbert,

N: R.J. Gilbert, REG: R.J. Gilbert (2009)

Fls 12/hemispherical truss, very broadly funnel-shaped, 45 × 82mm, with 6 slightly wavy lobes, buds a combination of deep apricot, lime green and brown, opening inside and out light yellow-green/pale yellowish green (2C/D), unmarked; stamens 13, filaments translucent lime green, anthers brown; style translucent lime green, stigma green. Calyx 7.5mm long, lobed, transparent lime green. Lvs elliptic, to 100 × 45mm, margins decurved, acute and apiculate at apex, narrowly rounded at base, hairless. Shrub 1.4 × 1.5m in 10 years. Early to mid May.

Nomenclatural standard: colour print provided by the registrant (WSY0112420). Etymology: named for the two dogs belonging to the registrant

(r) **'Lily Dance'**

cv. Elepidote rhododendron: *yakushimanum* 'Koichiro Wada' × 'Goldkrone' × 'Whitney's Tiger Lily'

H: T. Huisman (1995), G: T. Huisman (2001),

N: T. Huisman (2009), I: T. Schröder (2010),

REG: T. Huisman (2009)

Fls 10–14/ball truss, funnel-campanulate, 80–90 × 50–65mm, with 5 wavy-edged lobes, buds vivid reddish orange (33A–40B), opening inside creamy white (159D) in the throat, shading to pale purplish pink (62D) at the margins, outside whitish at the base (N155), shading to strong purplish pink

(62A–68B) at the margins, with strong yellowish pink (38A) ribs; dorsal lobe with a 12mm wide basal blotch of deep red (185A), which breaks up into moderate red (180B) dots which spread left and right onto the adjacent lobes. Calyx 25–30mm long, petaloid, pale yellow (18C), with a basal blotch 20–25 × 8–10mm, ending in dots, the blotch at base strong red (46A), the spots vivid red (46C). Lvs elliptic, 75–125 × 45–55mm, broadly acute at apex, rounded at base, margins flat, matt green above, hairless. Shrub 1.5 × 1.5m in 10 years. Early to mid May.

Nomenclatural standard: colour print of an electronic image provided by the raiser

(WSY0108033). Grown as seedling no. Z 00–13.

Etymology: the flower looks a little lily-like and the hose-in-hose form recalls the tutu skirts worn by female dancers

(r) **'Limonanda'**

cv. Elepidote rhododendron: *brachycarpum* hybrid

H: T. Huisman (1994), G: T. Huisman (1999),

N: T. Huisman (2009), I: T. Schröder (2010),

REG: T. Huisman (2009)

Fls 10–15/ball truss, open funnel-shaped, 55–65 × 65–75mm, with 5(–6) wavy-edged lobes, buds brilliant orange (25C), with light yellow (20B) veins, opening inside brilliant greenish yellow (5B) in the throat, shading to pale greenish yellow (2D) at the margins, with inconspicuous markings in the dorsal throat of near yellowish white (156D), outside light greenish yellow (3C), shading to pale greenish yellow (2D) at the margins. Calyx 1.5mm long, light olive (152A). Lvs elliptic, 75–110 × 30–40mm, acute at apex, cuneate at base, margins flat, glossy green above, hairless. Shrub 1.2 × 1.2m in 10 years. Mid May.

Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0108034). Etymology: named on the basis of the flower colour

(r) **'Linda Malland'**

cv. Elepidote rhododendron: 'Scintillation' × unknown

H: D.A. Griswold (1969), G: Griswold Nursery

(1980), N: Griswold Nursery (2010),

REG: Griswold Nursery (2010)

Fls 5/flat truss, open funnel-shaped, *c.*65 × *c.*65mm, with 5 wavy-edged lobes, moderate purplish pink (65A) in bud, opening inside pale purplish pink (65D), with discrete spots of strong red (39A) on the dorsal lobe and adjacent portions of the contiguous lobes, outside pale purplish pink (65D) or paler, with pale purplish pink (65D) midribs. Calyx 25mm long, light yellow (18B), spotted. Lvs elliptic, 150 × 40mm, broadly acute at apex, cuneate at base, margins downcurved, dull and moderate yellowish green (138A) above, hairless. Shrub 1.4 × 1.4m in 41 years. Mid May.

Etymology: named for a granddaughter of the hybridiser

(v) **‘Little Duchess’**

cv. Lepidote rhododendron: *anagalliflorum* (s) × *pauciflorum*

H: J.L. Rouse, G: J.L. Rouse, N: L. Rouse (2009), REG: A. Rouse (2009)

Fls 2–3/open truss, slightly down-facing, shortly campanulate, tube 12mm long, lobes 5mm long, corolla 18mm wide, with 5 flat-edged lobes, inside pale pink (49D), with fine stripes of moderate pink (49B), outside moderate pink (49B) at base and in fine longitudinal grooves in the corolla, the remainder pale pink (49D), unmarked. Lvs obovate, 10–12 × 7–9mm, margins flat, glossy green above, obtuse at apex, tapered at base, scaly. Shrub 30 × 30cm in 10 years, forming a compact, multi-branched plant. October (i.e. spring in Australia) and occasionally in autumn.

Nomenclatural standard: colour print provided by the registrant (WSY0112401). Etymology: name suggested by the eldest daughter of the registrant

(v) **‘Little Maneau Ra’**

cv. Lepidote rhododendron: ((*macgregoriae* × *laetum*) × *saxifragoides*) F<sub>2</sub> (s) × ((*macgregoriae* × *laetum*) × *saxifragoides*) F<sub>2</sub>

H: E.W. Moyles (1999), G: R. Marques & S. Bertelmann (2003), N: R.F. Kruppa (2009), I: Pacific Island Nursery (2008), REG: R.F. Kruppa (2009)

Fls 3–7/flat truss, narrowly funnel-shaped, 20–25 × 25mm, with 5 flat-edged lobes, opening through

shades of strong pink (49A, 48D, 50C) and deep purplish pink (55A), blending into a light yellow-green (*c.*2C) tube, with reddish smudges between the lobes both inside and out; anthers brownish yellow, filaments very light yellow; style brown. Lvs elliptic, 65–70 × 55mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and dark yellowish green (139A) above, hairless. Shrub 0.6–0.7 × 0.3–0.5m in 5 years. January to March and September to November.

(r) **‘Lockington Superb’**

cv. Elepidote rhododendron: ‘Countess of Lockington’ (s) × ‘Lockington Sunset’

H: D.J. Dosser (2000), G: D.J. Dosser, N: D.J. Dosser (2010), REG: D.J. Dosser (2010)

Fls 15–19/slightly conical truss, funnel-shaped, 100mm wide, with 7–8 wavy-edged lobes, buds strong purplish red (58C), opening inside light purplish pink (62C) around the margins and turning cream with age throughout, outside deep purplish pink (68A), turning cream with age, with dorsal rays of moderate purplish red (58A) spots becoming solid towards the base; slightly scented. Calyx light yellowish pink (36A). Lvs oblanceolate, 100 × 25mm, acute at apex, cuneate at base, margins decurved, matt green above. Shrub 1.0 × 1.0m in 10 years. October (Australia).

Nomenclatural standard: colour print provided by the registrant (WSY)

(r) **‘Lockington’s Pearl’**

cv. Elepidote rhododendron: ‘Miss B.L. Jones’ (s) × unknown

H: D.J. Dosser (2000), G: D.J. Dosser, N: D.J. Dosser (2010), REG: D.J. Dosser (2010)

Fls 19/conical truss, campanulate, 50 × 65mm, with 6–7, wavy-edged lobes, strong purplish pink (62A) in bud and throughout on opening, not marked; slightly scented. Lvs lanceolate, 175 × 45mm, acute at apex, truncate at base, margins decurved, matt green above, hairless. Shrub 1.5 × 1.0m in 10 years. October (Australia).

Nomenclatural standard: colour print provided by the registrant (WSY)

(r) **'Logan Delight'**

cv. *Lepidote rhododendron*: selection from *edgeworthii*

G: Logan Botanic Garden, S: Logan Botanic Garden (2008), N: R. Baines (2008), REG: Logan Botanic Garden (2009)

Fls 2–4/ loose, open truss, mostly between out and down-facing, funnel-campanulate, 50 × 80mm, with 5 distinctly wavy-edged lobes which are often recurved at the tip, buds white, flushed pink, opening inside white with pink speckles, outside white with pink blotches towards the apex; filaments and style white, stigma pink, anthers very dark; strongly scented. Calyx 20–25mm long, bright red, contrasting strongly with the white corolla. Lvs oblong-lanceolate, 70 × 25mm, margins wavy, acute at apex, cuneate at base, matt green and strongly rugose above, with a woolly, rusty brown indumentum below. Shrub 1.0 × 1.0m in 20 years. Late April.

Etymology: named to highlight the plant's association with Logan Botanic Garden (Scotland)

(r) **'Logan Surprise'**

cv. *Lepidote rhododendron*: selection from *edgeworthii*

G: Logan Botanic Garden, S: Logan Botanic Garden, N: R. Baines (2008), REG: Logan Botanic Garden (2009)

Fls 2–4/truss, funnel-campanulate, 60–80 × 95mm, with 5 distinctly wavy-edged lobes, some reflexed at tips, buds white, flushed with pink, opening inside white, with yellow flecks below the dorsal lobes and a yellow basal blotch; outside white; filaments and style white, stigma pinkish, anthers very dark; very strongly scented. Calyx 15mm long, reddish. Lvs oblong-lanceolate, 80 × 30mm, margins wavy, acute at apex, cuneate at base, glossy green and rugose above, with a woolly, rusty brown indumentum below. Shrub 1.0 × 1.0m in 20 years. Late April.

Etymology: named to highlight the plant's association with Logan Botanic Garden (Scotland)

(r) **'Luscious Lora'**

cv. *Lepidote rhododendron*: *fortunei* (s) × 'Caractacus'

G: H.E. Reiley (1974), N: H.E. Reiley (2007), REG: Mrs M.L. Reiley (2010)

Fls 9–11/domed truss, broadly funnel-shaped, 55 × 75mm, with 6 wavy-edged lobes, buds strong purplish red (63B), edged light purplish pink (63C), opening inside pale purplish pink (62D), edged light purplish pink (63C), the throat strong purplish red (64A), with two rows of strong purplish red (64B) speckles extending upwards as far as 10–15mm on the dorsal lobes; outside light purplish pink (62C), with light purplish pink (63C) midribs; stigma moderate brown (165A), style light yellow-green (144D); ovary strong yellow-green (144A); moderately scented. Calyx *c.* 3mm long, brilliant yellow-green (149A). Lvs oblong, 125–140 × 65mm, obtuse at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (137B) above, veins strong yellow-green (144A); new leaves very glossy. Shrub 2.5 × 3.1m in 30 years. Early May. Grown from seed purchased from the F.W. Schumacher Co. (Sandwich, Maryland, USA) in the 1960s

(v) **'Lydia Ellen'**

cv. *Lepidote rhododendron*: *suaveolens* (s) × 'Great Scentsation'

H: B.E. Oldham (1996), G: B.E. Oldham (2003), N: B.E. Oldham (2009), REG: B.E. Oldham (2010)

Fls (8–)9(–12)/flat truss, trumpet-shaped, 60 × 50mm, with 5 slightly waved lobes, inside strong purplish red (59D) on lobes, to deep purplish pink (58D) at the tube; outside strong purplish red (59D) blending to deep or moderate purplish pink (66C–D) on the inner lobes and tube; unmarked; stamens strongly exerted when mature, filaments strong purplish red (58B), anthers moderate red (183C); style strong red (53C), stigma deep purplish red (59B); strongly scented. Calyx less than 1mm long, light yellow-green (143C). Fls subtended by pale purplish pink tipped bracts. Lvs broadly elliptic, 110 × 55mm, margins flat, acute at apex, broadly tapering at base, dark yellowish green (139A) above, with sparse, strong greenish yellow (153C) scales mostly above when young, lost with age. Shrub 1.1 ×

0.7m in 11 years. Repeat flowering; unusual in being long-lasting as a cut flower.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the registrant's granddaughter, who admired it

(r) **'Magenta Medley'**

cv. Elepidote rhododendron: 'Midnight Mystique' (s) × (*yakushimanum* 'Koichiro Wada' × 'Burgundy') × 'Purple Splendour'

H: F. Fujioka (1993), G: G.H. Kesterson (1998), N: G.H. Kesterson (2009), REG: G.H. Kesterson (2009)

Fls 27/conical truss, saucer-shaped, c.65 × 90mm, with 5 very wavy-edged lobes, strong reddish purple (70B) in bud, opening inside very pale purple (73D) near the centre, gradually darkening to strong reddish purple (72B) at margins, and with a strong red (53D) blotch in the throat; outside strong purplish pink (68B), darkening to a 25mm wide strong purple (77B) margin, and with strong reddish purple (72C) midribs; a few dark red (59A) spots on the dorsal lobes. Calyx negligible. Lvs elliptic, 175 × 50mm, acute at apex, cuneate at base, margins flat, semi-glossy and dark green (131B) above, hairless. Shrub 1.2 × 0.9m in 10 years. Mid May.

(r) **'Mahalon'**

cv. Elepidote rhododendron: 'Kernéostic' (s) × 'September Song'

H: Société Bretonne du Rhododendron (1997), G: J.-F. Saint-Jalm (2003), N: Société Bretonne du Rhododendron (2009), REG: Société Bretonne du Rhododendron (2009)

Fls 7–9/ domed truss, funnel-campanulate, 55 × 95mm, with 5–7 flat-edged lobes, vivid red (45B) in bud, opening inside light orange-yellow (19B); outside light orange (28C), with vivid red (45B) veining; inconspicuous, strong yellowish green (N144A), dorsal spotting; stamens 11–13; stigma green. Calyx 2–4mm long, strong yellow-green (144A). Lvs elliptic, 125 × 55mm, margins flat, obtuse and shortly apiculate at apex, broadly cuneate to narrowly and shallowly rounded at base, moderate olive green (137A) above, hairless. Shrub 1.2 × 1.2m in 11 years. Late April to early May.

Nomenclatural standard: colour print provided by the registrant (WSY0112417). Etymology: the epithet means "my heart" in Breton and is also the name of the town where the 5th show of the Société Bretonne du Rhododendron took place in 2009

(r) **'Mama Cook'**

cv. Elepidote rhododendron: *macrophyllum* × 'Britannia' (s) × 'Ruby Bowman' × *elliottii* 'War Paint'

H: D.W. Goheen (1997), G: S.B. Southerland (2000), N: S.B. Southerland (2000), REG: S.B. Southerland (2010)

Fls up to 18/domed truss, open funnel-shaped, 50 × 65mm, with 5 wavy-edged lobes, dark red in bud, opening inside and out close to moderate red (47A), with tiny dark red speckles scattered randomly on all lobes, a concentration of these speckles forming a dorsal blotch. Lvs elliptic, 145 × 65mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and dark green above, with a thin coating of very faint orange indumentum below. Shrub with an open habit, 1.5 × 1.8m in 12 years. Early May. Etymology: named for the registrant's mother-in-law

(v) **'Margaret Jane'**

cv. Lepidote rhododendron: 'Solar Flare', open-pollinated

H: D. & P.J. Brown (2002), G: D. & P.J. Brown (2007), N: D. & P.J. Brown (2009), I: Browns Nursery, REG: D. & P.J. Brown (2009)

Fls 7–8/lax truss, tubular funnel-shaped, 40 × 50mm, with 5 slightly wavy-edged lobes, buds deep yellowish pink (43C), opening inside pale yellow (18C) with strong yellowish pink (43D) on the margins of the lobes, unmarked, outside pale yellow (18C), suffused with vivid reddish orange to strong yellowish pink (43B–D), with a bright spot of vivid reddish orange (43A) at each lobe sinus; with a sweet, spicy scent. Calyx vestigial. Lvs ovate, 120 × 65mm, cuneate to rounded at base, acute at apex, margins flat, matt green above, hairless; new foliage bright pink/copper, aging to green. Shrub 1.5 × 1.0m in 5–7 years. Repeat-flowering.

Nomenclatural standard: colour print provided by the registrant (WSY01122254)

(v) **'Marshmallow Cream'**

cv. Lepidote rhododendron: 'Pastenello' hybrid (s) × ((*konori* var. *phaeopleplum* × *lochiae*) × *zoelleri*) × *leucogigas*

H: G.L.S. Snell (2001), G: R. Marques &

S. Bertelmann (2006), N: R. Marques &

S. Bertelmann (2009), I: Pacific Island Nursery (2008), REG: R.F. Kruppa (2009)

Fls 12–17/domed truss, elongated tubular funnel-shaped, 95 × 90mm, with 5 wavy-edged lobes, buds opening inside pale yellow-green (155A) to brilliant yellow-green (150C), with light green lines visible inside the tube; outside pale yellow-green (155A) at the margin, the tube becoming yellow-green to light green; extremely pale yellow and green blotches at base; stigma and style light green; anthers yellowish brown, filaments very light greenish yellow. Calyx vestigial. Lvs elliptic, 175 × 70mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and dark green (136A) above, with no visible indumentum. Shrub with an open, upright habit, 1.2 × 0.6m in 8 years. Flowering two to three times a year: usually January/February, July and September.

(a) **'Mary Lou Dority'**

cv. Evergreen azalea: 'Betty Christopher' (s) × 'Komo-kulshan'

H: J.F. Klimavicz (2000), G: J.F. Klimavicz (2002),

N: J.F. Klimavicz (2009), REG: J.F. Klimavicz (2009)

Fls in a ball truss, c.45 × 70mm, saucer-shaped, with 5 wavy-edged lobes, very pale purple (69A) in bud, opening inside very pale purple (69A) on the tips of the lobes, fading to white in the centre, with a tinge of pale green in the throat and a large area of discrete, strong purplish red (60B) spots on the dorsal lobes; outside very pale purple (69A) between the greenish white midribs. Calyx 10mm long, strong yellow-green (144B). Lvs elliptic, 125 × 50mm, broadly acute at apex, cuneate at base, margins flat, glossy and moderate olive green (146A) above, hairless. Shrub with a dense habit, 0.5 × 0.5m in 5 years. Mid May.

Etymology: named for the mother-in-law of the registrant

(r) **'Mascara'**

cv. Elepidote rhododendron: 'Queen Souriya' (s) × 'Rotgold'

H: K.-H. Hübbbers (1989), G: K.-H. Hübbbers (1996),

N: Mrs L. Hübbbers (2009), REG: J. Westhoff (2009)

Fls c.8/loose, open truss, broadly funnel-shaped, 55 × 80mm, with 6 wavy-edged lobes, light greenish yellow to pale yellow-green (4C–D) both inside and out; a very few greenish spots in the dorsal third of the throat; the striking dark anthers contrast well with the corolla; scented when opening. Calyx 2mm long, green, with a red rim. Lvs elliptic-oblong, 80 × 50mm, rounded and with a point at the tip, truncate at the base, margins flat, glossy above, hairless.

Shrub 2.2 × 2.7m in 20 years. Early April.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112993).

Etymology: the anthers are said to look like eyelashes coloured and shaped by a mascara wand

(v) **'Maui Maiden'**

cv. Lepidote rhododendron: (*macgregoriae* × *laetum*) × *saxifragoides*, F<sub>2</sub> (s) × (*macgregoriae* × *laetum*) × *saxifragoides*, F<sub>2</sub>

H: E.W. Moyles (1999), G: R. Marques &

S. Bertelmann (2004), N: R. Marques &

S. Bertelmann (2009), I: Pacific Island Nursery (2008), REG: R.F. Kruppa (2009)

Fls 3–5/flat to lax truss, tubular funnel-shaped, 45 × 40mm, with 5 flat-edged lobes, moderate purplish pink (68C) in bud, opening inside and out strong red (41B) at margins, shading to light yellowish pink (38C) towards the base; unmarked; filaments pink, anthers brownish red; style red, stigma dark red. Lvs elliptic, 50–55 × 20–25mm, acute at apex, cuneate at base, margins upcurved, semi-glossy and dark yellowish green (139A) above, with no visible indumentum. Shrub 0.4–0.5 × 0.3–0.4m in 5 years. Flowering three times a year, for two to three months each time.

(r) **'Mesha'**

cv. Elepidote rhododendron: 'Janet Blair' (s) × unknown

H: unknown (c.1990), G: Alan Suchy (c.1994),

N: Angela Suchy (c.1996), REG: Alan Suchy (2009)

Fls 9/domed, but open-topped truss, tubular funnel-shaped, 45 × 70mm, with 5 wavy-edged lobes, pink in bud, opening inside yellowish cream, the lobes with pink borders, with a dorsal blotch of small, red speckles flaring outward from the base; outside pink. Lvs elliptic, 125 × 50mm, broadly acute at apex, cuneate at base, margins flat, dull and mid green above, hairless. Shrub 1.2 × 0.6m in c.5 years. Mid May.

Etymology: named for the ballet dancer Mikhail Baryshnikov (nick-named “Mesha”) by the registrant’s daughter

(r) ‘**Metarose**’

cv. Elepidote rhododendron: ‘Scintillation’ × ‘Furnivall’s Daughter’ (s) × ‘Queen Souriya’  
H: K.-H. Hübbbers (1989), G: K.-H. Hübbbers (1996), N: K.-H. Hübbbers (2004), REG: J. Westhoff (2009)  
Fls c.12/rounded truss, broadly funnel-shaped, 60 × 90mm, with 8 wavy-edged lobes, buds strong purplish red (63B), opening inside light purplish pink (62C), becoming paler in the centre, outside strong purplish red to strong purplish pink (63B–C); dense, strong greenish yellow (151B) spots covering the dorsal throat. Calyx 2mm long, translucent green. Lvs oblong-oblanceolate, 80 × 40mm, rounded and with a point at the apex, truncate at the base, margins flat, glossy above, hairless. Shrub 2.3 × 2.0m in 15 years. April.  
Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0112982). Etymology: the original plant is growing next to a *Metasequoia* in the raiser’s garden and it has pink flowers

(r) ‘**Mister Benoit**’

cv. Elepidote rhododendron: ‘Fred Peste’ (s) × unknown  
H: J.-F. Saint-Jalm (1996), G: J.-F. Saint-Jalm (2002), N: J.-F. Saint-Jalm (2009), REG: J.-F. Saint-Jalm (2009)  
Fls 10/domed truss, funnel-campanulate, 37 × 52mm, with 5 smooth-edged lobes, strong purplish red (58B) in bud, opening very pale purple (69C), with broad margins of strong purplish red (58B), outside strong purplish red (58C); without blotch or

spots; stamens very short (c.12mm long); ovary white hairy. Calyx vestigial. Lvs elliptic, 125 × 48mm, margins decurved, broadly acute at apex, cuneate/rounded at base, glossy and dark greyish green (N189A) above, hairy below (hairs yellowish grey (198D) at first, aging to deep orange-yellow (163A). Shrub 0.95 × 1.2m in 12 years. Early May to early June.

Nomenclatural standard: colour print provided by the registrant (WSY0112748). Etymology: named for the boyfriend of the registrant’s daughter

(v) ‘**Moyles’s Marvel**’

cv. Lepidote rhododendron: ((*macgregoriae* × *laetum*) × *saxifragoides*) F<sub>2</sub> (s) × ((*macgregoriae* × *laetum*) × *saxifragoides*) F<sub>2</sub>  
H: E.W. Moyles (1999), G: R. Marques & S. Bertelmann (2004), N: R. Marques & S. Bertelmann (2009), I: Pacific Island Nursery (2008), REG: R.F. Kruppa (2009)  
Fls 5–6/flat truss, narrowly funnel-shaped, 50 × 50mm, with 5 flat-edged lobes, dark purple in bud, opening white (155D) throughout, blending to yellowish white (158C) and light greenish yellow (4C) at the base of the tube; entire corolla ages to yellowish white (158C) in 3–4 days, the lobes edged with moderate to light pink (49B–C), the pink colour spreading and darkening with age; filaments white, anthers greenish; style very light green, stigma reddish brown. Calyx vestigial. Lvs elliptic, 90 × 50mm, broadly acute at apex, slightly rounded at base, margins upcurved, semi-glossy and dark yellowish green (139A) above, with no visible indumentum. Shrub with an open habit, 0.6 × 0.5m in 5 years. Usually flowering in winter, spring and summer, for periods of two months each.

(r) ‘**Nancy Steele**’

cv. Elepidote rhododendron: *aureum* (s) × ‘Prelude’  
H: R.M. Steele (1968–1969), G: R.M. Steele, N: R.M. Steele, I: Bayport Plant Farm (mid 1980s), REG: R.M. Steele (2009)  
Fls 11/lax, domed truss, funnel-campanulate, 38 × 60mm, with 6 flat to very slightly wavy-edged lobes, brilliant greenish yellow (3A–B) in bud, opening inside light yellow-green (2C), with barely detectable

darker yellow/green spotting on the dorsal lobes, outside light yellow-green (2C) to light greenish yellow (1C). Calyx 0.1–0.2mm long, green, edged red. Lvs elliptic, 105 × 45mm, obtuse at apex, cuneate/rounded at base, slightly convex, with smooth edges which are narrowly rolled over, slightly matt and moderate yellow-green (147C) above, close to the same colour below in shade but becoming more yellowish in the sun, hairless. Shrub with a dense mounding habit, 2.3 × 2.7m in 25 years. Very early spring (with *mucronulatum* and *dauricum* or even earlier).

Etymology: named for the wife of the registrant. The shrub size is of a plant in Yarmouth, Nova Scotia (US hardiness zone 7a); the Bayport Nursery Plant (zone 6a) used for propagation is 1.2 × 2.4m in 34 years. Previously distributed under parentage and the breeder's reference Bpt 80–5

(r) **'Ned's Number One'**

cv. Elepidote rhododendron: *yakushimanum* × *fortunei* subsp. *discolor* × *catawbiense*  
H: N. Schrope (1960), G: N. Schrope (1966),  
N: N. Schrope, I: Brouse Nursery (2005),  
REG: F.W. Brouse (2010)  
Fls 13/lax ball truss, funnel-shaped, 40–50 × 40–55mm, with very wavy-edged lobes, pink in bud, opening very light pink, with a light yellow dorsal blotch. Calyx c.5mm long, tan. Lvs elliptic, 100 × 25mm, apiculate at apex, cuneate at base, margins upcurved, dark green above, hairless. Dense shrub, 0.6 × 0.9m in 6 years. Mid May.

(a) **'Niamh'**

cv. Deciduous azalea: parentage unknown  
H: E.G. Millais (c.1990), G: E.G. Millais (c.1994),  
N: D.G. Millais (2010), I: D.G. Millais (2010),  
REG: D.G. Millais (2010)  
Fls single, 12–14/rounded truss, tubular funnel-shaped, 60–70 × 80mm, with 5 wavy-edged lobes, buds deep pink, opening inside deep pink (50B) and shading to strong pink (52D), outside vivid red (52A) on the tube, the lobes deep pink (52B); prominent vivid orange-yellow (23A) dorsal blotch; stigma green; strongly scented. Calyx 3–5mm long, pale green. Lvs elliptic, 80–90 × 40mm, glossy

above, margins decurved, hairy; new growth/foilage reddish, appearing with the flowers. Shrub 1.2 × 1.2m in 10 years. Late May to early June.

Nomenclatural standard: herbarium specimen provided by the registrant (WSY). Etymology: named for the registrant's daughter

(v) **'Oldham's Theresa'**

cv. Lepidote rhododendron: *suaveolens* (s) × 'Great Scentsation'

H: B.E. Oldham (1996), G: B.E. Oldham (2004),  
N: B.E. Oldham (2007), REG: B.E. Oldham (2010)  
Fls (9–)11(–13)/ loose, upright, umbellate truss, 65–70 × 45–55mm, with 5 flat-edged lobes, dark red (59A) in bud, opening inside strong purplish red (58B) on the lobes, the tube white (155D) seasonally tending to pale purplish pink (56C), outside strong purplish red (58B) blending to strong purplish pink (63C) on the lobes, the tube white (155D) blending to pale purplish pink (56C); unmarked; anthers dark purple (83A); styles vivid red (45B); moderately scented. Lvs elliptic to broadly elliptic, 105 × 60–65mm, margins flat, acute at apex, obtuse to truncate at base, with sparse, slightly unequal, vivid reddish orange (34B) scales, mostly lost with age; young foliage greyish red (182B) above, aging through light olive (152A) to moderate olive green (137A), lower surface greyish red (182B) with a strong yellow-green (144B) midrib, aging to moderate yellow-green (147C–D). Dense, rounded shrub, 1.2 × 1.3m in 12 years. Frequently repeat flowering. Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the daughter-in-law of the registrant

(r) **'Oliver James'**

cv. Elepidote rhododendron: 'Marcus of Lockington' (s) × 'Australian Sunset'  
H: D.J. Dosser, G: D.J. Dosser (2005),  
N: D.J. Dosser (2010), REG: D.J. Dosser (2010)  
Fls 13–15/ball truss, funnel-shaped, 40 × 52mm, with 6 strongly waved lobes (sometimes with irregularly reflexed margins and apices), buds deep purplish pink (55A), opening inside deep purplish pink (55A) and light purplish pink (55C), outside deep purplish pink (55A), with a 12mm long ray of

moderate purplish red (64A); scented. Calyx 40mm long, deep purplish pink (55A), with reflexed lobes. Lvs lanceolate, 150 × 30mm, broadly acute at apex, truncate at base, margins decurved, hairless. Shrub 2.0 × 1.5m in c.15 years. October (Australia). Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the deceased baby son of the registrant's neighbour

(v) **'Olivia Knight'**

cv. Lepidote rhododendron: *tuba* (s) × 'Rangitoto Rose'

H: B.E. Oldham (1993), G: B.E. Oldham (1998), N: B.E. Oldham (2008), REG: B.E. Oldham (2009)  
Fls 11/flat truss, trumpet-shaped, with a slightly curved tube, 90 × 70mm, with 7 smooth-edged lobes, buds deep purplish pink (61D), with deep purplish pink (64D) on the bracts, opening inside light purplish pink (62C) at the margins, blending to moderate purplish pink (62B) at the tube, with a small blotch of strong purplish red (63B) by the sinus between the lobes, outside pale purplish pink (62D) at margins, streaked and shaded to deep purplish pink (58D) on the tube; style strong reddish orange (42C), stigma brilliant greenish yellow (151D); anthers light yellow-green (150D), filaments white, tinged light yellowish pink (36C) at each end; strongly scented. Calyx disc 2mm long, indistinctly and minutely lobed, moderate purplish red (59C). Lvs very broadly elliptic, 105 × 55–60mm, margins flat, young leaves matt and strong to dark greenish yellow (151A–152D) above, acute at apex, truncate at base, with moderate orange scales when young, without scales when mature. Upright shrub, 1.7 × 0.8m in 15 years, with foliage concentrated near the top of a single stem. Frequent repeat-flowering. Nomenclatural standard: colour print provided by the registrant (WSY0112744). Etymology: named after the registrant's grand-daughter

(r) **'Palni Fire'**

cv. Elepidote rhododendron: *arboreum* subsp. *nilagiricum* (s) × *griersonianum*  
H: M.S. Viraraghavan (1987), G: M.S. Viraraghavan (2001), N: M.S. Viraraghavan (2010), REG: M.S. Viraraghavan (2010)

Fls 9/rounded truss, tubular-campanulate, 35 × 30mm, with 5 slightly wavy-edged lobes, orange-red in bud and throughout on opening, with relatively inconspicuous darker dorsal spotting. Calyx 10mm long, light brown. Lvs oblanceolate, 100 × 40mm, broadly cuneate at base, pointed at apex, margins flat, matt green above, with a creamy brown indumentum below which ages to light brown. Shrub 1.5 × 1.5m in 20 years. Late April to early May.

Etymology: Palni is in reference to the Palni Hills in southern India where the registrant lives and where this plant was raised

(r) **'Palni Sunrise'**

cv. Elepidote rhododendron: *arboreum* subsp. *nilagiricum* (s) × 'Trude Webster'  
H: M.S. Viraraghavan (1987), G: M.S. Viraraghavan (2005), N: M.S. Viraraghavan (2010), REG: M.S. Viraraghavan (2010)  
Fls 11/rounded truss, open campanulate, 50 × 60mm, with 5 flat-edged lobes, orange-red in bud and throughout on opening, aging to salmon pink; unmarked. Calyx 20mm long, light brown. Lvs broadly elliptic, 120 × 45mm, cuneate at base, broadly acute at apex, margins wavy, matt green above, with an off-white indumentum below which ages to light brown. Shrub 2.5 × 1.5m in 20 years. Early May.  
Etymology: Palni refers to the Palni Hills in southern India, where the registrant lives and where this plant was raised

(r) **'Paotrig Skaer'**

cv. Elepidote rhododendron: 'Kluis Sensation' (s) × 'Karen Triplett'  
H: L. Le Duigou (2000), G: A. Le Duigou (2006), N: L. & A. Le Duigou (2010), REG: L. & A. Le Duigou (2010)  
Fls 19/ball truss, funnel-campanulate, 60 × 80mm, with 5 wavy-edged lobes, strong red (51A) in bud, opening inside pale yellow (18D) and outside pale yellow (20D); small spots of strong reddish orange (31A) on the dorsal lobe; stamens 11; style deep pink (51C). Calyx petaloid, 10–35mm long, light orange-yellow (19B), with vivid yellowish orange

(30C) spots on the dorsal lobe. Lvs elliptic, 140 × 50mm, rounded at base, broadly acute at apex, glossy green above, margins decurved, hairless. Compact, rounded shrub, 0.95 × 0.95m in 9 years. May. Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: “Paotrig” means “small boy” in Breton, Skaer is the Breton spelling of their local town. This plant and ‘Plarc’hig Skaer’ are seedlings from the same cross

(r) **‘Papa Cook’**

cv. *Elepidote* rhododendron: ‘Red River’ (s) × *hyperythrum*

H: S.B. Southerland (1997), G: J.W. Cook (2001), N: S.B. Southerland (2001), REG: S.B. Southerland (2010)

Fls 16/domed truss, broadly funnel-shaped, 50 × 70mm, with 5 flat-edged lobes, vivid purplish red (N57A) in bud, opening inside deep purplish pink (68A) at margin, fading to lighter pink and then white towards the centre, with a spotted vivid red (45A) dorsal blotch; outside deep purplish pink (73A), fading to white at the base. Lvs elliptic, up to 150 × 70mm, broadly acute at apex, rounded at base, margins downcurved, thick and with a depressed midrib, hairless. Shrub with a dense habit, 1.2 × 1.2m in 13 years, heat-tolerant. Early April.

Etymology: named for the registrant’s father-in-law

(v) **‘Pastel Splendour’**

cv. *Lepidote* rhododendron: ‘Gardenia Odyssey’ (s) × ‘Rangitoto Rose’

H: B.E. Oldham (1993), G: B.E. Oldham (2000), N: B.E. Oldham (2009), REG: B.E. Oldham (2009) Fls 13–16/semi-erect, fairly dense, more or less domed truss, tubular funnel-shaped, 90 × 95mm, with 5 wavy-edged, broadly overlapping lobes, buds light yellow-green (145C), with strong red (46A) on the bract margins, opening moderate yellowish pink (39C) at margins, shading to light orange-yellow (22D) in the tube, outside light yellowish pink (29C), shading though light orange-yellow (22C) to brilliant yellow-green (150B) at the base; small streaks of strong red (47B) inside the tube; strongly scented; flower colour varying somewhat with the

seasons. Calyx 1mm long, deep purplish pink (61D). Lvs narrowly elliptic, 115 × 35–40mm, margins flat, matt grey-green above, acute to attenuate at apex, narrowly tapering at base, scaly, the scales becoming brown and more sparse with age. Shrub 1.2 × 0.75m in 9 years. Repeat-flowering.

Nomenclatural standard: colour print provided by the registrant (WSY0112745). Etymology: name descriptive of “this outstanding hybrid”

(a) **‘Patsy’s Choice’**

cv. *Evergreen* azalea: ‘Hinodegiri’, open-pollinated (s) H: J.O. Thornton (1985), G: J.O. Thornton, N: J.O. Thornton, REG: J.O. Thornton (2009)

Fls 3/flat truss, tubular funnel-shaped, 30 × 40mm, with 5 flat-edged lobes, strong pink (48C) in bud, opening strong pink (48C) throughout. Calyx c.5mm long, light green. Lvs obovate, 20 × 10mm, obtuse at apex, cuneate at base, margins flat, glossy and mid green above, changing to bronze in winter, with no visible indumentum. Shrub with a dense habit, 1.2 × 1.2m in 25 years. Early.

Etymology: named for the wife of the raiser to commemorate their 50th wedding anniversary

(r) **‘Pemako Fairy’**

cv. *Lepidote* rhododendron: *pemakoense* (s) × *keiskei* var. *ozawae* ‘Yaku Fairy’

H: B.N. Starling (1995), G: B.N. Starling (1999), N: B.N. Starling (1999), I: B.N. Starling (2001), REG: B.N. Starling (2009)

Fls 3/truss, funnel-shaped, 15–18 × 30mm, with 5 wavy-edged lobes, buds pale purplish pink (55D), opening pale purplish pink (56D) and fading to white; unmarked; with a prominent darker style and stigma and dark anthers. Calyx 5mm long, green. Lvs elliptic to oblong, 15–20 × 5–7mm, rounded to pointed at apex, cuneate to rounded at base, margins flat, matt green above, with scattered (non-touching), pale but dark-centred scales below. Shrub 25 × 45cm in 10 years, completely covered by flowers when grown in an open situation. Early April.

Etymology: name derived from those of its parents

(r) **'Phillipa J.'**

cv. Elepidote rhododendron: *wardii*, open-pollinated

S: E. Jago (2010), G: E. Jago (2000), N: E. Jago (2010), REG: E. Jago (2010)

Fls 8/loose rounded truss, open funnel-shaped, 60 × 40mm, with 7 wavy-edged lobes, buds lemon tinged with green, opening inside lemon to deeper lemon green at the base, outside lemon to deeper lemon at the base, unmarked; anthers brown; style lemon green, stigma light yellow. Calyx 4mm long, green, with amber brown margins. Pedicels green, tinged pinkish red. Lvs oblong, 80–90 × 30–40mm, margins upcurved, broadly acute at apex, matt mid green above, hairless. Shrub 1.6 × 2.0m in 14 years. Early to mid November (New Zealand).

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the daughter of the registrant

(r) **'Pink Sunset'**

cv. Elepidote rhododendron: parentage unknown, but possibly a selection from the cross that produced 'Mrs Eddy'

H: E. de Rothschild, G: Mrs Stent (c.1958), S: D. Millais (c.2000), N: D. Millais (2009), I: Millais Nurseries (2009), REG: D. Millais (2009)  
Fls 13–17/ low-domed truss, funnel-shaped to funnel-campanulate, up to 80 × 120mm, with 5 wavy-edged lobes, strong red (53B) in bud, opening inside strong purplish pink (63C) on dorsal lobes and slightly paler, light purplish pink (62C), on ventral lobes, aging to almost white, prominent speckling of strong to dark red (53B/187B) in the throat and on most of the dorsal lobe and adjacent lobes and spreading throughout the flower but fainter and less abundant on the other lobes; outside opening strong purplish red (63A) on dorsal lobes, a little paler on the ventral lobes (moderate purplish pink, 62B), with a darker pink line remaining down the centre of the lobes as the edges fade to almost white; stamens 10–12, filaments white, anthers pale biscuit yellow (paler than pale yellow, 158A); style white, flushed pale green, stigma red; ovary bright yellow-green, glandular, hairy. Calyx irregular, up to 3mm long, yellow-green, flushed red on exposed

surface, glandular hairy. Lvs oblong-elliptic, 85–180 × 30–55mm, margins decurved, mucronate at apex, matt and dark green above, hairless; petioles up to 50mm long, yellow-green, sometimes flushed red on exposed surface. Shrub 1.5 × 1.5m in 10 years. Late May to early June.

Nomenclatural standard: herbarium specimen provided by the registrant (WSY0112559).

Etymology: the pattern of spotting appears like the rays of a sunset. One of a batch of seven un-named plants bought from Exbury

(v) **'Pixie Star'**

cv. Lepidote rhododendron: *macgregoriae* × *rubineiflorum* (s) × *rubineiflorum* × *anagalliflorum*

H: A. Rouse (2001), G: A. Rouse (2008),

N: A. Rouse (2009), REG: A. Rouse (2009)

Fls 2–3/open truss (fls out to slightly down-facing), shortly tubular-campanulate, tube 8mm long, lobes 10mm long, corolla 25mm wide, with 5, rather narrowly oblong-obovate, wavy-edged, widely separated lobes, inside light pink (49C), outside pale pink (49D) to light pink (49C), unmarked; stamens and style projecting prominently from the tube, deeper coloured. Lvs narrowly elliptic, 12 × 5mm, margins flat, matt green above, obtuse at apex, tapered at base, scaly. Shrub 20 × 20cm in 8 years, forming a compact, multi-branched plant. October (Australia).

Nomenclatural standard: colour print provided by the registrant (WSY0112400). Etymology: name suggested by the small, star-like flowers

(r) **'Plarc'hig Skaer'**

cv. Elepidote rhododendron: 'Kluis Sensation' (s) × 'Karen Triplett'

H: L. Le Duigou (2000), G: A. Le Duigou (2006),

N: L. & A. Le Duigou (2010), REG: L. & A. Le Duigou (2010)

Fls 15/ball truss, funnel-shaped, 52 × 80mm, with 5 wavy-edged lobes, strong purplish red (61B) in bud, opening inside pale orange-yellow (19D) which fades with age, and with margins flushed pale purplish pink, outside light pink (49C); dorsal lobe with dark red (183A) spots and a deep purplish pink blotch at the base; stamens 11, filaments light yellow-green

(2C); stigma strong purplish red (61B). Calyx 2mm long, strong purplish red (59D). Lvs elliptic, 115 × 40mm, broadly acute at apex, cuneate at base, matt green above, margins decurved, hairless. Small compact shrub, 0.5 × 0.7m in 9 years. May. Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: “Plarc’hig” means “small girl” in Breton, Skaer is the Breton spelling of their local town. This plant and ‘Paotrig Skaer’ are seedlings from the same cross

(v) **‘Porcelain Doll’**

cv. *Elepidote rhododendron*: ‘Gardenia Odyssey’ (s) × ‘Bobs’s Crowning Glory’  
H: G.L.S. Snell (1999–2000), G: R. Marques & S. Bertelmann (2004), N: R. Marques & S. Bertelmann (2009), I: Pacific Island Nursery (2008), REG: R.F. Kruppa (2009)  
Fls 10–14/flat truss, tubular funnel-shaped, 100 × 75mm, with 7 flat-edged lobes, deep purplish red (59B) in bud, opening inside and out shades of pink varying from deep purplish pink (55A) through strong purplish pink (55B) and light purplish pink (55C) to pale purplish pink (56A) and almost white; the base of the tube inside is greenish yellow; some red spots at the base of the lobe sinuses (outside); anthers off-white, filaments very light yellow; stigma light green, style light brown; moderately scented. Lvs elliptic, 150 × 80mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (137A) above, with no visible indumentum. Shrub with an open habit, 1.2 × 0.6m in 8 years. Spring and summer.

(r) **‘Princess Michael of Kent’**

cv. *Elepidote rhododendron*: *fortunei* (s) × Idealist Group  
H: E. de Rothschild, G: E. de Rothschild, N: Leopold de Rothschild (2010), REG: Leopold de Rothschild (2010)  
Fls 9–11/lax truss, funnel-shaped, 60 × 90mm, with 5–7 slightly wavy-edged lobes, buds strong purplish red (60C–D), opening inside creamy yellow (near 155A), with a yellow basal blotch, outside creamy yellow (near 155A), with a pronounced deep pink (50B) blush around the edges; filaments white,

anthers brown; style creamy yellow, stigma green. Calyx 35mm long, strong red (46A). Pedicels 40–50mm long. Lvs oblanceolate, c.110 × 50mm, obtuse and mucronate at apex, cuneate at base, margins slightly upcurved, moderate olive green (147A) above, hairless. Shrub 4.0 × 3.0–4.0m in 25 years. April to May. Nomenclatural standard: colour print provided by the registrant (WSY)

(r) **‘Prinses Máxima’**

cv. *Elepidote rhododendron*: (‘Goldkrone’ × *yakushimanum*) × (‘Cosmopolitan’ × ‘Scintillation’) × ‘Whitney’s Tiger Lily’  
H: T. Huisman (1996), G: T. Huisman (2001), N: T. Huisman (2009), I: T. Schröder (2010), REG: T. Huisman (2009)  
Fls 7–10/ball truss, broadly funnel-shaped, 55–65 × 70–80mm, with 5 flat-edged lobes, buds light purplish pink (65B), opening inside very light pink at first, then becoming white (N155D), outside light pinkish white (N155C) at first, aging to almost white (N155D), spotted all over the lobes, the spots deep reddish purple (77A) to deep purplish pink (N57C) and spreading on the dorsal lobes into the 10mm wide white margin and on the lower lobes onto the 20mm wide white margin. Calyx 50–60mm long, petaloid and sometimes longer than the corolla, coloured and spotted like the corolla. Lvs elliptic to oblong, 85–145 × 35–55mm, broadly acute at apex, cuneate to rounded at base, margins flat, glossy green above, hairless. Shrub 1.6 × 1.2m in 10 years. Mid May.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0108035). Etymology: named, with permission, for the wife of the Crown Prince of the Netherlands (Prince William Alexander)

(r) **‘Quick Flick’**

cv. *Elepidote rhododendron*: ‘Autumn Gold’ (s) × Dido Group  
H: R.J. Coker (c.1985), G: R.J. Coker, N: B.J. & H. Coker (2010), REG: B.J. & H. Coker (2010)  
Fls 13/open, rounded truss, open funnel-shaped, 55 × 80mm, with 6 wavy-edged lobes, vivid reddish

orange (33A) in bud, opening vivid yellow (17C) in the throat, shading to broad pink margins; outside vivid yellow (17C) at the base, shading to vivid reddish orange (33A) suffused with pink at the base (under the calyx); with distinct brown spotting on the dorsal lobes; flower becomes more pink with age. Calyx irregular, 20–30mm long, vivid yellow (17C) at the base, shading to vivid reddish orange (33A) at the apex, with similar spotting to the corolla. Lvs oblanceolate, 105 × 40mm, margins decurved, obtuse at apex, cuneate at base, matt green above, hairless. Shrub 1.5 × 1.0m in 10 years. Midseason to late.

Nomenclatural standard: colour print provided by the registrants (WSY). Etymology: named in memory of Ron Coker who used to refer to a last drink as a “quick flick”

(v) ‘**Rangitoto Blush**’

cv. *Lepidote rhododendron*: ‘Gardenia Odyssey’ (s) × *laetum*  
H: B.E. Oldham (1993), G: B.E. Oldham (1998), N: B.E. Oldham (2009), REG: B.E. Oldham (2010)  
Fls (7–)8(–12)/flat, umbellate truss, tubular funnel-shaped, 80 × 70mm, with 5, flat-edged, markedly overlapping lobes, buds moderate red (185B), opening light yellow (18B), with strong purplish pink (55B) margins, heavily but minutely streaked strong pink (54C), blending to a streaky deep pink (50B) deep in the tube; outside light yellow (18A), with strong purplish pink (55B) margins, shading to vivid reddish orange (44B) mid tube and to light yellow-green (145B) at the base; small spots of strong red (50A) at the base of the lobes externally; style deep purplish red (61A), stigma light yellow-green (145B); scented. The flower colour varies with the time of year and climatic conditions. Calyx less than 1mm long, dark red (59A) to vivid purplish red (66A). Lvs elliptic, 150 × 65mm, margins flat, acute/attenuate at apex, tapering at base, glossy green above, new growth strong yellow-green (144A) above, tending to take on a purplish hue in cold weather, with sparse light olive (152B) scales when young, these lost with age. Upright shrub 1.8 × 1.3m in 15 years. Frequent repeat flowering.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: the plant is one of a series of this cross raised at the registrant’s previous address at Rangitoto Avenue, Remuera, New Zealand

(v) ‘**Rangitoto Dawn**’

cv. *Lepidote rhododendron*: ‘Gardenia Odyssey’ (s) × *laetum*  
H: B.E. Oldham (1993), G: B.E. Oldham (1998), N: B.E. Oldham (2009), REG: B.E. Oldham (2010)  
Fls 7/semi-erect, umbellate truss, tubular funnel-shaped, 60 × 75mm, with 5, slightly wavy-edged, slightly overlapping lobes, strong yellow-green (143C) in bud, opening inside with margins moderate pink (51D) to strong yellowish pink (37B) centrally, the tube brilliant to vivid yellow (13C–A), outside brilliant greenish yellow (6B) shading to brilliant yellow-green (154B) at the base; unmarked; style brilliant yellow-green (149C), stigma brilliant yellow-green (149B); not scented. Calyx very short, strong yellow-green (145A). Lvs elliptic, 120 × 55mm, margins flat, acute at apex, tapering at base, new growth deep greenish yellow (153A), maturing to moderate olive green (137A) above, with moderate reddish orange (35B) scales above and below, eventually lost from the upper surface. Upright shrub, 2.0 × 1.1m in 15 years. Irregularly flowering twice a year (for a couple of months each time).

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: one of a series of plants of this cross raised at the former address of the registrant in Rangitoto Avenue, Remuera, New Zealand

(v) ‘**Rangitoto Glow**’

cv. *Lepidote rhododendron*: ‘Gardenia Odyssey’ (s) × *laetum*  
H: B.E. Oldham (1993), G: B.E. Oldham (1999), N: B.E. Oldham (2006), REG: B.E. Oldham (2009)  
Fls 9–13/flat truss, tubular funnel-shaped, 90 × 75mm, with 5 wavy-edged lobes, opening brilliant yellow (15C) at apex of lobes, shading to vivid yellow (16A) centrally, unmarked; outside vivid yellow (17C); not scented; style moderate yellow-green

(143D), stigma strong yellow-green (143C). Calyx very short, 5-lobed, strong yellow-green (143B). Lvs elliptic, 115 × 55mm, margins flat, acute at apex, broadly tapering at base, glossy green above, with light yellow-green (144D) scales when young (lost with age). Fastigiate shrub, 2.75 × 1.25m in 12 years. Repeat-flowering.

Nomenclatural standard: colour print provided by the registrant (WSY0112807). Etymology: derived from a cross made when the registrant was living in Rangitoto Avenue, Remuera, New Zealand

(v) **'Rangitoto Sunrise'**

cv. Lepidote rhododendron: 'Gardenia Odyssey' (s) × *laetum*

H: B.E. Oldham (1993), G: B.E. Oldham (1999), N: B.E. Oldham (2003), REG: B.E. Oldham (2009)

Fls 6–8/flat truss, tubular funnel-shaped, 60 × 75mm, with 5 wavy-edged lobes, buds deep red (185A), blending to moderate purplish red (59C), opening inside light orange-yellow (22C), margins deep yellowish pink (39B), generally streaked light pink (39D) into the tube and with patches of vivid reddish orange (42A) at the lobe sinuses; outside moderate reddish orange (35A), blending into vivid yellow-green (154A) at the base; slightly scented. Calyx very short, vivid red (45B). Lvs elliptic, 135 × 60mm, margins flat, acute at apex, broadly tapering at base, hairless. Fastigiate shrub, 1.7 × 1.1m in 12 years. Repeat-flowering.

Nomenclatural standard: colour print provided by the registrant (WSY0112809). Etymology: the cross was made in the registrant's previous garden in Rangitoto Avenue, Remuera, New Zealand

(v) **'Rangitoto Sunset'**

cv. Lepidote rhododendron: 'Gardenia Odyssey' (s) × *laetum*

H: B.E. Oldham (1993), G: B.E. Oldham (1999), N: B.E. Oldham (2000), REG: B.E. Oldham (2009)

Fls 7–9/flat truss, tubular funnel-shaped, 70 × 110mm, with 5 wavy-edged lobes, opening inside light yellow (17D) at the centre, shading to deep yellowish pink (44D) at the margins, outside

brilliant yellow (11A) in centre of the lobes, strong yellowish pink (37A) at the margins, the tube strong reddish orange (42B), blending to brilliant greenish yellow (4A) at the base; outside of the tube streaked vivid reddish orange (33B); sinuses sometimes blotched vivid reddish orange (34A); not scented. Calyx less than 1mm long, minutely 5-lobed, strong purplish red (61B). Lvs narrowly elliptic, 140 × 55mm, margins flat, acute at apex, tapered at base, with moderate numbers of dark greenish yellow (152C) scales when young, but these lost with age, glossy green above (turning dark purple (79A) in winter on old growth). Fastigiate shrub, 2.25 × 1.25m in 15 years. December to June.

Nomenclatural standard: colour print provided by the registrant (WSY0112810). Etymology: derived from crosses made by the registrant when living in Rangitoto Avenue, Remuera, New Zealand

(r) **'Reich's Signifikant'**

cv. Lepidote rhododendron: *ambiguum* (s) × 'Barnaby Sunset'

H: W. Reich (1990), G: W. Reich (1995), N: W. Reich (1999), I: W. Reich (2000), REG: W. Reich (2010)

Fls 7/open to loose ball-shaped truss, open funnel-shaped, 46–49 × 32–34mm, with 5 flat-edged lobes, greenish yellow in bud, opening light yellow throughout, with some darker marking on the dorsal lobe. Lvs narrowly to broadly elliptic, 15–50 × 7–22mm, pointed at apex, cuneate at base, margins flat, with a scaly indumentum. Shrub 1.7 × 1.2m in 16 years. Mid to late April.

Nomenclatural standard: colour print provided by the registrant (WSY)

(r) **'Reshma'**

cv. Elepidote rhododendron: 'Graf Lennart' (s) × 'Blewbury'

H: K.-H. Hübbbers (1987), G: K.-H. Hübbbers (1992), N: Mrs L. Hübbbers (1999), REG: J. Westhoff (2009)

Fls c.12/loose, flat-topped, rounded truss, funnel-campanulate, 35 × 40mm, with 5 wavy-edged lobes, inside greenish white (157D), outside pale yellow-green (157B), with deep purplish red (187D) spots

on two-thirds of the dorsal throat. Calyx minute, greenish, with a small red rim. Lvs elliptic, 60 × 25mm, acute at apex, cuneate at base, margins decurved, glossy above, hairless. Shrub 1.0 × 0.8m in 15 years. Early to mid April.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112992). Etymology: named for a niece of the raiser's wife

(r) **'Rhody Gus'**

cv. Elepidote rhododendron: parentage unknown  
G: Heritage Museums and Gardens (1970s),  
S: Sandwich Club (2009), N: Sandwich Club (2009), REG: Sandwich Club (2009)  
Fls 12/conical truss, open funnel-shaped, c.65 × 100mm, with 7 wavy-edged lobes, between two shades of light purple (75A–B) in bud, opening inside very pale purple (75D), with a 10mm vivid reddish purple (74A) mark in the throat, fading to nearly white with age, and with irregularly spaced spots of vivid to light reddish purple (74A–C) and moderate purplish pink (74D); outside very light purple (75C); stamens rudimentary; stigma strong purplish red (71B). Pedicels moderate reddish brown (166B). Lvs oblong, 150 × 65mm, obtuse at apex, oblique at base, margins wavy, semi-glossy and moderate olive green (147A) above, hairless. Shrub 2.7 × 2.1m in 40 years. Late May.

(r) **'Robert Hale'**

cv. Elepidote rhododendron: 'Van Nes Sensation' (s) × *falconeri*  
H: K.I. Sifferman (c.1969), G: Griswold Nursery (1980), N: Griswold Nursery (2010), REG: Griswold Nursery (2010)  
Fls 14/ball truss, funnel-campanulate, c.50 × c.75mm, with 5 wavy-edged lobes, light purplish pink (73C) in bud, opening greenish white (155C) throughout, with a moderate purplish red (184C) blotch at the base of the dorsal lobe, and tiny markings of the same colour extending upwards in straight lines from the edge of the blotch; anthers very dark red. Lvs oblong, 140 × c.70mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and dark green above, with moderate

yellow (161A) hairs below. Shrub with an open habit, 3.4 × 3.4m in 41 years. Mid May.

Etymology: named for a lawyer and long-time friend of the Griswold Nursery

(r) **'Romantika'**

cv. Elepidote rhododendron: parentage unknown  
H: K.-H. Hübbers (1988), G: K.-H. Hübbers (1995), N: K.-H. Hübbers (2000), REG: J. Westhoff (2009)  
Fls c.14/loose, roundish truss, broadly funnel-shaped, 50 × 80mm, with 6 very wavy-edged lobes, opening inside pale yellow to pale orange-yellow (19C–D) in the throat, deep purplish pink (55A) at the margins, the dorsal lobe darker orange-yellow and with two faint bands of brownish orange (165B) spots; outside deep purplish pink (55A). Calyx 2mm long, green, with a red rim. Lvs oblong, 100 × 45mm, emarginate and with a minute point at apex, rounded at base, margins slightly decurved, matt above, with silver hairs below when mature. Shrub 1.0 × 0.9m in 10 years, with white bark on older plants. April.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112990)

(r) **'Rose Tango'**

cv. Elepidote rhododendron: 'Crimson Constellation' (s) × 'Plum Passion'  
H: J. Barlup (2002), G: J. Barlup (2006), N: J. Barlup (2010), REG: J. Barlup (2010)  
Fls 17/ball truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, dark red (59A) in bud, opening inside very pale purple (76C), except for the dorsal lobe which is very pale purple (76D), all lobes edged strong purple (77B); outside very pale purple (76C), with darker midribs and edging of strong purple (77B); brilliant yellow-green (150C) spotting on dorsal lobe, beginning 25mm from the base and extending almost to the darker edging. Lvs elliptic, 110 × 55mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.6 × 0.9m in 7 years. Early May.

(r) **'Rosilac'**

cv. Elepidote rhododendron: 'Paprika Spiced' (s) × 'Deenah'

H: K.-H. Hübbbers (1993), G: K.-H. Hübbbers (2000), N: Mrs L. Hübbbers (2007), REG: J. Westhoff (2009)

Fls 10–16/roundish truss, broadly funnel-shaped, 35 × 50mm, with 7 very wavy-edged lobes, buds strong red (53C), opening inside light purplish pink (55C) in the throat, pale pink (49D) at the margins, outside strong red (51A), with light pink (50D) margins, dense deep red (53A) spots on three-quarters of the four dorsal lobes: the bicoloured effect strengthens as the flower ages. Calyx minute, light green. Lvs elliptic, 85 × 40mm, tapered apex, rounded at base, margins decurved, glossy above, hairless. Shrub 1.2 × 1.0m in 15 years. April. Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112988). Etymology: named after a suburb of San Fernando, Trinidad

(r) **'Roter Samurai'**

cv. Elepidote rhododendron: *makinoi* (s) × *yakushimanum* × *microgynum* Gymnocarpum Group

H: W. Reich (1990), G: W. Reich (2004), N: W. Reich (2009), REG: W. Reich (2010)

Fls up to 11/loose, ball-shaped truss, broadly funnel-campanulate, 38–41 × 50–55mm, with 5 very wavy-edged lobes, strong red (53B) throughout, with dark reddish brown spots on the three dorsal lobes. Calyx 6–7mm long, lobes reflexed, pinkish red. Leaves narrowly oblong-lanceolate, 28–127 × 8–30mm, pointed at apex, cuneate at base, margins decurved, young leaves hairy above but eventually glossy green and with impressed veins, with a whitish ochre indumentum below which ages to ochre orange. Shrub 0.6 × 0.6m in 9 years. May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: the name alludes to the colour of the flowers and to the Japanese origin of two of the parents species; samurai being the highest class of Japanese warrior

(v) **'Rowena Knight'**

cv. Lepidote rhododendron: *tuba* (s) × 'Rangitoto Rose'

H: B.E. Oldham (1993), G: B.E. Oldham (1997), N: B.E. Oldham (2000), REG: B.E. Oldham (2009)

Fls 9–11/flat truss, trumpet-shaped, the tube slightly curved, 100 × 55mm, with 5 slightly wavy-edged lobes, buds pale purplish pink (62D), with bract margins strong purplish red (58B), opening inside very pale purple (69C), rapidly fading to white through very pale purple (69D), with strong purplish pink (68B) spots at the lobes bases, outside similarly coloured with streaks of strong purplish pink (68B) at the base of the lobes; strongly scented. Calyx disc-like, 1.5mm long, irregularly lobed, strong purplish pink (68B). Lvs elliptic, 150 × 65mm, margins flat, matt moderate yellow-green (146C) above, acute to attenuate at apex, rounded at base, with sparse scales below (which become more sparse with age). Upright shrub, 1.7 × 1.5m in 15 years. Flowering mainly February to June (New Zealand), but occasional flowers at any period.

Nomenclatural standard: colour print provided by the registrant (WSY0112746). Etymology: named for the registrant's daughter-in-law

(r) **'Rustic Cloud'**

cv. Elepidote rhododendron: parentage unknown  
S: Sandwich Club (2009), G: Heritage Museums and Gardens (1970s), N: Sandwich Club (2009), REG: Sandwich Club (2009)

Fls 8/ball truss, broadly funnel-shaped, c.85 × 115mm, with 7 very wavy-edged lobes, moderate purplish pink (65A) in bud, opening inside very pale purple (76D), with a pale yellow-green (4D) centre, the dorsal lobes with linear, speckled markings of vivid reddish orange (34B); outside fades with age from moderate purplish pink (65A) to very pale purple (76D). Lvs elliptic, 115 × 45mm, broadly acute at apex, rounded at base, with flat margins, dull and moderate olive green (147A) above, hairless. Shrub 1.8 × 2.1m in 10 years. Late May.

(r) **'Sammy Brim'**

cv. Elepidote rhododendron: 'Tiana' (s) × 'Helen Vieira'

H: J.D. Brim (2003), G: J.D. Brim (2006),

N: J.D. Brim (2009), REG: J.D. Brim (2009)

Fls 14/ball truss, open funnel-shaped, c.65 × 90mm, with 7 wavy-edged lobes, greenish white throughout, with a large, central, dark red (59A) blotch, resembling the projection of a starburst firework explosion. Lvs oblong, 200 × 60mm, broadly acute at apex, rounded at base, margins flat, glossy and dark yellowish green (139A) above, hairless. Shrub 1.2 × 1.2m in 6 years. Early May. Etymology: named for the registrant's son

(r) **'Sangeet'**

cv. Lepidote rhododendron: 'Biskra' (s) × 'Conroy'

H: K.-H. Hübbers (1990), G: K.-H. Hübbers

(1997), N: Mrs L. Hübbers (2006),

REG: J. Westhoff (2009)

Fls up to 6/loose truss, funnel-shaped, 40 × 50mm, with 5(-6) flat-edged lobes, inside strong orange-yellow (24B), outside vivid reddish orange (40A), fading to light orange (26C) at the margins, with darker veins from the centre to the margin. Calyx 5mm long, violet, with green scales. Lvs lanceolate, 70 × 30mm, acute at apex, cuneate at base, margins flat, glossy above, with small yellowish scales when mature. Upright shrub, 1.6 × 1.2m in 15 years, with aromatic foliage. April.

Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112987). Etymology: the word "sangeet" in Hindi means "song"

(r) **'Satan's Cushion'**

cv. Elepidote rhododendron: *forrestii* Repens Group × 'Popocatepetl' (s) × *forrestii* Repens Group

H: B.N. Starling (1993), G: B.N. Starling (1999),

N: B.N. Starling (1999), I: B.N. Starling (2002),

REG: B.N. Starling (2009)

Fls 3-4/lax truss, tubular-campanulate, 40-65 × 30-35mm, with 5 wavy-edged lobes, buds vivid red (46B), opening vivid reddish orange (44B) both inside and out, unmarked. Calyx 1-2mm long, green. Lvs broadly elliptic, 50 × 20-25mm,

rounded at apex, slightly tapered at base, margins flat, matt green above, hairless. Very low and compact shrub, 0.3 × 1.0m in 15 years. Late March to early April.

Etymology: named for its habit and the fiery colour of the flowers ("hot as hell" according to the registrant)

(r) **'Scented Border'**

cv. Lepidote rhododendron: selection from *ciliatum*

G: Lord & Lady Falmouth (1965), S: Hon.

E. Boscawen (2000), N: Hon. E. Boscawen (2000),

I: Hon. E. Boscawen (2001), REG: Hon.

E. Boscawen (2009)

Fls 3-5/rather flattened, but rounded truss, often with groups of 5-8 trusses together, broadly tubular-campanulate, 38-50 × 30-55mm, with 5 emarginate, wavy-edged lobes, buds deep purplish pink (68A), to dark purplish pink (186C), fading to white at the base, opening inside white or white tinged pale pink, the outside blushed light purple/strong reddish purple (75B/70B) on the central portion of the lobes, stronger purplish pink on the three dorsal lobes (not as bright as deep purplish pink, 73A), fading almost to white with age; with a pleasant, delicate scent; stamens 10, filaments white, pubescent at base, anthers golden brown; style as long as the corolla, white, sometimes flushed pink at base and apex, stigma creamy yellow. Calyx irregular, 8-11mm long, strong yellow-green (145A), frequently flushed brownish red, with stiff marginal hairs. Pedicels strong yellow-green (145A), with spreading hairs. Lvs ovate or elliptic to obovate, 45-90 × 19-37mm, rounded or obtuse at apex, varying from obtuse to acuminate at apex, matt to slightly glossy green above, margins upcurved, ciliate and hairy (the hairs rufous to cream/white), lower surface with brown scales. Shrub 1.5-2.0 × 1.5-2.0m in 18 years. March to April. FCC 2009

Nomenclatural standard: herbarium specimen of the FCC plant (WSY0112836). Etymology: named for the border at Tregothnan where the selection was made

(r) **‘Shanti’**

cv. Lepidote rhododendron: ‘Biskra’ (s) × ‘Conroy’  
H: K.-H. Hübbbers (1990), G: K.-H. Hübbbers (1997), N: Mrs L. Hübbbers (2006),  
REG: J. Westhoff (2009)  
Fls up to 5/very loose truss, narrowly funnel-shaped, 50 × 50mm, with 5 wavy-edged lobes, buds moderate reddish orange (173B), opening inside strong yellowish pink (31C), outside vivid to strong reddish orange (34B–C), with darker veins from the centre to the margin. Calyx minute, greenish, with paler scales. Lvs oblong (curved like a banana), 60 × 35mm, mucronate at apex, rounded at base, margins wavy, glossy above, with brown scales when mature. Upright shrub 2.0 × 1.4m in 15 years. April.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112986). Etymology: the word “shanti” in Hindi means “peace”

(r) **‘Sharda’**

cv. Lepidote rhododendron: ‘Biskra’ (s) × ‘Conroy’  
H: K.-H. Hübbbers (1990), G: K.-H. Hübbbers (1997), N: Mrs L. Hübbbers (2006),  
REG: J. Westhoff (2009)  
Fls up to 6/flat, very loose truss, tubular funnel-shaped, 40 × 55mm, with 5 flat-edged lobes, greyish red (182B) in bud, opening inside deep yellowish pink (39B), outside vivid reddish orange to strong red (41A/B), with darker veins from centre to margin; slightly scented. Calyx 3mm long, brownish green, scaly. Lvs oblong-elliptic, 70 × 30mm, mucronate at apex, cuneate at base, margins flat, glossy above, with dense brown scales when mature. Upright shrub, 1.6 × 1.2m in 15 years, with slightly aromatic foliage. April.  
Nomenclatural standard: colour print of an electronic image provided by the registrant (WSY0112985). Etymology: named for a sister-in-law of the raiser’s wife

(r) **‘Silbermond’**

cv. Elepidote rhododendron: ‘Resonanz’ (s) × ‘Mrs A.T. de la Mare’  
H: H. Hesse (1998), G: H. Hesse (2006),  
N: H. Hesse, REG: H. Hesse (2009)

Fls 8–18/ball truss, open funnel-shaped, 45–60 × 50–85mm, with 6–7 wavy-edged lobes, buds light pink, opening inside white, with the margins more or less with a slight pink tinge, throat pale yellowish towards the base, with very faint brownish marking; outside white, tinged with pink towards the margins. Calyx 1–2mm long, light green. Lvs elliptic, 60–130 × 16–60mm, margins flat or decurved, matt green above, rounded to partly acute at apex, rounded at base. Shrub 0.75 × 0.6m in 10 years. Mid to late May.

Etymology: the name alludes to the colour of the flowers

(r) **‘Silver Frost’**

cv. Elepidote rhododendron: selection from *degronianum* subsp. *heptamerum* var. *kyomaruense*, grown from seed obtained by Greer from Koichiro Wada

S: H.E. Greer (1995), G: H.E. Greer (1972),  
N: H.E. Greer, I: Greer Gardens (1995),  
REG: H.E. Greer (2010)

Fls 15/ball truss, funnel-campanulate, 40 × 50mm, with 5 wavy-edged lobes, buds white, slightly tinged pink (lighter than pale yellowish pink, 36D), opening white inside, with a very light yellow-green spotting on the dorsal lobe; outside white; bud scales bronze green. Calyx minute, white. Lvs ovate, 150 × 45mm, acuminate at apex, rounded at base, margins downcurved, dull, deep green above, with a silvery white indumentum above which lasts into summer and an agglutinated indumentum below which is light tan when mature. Shrub with a dense habit, 1.5 × 2.4m in 25 years. Mid May.

The original seed was labelled as “R. metternianum” (now treated as ‘Metternianum’)

(r) **‘Sir Edmund Hillary’**

cv. Elepidote rhododendron: selection from *arboreum* subsp. *arboreum*  
S: Heritage Park Rhododendron Charitable Trust (2009), G: Heritage Park Rhododendron Charitable Trust (pre 1985), N: Heritage Park Rhododendron Charitable Trust (2009),  
REG: Heritage Park Rhododendron Charitable Trust (2010)

Fls 21–26/very compact, rounded truss, tubular-campanulate, 42–52 × 44–53mm, with 5 wavy-edged lobes, described as a mid orange-red (vivid reddish orange, 40B) in bud and throughout when open, with darker red spotting and darker red basal nectaries. Calyx 1–3mm long, insignificant, not coloured. Lvs oblanceolate, 80–140 × 35–40mm, margins flat, tapered at base, pointed at apex, glossy green above, with a thin, silvery white, plastered indumentum below; the leaves mostly displayed at an angle of 45° or less (unlike sister seedlings which have strongly drooping, paler green foliage). Shrub 3.5 × 2.5m in 40 years. August to October (New Zealand).

Raised from seed sent by the Queen of Nepal to the New Zealand Rhododendron Association about 40 years ago. Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the famous mountaineer, a New Zealander, with a strong connection to Nepal from whence the seed came

(r) **‘Sklerijenn’**

cv. Elepidote rhododendron: ‘Double Date’ (s) × ‘Karen Triplett’

H: L. Le Duigou (2000), G: A. Le Duigou (2006), N: L. & A. Le Duigou (2010), REG: L. & A. Le Duigou (2010)

Fls 8–9/ball truss, broadly funnel-shaped, 57 × 100mm, with 6–7 wavy-edged lobes, moderate red (47A) in bud, opening inside pale yellow (162D to 158A), outside moderate yellowish pink (179D) to deep pink (180D); all lobes with fine spotting of moderate red (179A); stamens 12–15; style strong yellow-green (144C). Calyx 5–20mm long, with spreading, unequal lobes, moderate orange-yellow (168D), edged moderate red (179A). Lvs elliptic, 120 × 40mm, cordate at base, acuminate at apex, matt green above, margins decurved. Compact, erect shrub, 1.5 × 1.2m in 9 years. May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: the epithet is the Breton word meaning a “source of light”

(r) **‘Sleeping Beauty’**

cv. Lepidote rhododendron: *cinnabarinum* subsp. *xanthocodon* Concatenans Group (s) × *keiskei* var. *ozawae* ‘Yaku Fairy’

H: B.N. Starling (1972), G: B.N. Starling (1976), N: B.N. Starling (1976), I: B.N. Starling (1980), REG: B.N. Starling (2010)

Fls 6–8/domed to spherical truss, funnel-shaped, 20 × 35–40mm, with 5 wavy-edged lobes, buds pale yellow (158A), opening inside and out pale yellow (158B), unmarked; stamens 10, up to 22mm long, anthers light brown to orange; style 20–25mm long, reddish. Calyx 3mm long, reddish. Lvs ovate to lanceolate, 50–60 × 20–25mm, pointed at apex, rounded at base, margins wavy. Shrub with a low, spreading habit, 0.75 × 1.8m in 30 years, the flowers often obscuring the foliage. April.

Etymology: refers to the Tchaikovsky ballet of this name and suggested by the habit of the plant

(r) **‘Snug Harbor’**

cv. Elepidote rhododendron: *macrophyllum* (s) × *insigne*

H: unknown, G: unknown, N: I.G. Snyder (2009), REG: I.G. Snyder (2009)

Fls 16/ball truss, open funnel-shaped, 50 × 75mm, with 5 very wavy-edged lobes, light reddish purple (74C) in bud, opening inside very light purple (75C) in the centre, the lobes edged strong reddish purple (70B) and the dorsal lobe spotted with moderate reddish brown (174A) and greyish reddish orange (174B); outside deep purplish pink (70C), with a strong purplish red (71B) stripe in the centre of each lobe. Lvs lanceolate, 125–165 × 40mm, acuminate at apex, cuneate at base, margins downcurved, moderate olive green (137A) above, with yellowish white (156D) hairs on the underside of the leaf when mature and on the stem. Shrub with an open habit, 2.1 × 2.7m in 10 years. Midseason.

(r) **‘So-horai’**

cv. Elepidote rhododendron: selection from *makinoi*  
S: E. Yoshimura (1986), G: E. Yoshimura (1986), N: K. Ohta (2009), REG: E. Yoshimura (2009)  
Fls double, 11(–12)/shallow, half-ball truss, broadly funnel-campanulate, 40 × 60mm, with a 5-lobed

outer and a 4-lobed inner corolla, the lobes smooth-edged or only very slightly wavy, buds deep pink, opening paler pink throughout, with deep pink spots on the dorsal side. Calyx 3–4mm long. Lvs linear-lanceolate, 120 × 25mm, margins slightly recurved, obtuse at base, acuminate at apex, glossy green above, with white hairs beneath when young, but hairless when mature. Shrub 2.0 × 2.0m in 30 years. Late April.

Nomenclatural standard: colour print provided by the registrant (WSY0112292). First raised as a seedling by Mr Yoshimura in 1979

(r) **‘Sonnenauge’**

cv. Elepidote rhododendron: ‘Golden Splendour’ (s) × ‘Edward Dunn’ × *auriculatum*  
H: K.-H. Hübbers (2000), G: K.-H. Hübbers (2010), N: K.-H. Hübbers (2010), REG: J. Westhoff (2010)  
Fls up to 10/lax domed truss, funnel-shaped, 90 × 100mm, with 6–7 wavy-edged lobes, buds light yellow (11B), the tips blushed pink, opening inside brilliant yellow (11A) in the throat, shading to pale yellow (11C) near the margin, outside light yellow (11B); dark red (187B) blotch in the throat; slightly scented. Calyx 40mm long, light yellow (11B). Lvs elliptic, 120 × 55mm, rounded and mucronate at apex, slightly auriculate at base, margins flat to slightly decurved, matt green above, hairless. Shrub 1.7 × 1.1m in 9 years. Mid to late June.  
Etymology: derived from the sun-like yellow flowers and the eye-like blotch (“sonnen” means sun in German, “auge” means eye)

(r) **‘Sooner’**

cv. Elepidote rhododendron: ‘Westbury’ × *hyperythrum*  
H: J.T. Thornton (1993), G: L.O. Miller, N: L.O. Miller, REG: L.O. Miller (2009)  
Fls 10/conical truss, broadly funnel-shaped, 50 × 70mm, with 5 wavy-edged lobes, pink in bud, opening inside pink, with heavy maroon stippling in the throat of the dorsal lobe, outside pink. Lvs oblong, 125 × 40mm, acute at apex, rounded at base, margins downcurved, glossy and dark green above, hairless. Dense shrub, 1.2 × 1.5m in 12 years. Mid April.

Etymology: named after the University of Oklahoma football team: the name refers back to the time of the late 19th century organised land grant programme in what is now Oklahoma State, when potential claimants lined up behind a starting line waiting for a gun to signal the start of a rush to a favoured plot of land. The “Sooners” were those who jumped the gun. The “Boomers” were those who waited for the gun

(r) **‘Split Up’**

cv. Elepidote rhododendron: ‘Furnivall’s Daughter’ × unknown  
H: T. Huisman (1985), G: T. Huisman (1991), N: T. Huisman (2009), I: T. Schröder (2010), REG: T. Huisman (2009)  
Fls 8–12/domed truss, open funnel-shaped, 45 × 40mm, often split to the base, with 5 flat-edged lobes, buds vivid red (57A), opening inside deep purplish pink (66C) to vivid purplish red (57C), outside vivid purplish red (66B) to deep purplish pink (66C), with moderate purplish pink (74D) between the darker ribs; both inside and out with a network of darker lines/veins; dorsal lobes with vivid reddish purple (74A) spots on a strong reddish orange (169A) ground. Calyx 2mm long, very pale purple (76D). Lvs lanceolate to oblong, 90–140 × 50–75mm, broadly acute at apex, oblique at base, margins flat, matt green above, hairless. Shrub 1.2–1.5m in 10 years. Early to mid May.  
Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0108037). Etymology: name suggested by the deeply divided corolla

(r) **‘Spring Fragrance’**

cv. Elepidote rhododendron: *yakushimanum* ‘Koichiro Wada’ × ‘Loderi Venus’ (s) × ‘Van Nes Sensation’  
H: F. Fujioka (1995), G: F. Fujioka (2002), N: F. Fujioka (2009), REG: F. Fujioka (2009)  
Fls 14/domed truss, broadly funnel-shaped, 75 × 115mm, with 6–8 wavy-edged lobes, buds moderate purplish pink (54D) at base, shading to pale purplish pink (56D) near the tip, opening inside pale purplish pink (56A), shading to pale purplish

pink (55D) near the throat, with a margin of c.12–15mm wide of pale purplish pink (56D) and with minimal spotting of strong reddish orange (169A) at the throat; outside pale purplish pink (56D), with streaks of moderate purplish pink (54D) radiating from the base; scented. Calyx minute. Lvs ovate, 125 × 60mm, broadly acute at apex, rounded at base, with wavy margins, semi-glossy and dark yellowish green (139A) above, hairless. Shrub 1.1 × 0.9m in 10 years. Late April.

(r) **‘SpringGold’**

cv. Lepidote rhododendron: ‘Saffron Prince’ (s) × *dalhousiae*

H: E.W. Moyles (1992), G: E.W. Moyles (1998), N: E.W. Moyles (2010), I: Enjoy Rhododendrons (2000), REG: E.W. Moyles (2010)

Fls 6/rounded truss with a flattened crown, open funnel-shaped, c.75–80 × c.85–90mm, with 5 wavy-edged lobes, cinnamon red in bud, opening inside medium saffron yellow, with occasional small, pale red or pink markings on the lobes and a basal blotch of dark saffron to red; outside medium saffron yellow. Calyx c.3mm long, pale green. Lvs oblong, 85–90 × 35–40mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and deep green above, with scales below. Shrub 1.2 × 1.2m in 10 years, the foliage entirely obscured by the flowers. Early April.

Syn: ‘The Winner’

The spelling as one word with the capital second “g” is deliberate

(v) **‘Stellar Flare’**

cv. Lepidote rhododendron: *suaveolens* (s) × *goodenoughii*

H: B.E. Oldham (1996), G: B.E. Oldham (2003), N: B.E. Oldham, REG: B.E. Oldham (2009)

Fls (13–)19(–21)/truss (initially erect, then spreading to half hanging), salverform, slightly curved, 70 × 35mm, with 5 wavy-edged lobes, opening inside pure white to pale pink (49D), outside very pale purple (73D) at apex shading through very pale purple (69A) to moderate purplish pink (74D) at the base; no markings; sweetly scented. Calyx negligible. Lvs elliptic, 105 ×

45mm, margins flat, obtuse at apex, broadly tapering at base, glossy green above (moderate olive green to moderate yellowish green, 137A–D), hairless. Shrub 1.4 × 0.75m in 12 years. Repeat-flowering with at least one intense burst of about 3 months’ duration.

Nomenclatural standard: colour print provided by the registrant (WSY0112812). Etymology: epithet descriptive of the huge stellate trusses covering the bush

(r) **‘Steredenn’**

cv. Elepidote rhododendron: ‘Kernéostic’ (s) × ‘Lem’s Monarch’

H: M. Colombel (1993), G: M. Colombel, N: M. Colombel, REG: M. Colombel (2009)  
Fls 13–16/full, shallowly domed truss, broadly funnel-shaped, 55–65 × 90–105mm, with 5 lightly wavy-edged lobes, buds strong purplish red (58B), opening inside nearly white with an obvious red marginal band, which fades to pale pink/pale purple, outside with many thin, pink, obvious veins; relatively inconspicuous green spots on the dorsal lobe, which turn to light yellow and with two, small, yellow marks in the throat; stamens 10, straight, unequal, filaments white (tinged pink apically), anthers light brown at first, later white; stigma light pink. Calyx 3–6mm long, light green, tinged pink. Lvs elliptic, 130–150 × 45–50mm, margins flat, matt green above, broadly acute at apex, rounded at base, hairless. Shrub 1.3 × 2.4m in 15 years. Late April to early May.

Nomenclatural standard: colour print provided by the registrant (WSY0112407). Etymology: the epithet means “star” in the Breton language

(r) **‘Stuart’s Eye Candy’**

cv. Elepidote rhododendron: ‘Naselle’ (s) × ‘Hachmann’s Polaris’

H: S. Imrie, G: S. Imrie (2001), N: S. Imrie (2009), I: Lake Tapps Rhododendron Nursery (2009), REG: S. Imrie (2009)

Fls 19/ball truss, broadly funnel-shaped, 50 × 90mm, with 7 very wavy-edged lobes, vivid red (57A) in bud, opening inside pale greenish yellow (1D) in the centre of the corolla and on the centre of

all lobes, with a broad margin of moderate purplish pink (66D); outside moderate purplish pink (66D); tiny speckles of strong greenish yellow (153B) on the dorsal lobe. Calyx *c.*40mm long, vivid purplish red (61C). Lvs elliptic, 130 × 50mm, acute at apex, rounded at base, margins flat, dull and dark green (136A) above, hairless. Shrub with a dense habit, 0.9 × 0.9m in 10 years. Early April.

(r) ‘**Stuart’s Treasure**’

cv. Elepidote rhododendron: ‘Lemon Dream’ × ‘Eunie’  
H: S. Imrie (2000), G: S. Imrie (2003), N: S. Imrie (2009), I: Lake Tapps Rhododendron Nursery (2009), REG: S. Imrie (2009)  
Fls 15/ball truss, broadly funnel-shaped, 50 × 80mm, with 5 wavy-edged lobes, brilliant greenish yellow (1A) in bud, opening throughout light greenish yellow (1C); with several short, vivid reddish orange (40A), petaloid stamens at the base in addition to some normal stamens. Calyx *c.*3mm long, strong yellow-green (43B). Lvs obovate, 115 × 65mm, obtuse at apex, cuneate at base, margins flat, dull and dark bluish green (131A) above, hairy below (the hairs greyish reddish orange (177C) on the mature leaf). Shrub with a dense habit, 1.2 × 0.75m in 10 years. Mid April.

(r) ‘**Sugan**’

cv. Lepidote rhododendron: ‘Biskra’ (s) × ‘Conroy’  
H: K.-H. Hübbers (1990), G: K.-H. Hübbers (1997), N: Mrs L. Hübbers (2009), REG: J. Westhoff (2009)  
Fls *c.*4/loose truss, tubular funnel-shaped, 50 × 45mm, with 5 flat-edged lobes, opening inside deep pink (48A/B), outside vivid reddish orange to strong red (41A–C), with darker veins from centre to margin. Calyx 3mm long, light green. Lvs lanceolate, 80 × 35mm, mucronate at apex, cuneate at base, margins decurved, glossy above, with light brown scales when mature; slightly aromatic. Upright shrub, 2.2 × 1.4m in 15 years. April.  
Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0112980). Etymology: the word “sugan” in Hindi means “beautiful” and is also used as a female first name

(r) ‘**Sumari**’

cv. Elepidote rhododendron: ‘Good News’ × ‘Erato’ (Hachmann) (s) × *Fortunei* subsp. *discolor* hybrid  
H: K.-H. Hübbers (1994), G: K.-H. Hübbers (2001), N: Mrs L. Hübbers (2007), REG: J. Westhoff (2009)  
Fls up to 8/loose truss, funnel-shaped, 60 × 80mm, with 5 wavy-edged lobes, strong red (46A) in bud, opening inside vivid red (45C), outside vivid reddish orange (44C), with faint, darker red spots on the dorsal part of the throat: colour does not fade or burn in sunlight. Calyx up to 6mm long, lobed, red, with greenish scales. Lvs elliptic, 120 × 60mm, obtuse but minutely apiculate at apex, truncate at base, margins flat, matt above, hairless. Shrub 1.1 × 0.8m in 10 years. Late May. Reported to be resistant to bud blast.  
Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0112979). Etymology: named for the mother of the raiser’s wife

(r) ‘**Sun Fire**’

cv. Elepidote rhododendron: ‘Rotgold’ × ‘Amaretto’ × *dichroanthum*, yellow form  
H: T. Huisman (1993), G: T. Huisman (1998), N: T. Huisman (2009), I: T. Schröder (2010), REG: T. Huisman (2009)  
Fls 8–10/ball truss, open funnel-shaped, 50–65 × 65–80mm, with 5 wavy-edged lobes, the dorsal lobes curved and bent backwards, buds vivid reddish orange (40B), opening pale yellow (12D) to almost white (N155D), with a broad, 8–10mm wide margin of strong red (41B), outside with broad veins of pale greenish yellow (2D), with deep pink (48B) at the margins; many spots of strong orange (163B) over the dorsal lobes. Calyx 1.5–3.5mm long, a mixture of light greenish yellow (4C) to light orange (28D). Lvs elliptic, 75–110 × 30–40mm, acute at apex, cuneate at base, margins flat, glossy green above, hairless. Shrub 1.2 × 1.4m in 10 years. Early to mid May.  
Nomenclatural standard: colour print of an electronic image provided by the raiser (WSY0108038). Known as seedling no. Z 515.  
Etymology: name suggested by the flowers appearing bright orange at a distance

(r) **‘Sun Silk’**

cv. Elepidote rhododendron: ‘Command Performance’ (s) × ‘Goldprinz’

H: J. Barlup (2003), G: J. Barlup (2007), N: J. Barlup (2010), REG: J. Barlup (2010)

Fls 24/ball truss, broadly funnel-shaped, 60 × 85mm, with 7 wavy-edged lobes, light orange-yellow (16C) in bud, opening inside pale yellow (8D) with the dorsal lobe a deeper shade of light greenish yellow (8B); outside colouring similar; discrete patches of strong red (53C) spots inside on each of the two dorsal lobes extending 40mm from the base, and brush-like strong red (53C) flares on the dorsal lobes, extending c.10mm from the base. Calyx c.10mm long, light greenish yellow (8D), with a touch of strong red (53C) at the edge. Lvs elliptic, 115 × 55mm, broadly acute at apex, rounded at base, margins flat, glossy and moderate olive green (147A) above, hairless. Shrub with an open habit, 0.9 × 0.6m in 6 years. Early May.

(r) **‘Superflimmer’**

cv. Elepidote rhododendron: sport from ‘Goldflimmer’

H: S. Imrie, G: S. Imrie (2002), N: S. Imrie (2009), I: Lake Tapps Rhododendron Nursery (2010), REG: S. Imrie (2009)

Fls 9/ball truss, saucer-shaped, 35 × 50mm, with 5 wavy-edged lobes, moderate purplish red (70A) in bud, opening inside light purple (80D), with a speckled, deep greenish yellow (153D) dorsal blotch; outside light purple (80C). Lvs elliptic, 115 × 45mm, broadly acute to acute at apex, cuneate at base, margins upcurved, dull and dark bluish green to dark green (131A–B) above and variegated with prominent light to pale yellow (8B–D) markings, hairless. Shrub with a dense habit, 0.9 × 0.9m in 10 years. Late April to May.

(r) **‘Susie Lynn Jones’**

cv. Elepidote rhododendron: ‘Countess of Lockington’ (s) × ‘Australian Sunset’

H: D.J. Dosser (2004), G: D.J. Dosser (2008), N: D.J. Dosser (2010), REG: D.J. Dosser (2010)

Fls 12/ball truss, funnel-shaped, 40 × 100mm, with 6 wavy-edged lobes, buds moderate red (47A),

opening inside pale yellow (11D), with strong purplish red (58C) margins, outside pale yellow (11D), with a pink tinge; deep purplish red (71A) ray on the two dorsal lobes; slightly scented. Calyx 14mm long, strong red (47B). Lvs lanceolate, 150 × 25mm, broadly acute at apex, cuneate at base, margins flat, glossy green above, hairless. Shrub 0.75 × 0.5m in 6 years. November (Australia).  
Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for a friend of the registrant

(r) **‘The Pink Ribbons’**

cv. Elepidote rhododendron: ‘Paprika Spiced’ (s) × ‘Jezebel’

H: E. Cockeram (1995), G: E. Cockeram (2000), N: E. Cockeram (2009), I: Log Cabin Nursery (2010), REG: E. Cockeram (2009)

Fls 10–12/domed truss, funnel-campanulate, 60 × 85mm, with 5 wavy-edged lobes, opening inside light purplish pink (55C), with sparsely distributed speckling of deep purplish pink (55A); outside strong purplish pink (55B). Calyx less than 2mm long, brilliant greenish yellow (5A). Lvs ovate, 125 × 35mm, acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive green (137A) above, hairless. Shrub with a dense habit, 1.4 × 1.2m in 14 years. Mid May.

Etymology: the epithet relates to the two pink ribbons used as a logo by the Susan G. Komen Foundation, a national organisation in the USA supporting breast cancer research; funds derived from the sale of plants of this cultivar are being donated to this cause

(v) **‘Thirtieth of June’**

cv. Lepidote rhododendron: *leucogigas* × *konori* × *laetum*

H: P. Sullivan (1973), G: P. Sullivan (1979), N: P. Sullivan (c.1979), I: Pacific Island Nursery (2004), REG: R. Kruppa (2010)

Fls 7/flat truss, elongated tubular funnel-shaped (with a straight tube), 125 × 100mm, with 7 wavy-edged lobes, reddish purple in bud, opening off-white (close to 159D) throughout, with streaks of moderate to light and pale pink (close to 49B–D)

on the outside of the tube; with a small, strong pink (49A) spot at each lobe sinus; anthers white, filaments very pale pink; stigma very light green, style mainly white except for red immediately below the stigma. Lvs 170–175 × 70–75mm, broadly acute at apex, cuneate at base, margins upcurved, glossy and moderate olive green (147A) above, with no visible indumentum. Very slow-growing shrub, with an open, upright habit, 0.7 × 0.4–0.5m in 6 years. May to June.

Etymology: the name commemorates the first day the plant flowered (in San Francisco); the flowering period given relates to where the registration description was taken (Hawaii)

(r) ‘**Thomas Le Duigou**’

cv. Elepidote rhododendron: ‘Double Date’ (s) × ‘Karen Triplett’

H: L. Le Duigou (2000), G: A. Le Duigou (2005), N: L. & A. Le Duigou (2009), REG: L. & A. Le Duigou (2009)

Fls 10/rounded truss, open funnel-shaped, 70 × 120mm, with 6 wavy-edged lobes, deep yellowish pink (39B) in bud, opening inside light greenish yellow (1C), with greenish spots on all the lobes and around the centre of the flower (heavier on the dorsal lobes), outside light greenish yellow (3D), the lobes soon fade to white around a broad margin; stamens 14, anthers pale pink; style greyish yellow-green (148C), stigma green. Calyx 2–5mm long, moderate yellow-green (146D). Lvs elliptic, 120 × 45mm, margins decurved, cuspidate at apex, obtuse at base, hairless. Shrub 1.2 × 0.7m in 8 years. May.

Nomenclatural standard: colour print provided by the registrant (WSY0112425). Etymology: named for the grandson of the registrant

(v) ‘**Tickety Boo**’

cv. Lepidote rhododendron: ‘Saxon Glow’ (s) × ‘Vladimir Bukovsky’

H: D. & P.J. Brown (2002), G: D. & P.J. Brown (2007), N: D. & P.J. Brown (2009), I: Browns Nursery, REG: D. & P.J. Brown (2009)

Fls 5–6/lax truss, tubular funnel-shaped, 30 × 45mm, with 5 flat-edged lobes, buds strong yellowish pink (33C), opening inside vivid reddish orange

(40B), shading to light orange (28C) at the base of the corolla, outside strong yellowish pink (32C), shading to vivid red (45C) on the lobe margins, unmarked. Calyx vestigial. Lvs ovate, 75 × 43mm, rounded at base, acute at apex, margins flat, glossy green above, hairless. Compact, much-branched shrub 1.2 × 1.0m in 10 years. Repeat-flowering. Nomenclatural standard: colour print provided by the registrant (WSY01122253)

(r) ‘**Tisa**’

cv. Elepidote rhododendron: ‘Nancy Evans’ (s) × ‘Bambi’ × *proteoides*, no. 12

H: J. Barlup (2000), G: J. Barlup (2004),

N: C.A. Lehmann (2008), REG: J. Barlup (2009)

Fls 13/domed truss, funnel-shaped, 40 × 50mm, with 5 wavy-edged lobes, vivid red (46C) in bud, opening inside light yellow (20B), with edges of lobes lightly tinged strong red (53D); outside light yellow (20B), with tones of strong red (53D) on the midrib and deep red (53A) veining on the midrib; deep red (53A) spotting on the dorsal lobe extending 40mm from the base and deep red (53A) nectar pouches extending c.5mm from the base. Calyx c.6mm long, light yellow (20B), with deep red (53A) spotting. Lvs elliptic, 65 × c.30mm, acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.25 × 0.4m in 4 years. Late April.

(r) ‘**Tom Smith**’

cv. Elepidote rhododendron: parentage unknown

H: unknown, G: T. Smith (2008), N: S. Lord

(2009), REG: S. Lord (2009)

Fls 8–9/domed truss, broadly funnel-shaped, 75 × 90–100mm, with 6 wavy-edged lobes, deep peachy pink in bud, opening inside creamy peachy pink, with small, deep pink speckles scattered randomly on all lobes, and a midvein on each lobe consisting of a straight line of deep pink speckles, longest on the dorsal and shorter on other lobes; outside peachy orange-pink. Lvs elliptic, 125–150 × 65–75mm, obtuse at apex, rounded at base, margins flat, semi-glossy above, hairless. Shrub with an open habit, 1.4m in c.10 years. May.

Etymology: named for the father of the registrant

(r) **‘Tony Magallon’**

cv. Elepidote rhododendron: ‘Odee Wright’ (s) × ‘Crest’

H: D.A. Griswold (1969), G: Griswold Nursery (1980), N: Griswold Nursery (2010), REG: Griswold Nursery (2010)

Fls 8/lax truss, funnel-campanulate, *c.*50 × *c.*65mm, with 5 wavy-edged lobes, brilliant orange-yellow (21B) in bud, opening light yellow-green (2C) throughout; anthers brown. Lvs oblong, 65 × 40mm, broadly acute at apex, cordate at base, with upcurved margins, dull and moderate olive green (137B) above, hairless. Shrub with an open habit, 2.1 × 1.8m in 41 years. Mid May.

Etymology: named for a long-time employee of the Griswold Nursery

(r) **‘Tough Buck’**

cv. Elepidote rhododendron: unknown (s) × *hyperythrum*

H: J.T. Thornton (1995), G: L.O. Miller (2004), N: L.O. Miller (2004), I: Briggs Nursery, REG: L.O. Miller (2009)

Fls 11/conical truss, broadly funnel-shaped, *c.*40 × 75mm, with 5 wavy-edged lobes, red in bud, opening inside light pink and fading to white, with heavy maroon stippling on the dorsal lobe; outside light pink, fading to white. Lvs oblong, 100 × 50mm, acute at apex, rounded at base, margins downcurved, semi-glossy and dark green above, hairless. Dense shrub, 1.2 × 1.5m in 12 years. Mid April.

Etymology: named to reflect the nature of the plant growing in an inhospitable location

(r) **‘Treuscoat’**

cv. Elepidote rhododendron: ‘Lord Roberts’ (s) × ‘Karen Triplett’

H: L. Le Duigou (2000), G: A. Le Duigou (2006), N: A. Le Duigou (2009), REG: L. Le Duigou (2009)

Fls 19/rounded truss, broadly funnel-campanulate, 55 × 80mm, with 5–6, slightly wavy-edged lobes, strong purplish red (58B) in bud, opening inside light pink (49C), outside strong purplish pink (55B), with sparse purplish red spots on the dorsal lobe and a vivid reddish orange (30B) suffusion at the base of

the throat especially on the dorsal side; stamens 12; style deep yellowish pink (39B). Calyx 10–30mm long, pale yellow (20D). Lvs narrowly elliptic, 140 × 48mm, margins wavy and decurved, acute at apex, obtuse at base, hairless. Shrub 0.95 × 1.25m in 8 years. April to May.

Nomenclatural standard: colour print provided by the registrant (WSY0112395). Etymology: named for the village in Brittany, France, where the plant has been grown

(r) **‘Trewithen Purple’**

cv. Lepidote rhododendron: parentage unknown - probably a *russatum* hybrid

R: Trewithen Gardens (pre 1988), N: Trewithen Gardens, I: Trewithen Gardens, REG: Trewithen Gardens (2010)

Fls up to 7/individual truss, with up to 7 trusses clustered in a group, broadly funnel-shaped, *c.*18 × 28mm, 5-lobed, overall near strong to light purple (N80B–D), with on the outside near reddish purple/purplish pink (N74) at the base, extending up the lobes and fading out; inside occasionally with a few fawn speckles which appear usually at the junctions of a couple of petals, but their position seems variable; stamens 10, filaments near strong purple (N80B), anthers brown, not exerted; style white to near strong red (51A), stigma red. Calyx *c.*1mm long with unequal lobes, white scaly, like the ovary and pedicels. Lvs elliptic, up to 23 × 10mm, prominently scaly, the scales brown on the lower surface and white above; petioles with brown scales. Shrub 1.0–1.5 × 1.0–1.5m in 10 years. Late April to May (with a second flush in September/October). AM 2009

(r) **‘Trina’**

cv. Elepidote rhododendron: ‘Coral Blossom’ × ‘Tia’  
H: J. Barlup (2000), G: J. Barlup (2005), N: J. Barlup (2008), REG: J. Barlup (2009)

Fls 16/domed truss, broadly funnel-shaped, 50 × 70mm, with 6 wavy-edged lobes, vivid red (46A) in bud, opening inside pale orange-yellow (16D), with the dorsal lobe a slightly darker shade of light orange-yellow (16C); outside pale orange-yellow (16D), with a light pink (48C) tone in the centre of

the lobes and deep pink (48A) veining on the midribs; a strong red (53B) solid flare on the dorsal lobe, extending c.15mm from the centre and changing to spotting of the same colour for another c.15mm. Calyx 10mm long, pale orange-yellow (16D), with a strong red (53B) flare. Lvs elliptic, 125 × 45mm, broadly acute at apex, rounded at base, with flat margins, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 0.9m in 8 years. Early May.

(r) **‘Ute Syc’**

cv. *Elepidote rhododendron*: ‘Razorbill’ × *dendrocharis*

H: W. Reich (2001), G: W. Reich (2005),  
N: W. Reich (2007), REG: W. Reich (2010)  
Fls 20–30/ball truss, campanulate to tubular funnel-shaped, 17–20 × 24–27mm, with 5 flat-edged lobes, buds white, with strong purplish red (71B–C) tips, opening white, with strong purplish red (71B–C) edges; unmarked. Calyx 4mm long, brownish green. Pedicels brownish green. Lvs broadly elliptic, 21–42 × 6–14mm, obtuse at apex, broadly cuneate at base, margins flat but leaf convex, glossy green above, with a scaly indumentum. Upright shrub, 57 × 29cm in 6 years. Early to mid May.

Nomenclatural standard: colour print provided by registrant (WSY). Etymology: named for the registrant’s sister-in-law

(r) **‘Vanilla Torch’**

cv. *Elepidote rhododendron*: ‘Capistrano’ (s) × ‘Tia’

H: J. Barlup (2000), G: J. Barlup (2004),  
N: J. Barlup (2010), REG: J. Barlup (2010)  
Fls 13/ball truss, broadly funnel-shaped, 45 × 70mm, with 5 wavy-edged lobes, deep pink (48A) in bud, opening inside and out pale yellow (11C), blending to pale yellow (4D) at the margins; strong red (53C) flares on all lobes extending c.25mm from the base, much more pronounced on dorsal lobes. Lvs elliptic, 100 × 40mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 0.8 × 0.9m in 9 years. Early May.

(r) **‘Variegated Unique’**

cv. *Elepidote rhododendron*: sport from ‘Unique’

S: S.T. Imrie (2000), G: S.T. Imrie (2003),  
N: S.T. Imrie, I: Lake Tapps Rhododendron Nursery, REG: S.T. Imrie (2009)  
Fls 12/ball truss, broadly funnel-shaped, 45 × 75mm, with 5 wavy-edged lobes, light purplish pink (62C) in bud, opening inside pale greenish yellow (2D), at first lightly tinged pink which fades to 2D with age, with randomly distributed, inconspicuous pink flecks; outside brilliant greenish yellow (2B). Calyx c.3mm long, light yellow (12C). Lvs elliptic, 115 × 50mm, broadly acute at apex, rounded at base, margins irregularly wavy, dark yellowish green (136B) above, variegated with a distinctive, continuous, pale yellow (160D) margin, hairless. Shrub with a dense habit, 0.9 × 0.9m in 10 years. Mid April.

(r) **‘Vicky J.’**

cv. *Elepidote rhododendron*: *wardii*, open-pollinated

S: G. Masters (2010), G: E. Jago (2000),  
N: G. Masters (2010), REG: G. Masters (2010)  
Fls 11/rounded to loose truss, open funnel-shaped, 50 × 30mm, with 7 wavy-edged lobes, buds primrose yellow, opening inside primrose yellow with a tinge of green at the base and with a blotch made up of light mustard yellow spots, outside yellow at base, shading to lemon at the apex; anthers mid brown; slightly scented. Calyx 2mm long, green. Pedicels tinged red. Lvs broadly elliptic, 80–130 × 40–50mm, margins slightly upcurved, very broadly acute at apex, hairless. Shrub 1.8 × 0.9m in 14 years. Early to mid November (New Zealand).

Nomenclatural standard: colour print provided by registrant (WSY). Etymology: named for a daughter of Erica Jago

(r) **‘Violettes Impériales’**

cv. *Lepidote rhododendron*: ‘Vibrant Violet’ (s) ×

*calostrotum* subsp. *riparium*

H: M. Colombel (2002), G: M. Colombel (2006),  
N: M. Colombel (2010), REG: M. Colombel (2010)  
Fls 3–4/flat truss, saucer-shaped (almost flat), 50–55 × 50–55mm, with 5 flat-edged lobes, buds

deep reddish purple (77A), opening inside deep to strong purple (83B–C), outside vivid violet (88A), with a few purple spots on the dorsal lobe; stamens 10, hairy at the base, filaments coloured as the corolla, anthers black; style and stigma dark red at first, aging to light purple. Calyx 6–8mm long, light green, the lobes with scales in the centre. Lvs elliptic, 25–30 × 13–15mm, margins flat, broadly acute at apex, rounded at base, matt green above, scales glaucous at first, aging to brown with a darker centre and placed their own diameter apart. Shrub 0.75 × 1.1m in 8 years. Mid to late April.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: the epithet is the name of a light opera by Vincent Scotto and alludes to the colour of the flowers

(r) **‘White Rays’**

cv. Lepidote rhododendron: ‘Dora Amateis’ × *moupinense*

H: W. Reich (1995), G: W. Reich (1999–2000), N: W. Reich (2006), REG: W. Reich (2010)  
Fls 5/truss, radiating in a whorl, tubular funnel-shaped, 40 × 35–38mm, with (3–4–)5 smooth-edged, lobes which sometimes reach to the base of the corolla (the deepest lobes being much narrower), white in bud and throughout on opening. Calyx 4.5mm long, whitish green. Lvs lanceolate, 15–55 × 5–16mm, pointed at apex, cuneate at base, margins flat, slightly glossy above, with a scaly indumentum; some slightly twisted along long axis. Shrub 0.2 × 0.35m in 11 years. Late April to early May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: derived from the impression given by the radiating white flowers

(r) **‘Wilma Hentschel’**

cv. Lepidote rhododendron: ‘Razorbill’ (s) × ‘Laurin’

H: W. Reich (1996), G: W. Reich (2002–2003), N: W. Reich (2007), I: Glendoick Gardens (2010), REG: W. Reich (2010)

Fls 5–6/loose, open truss, funnel-campanulate, 35–44 × 32–36mm, with 5 wavy-edged lobes, light purple (81D) in bud, opening vivid to light purple (81B–C) throughout, unmarked. Calyx 5mm long,

whitish green. Lvs elliptic, 16–42 × 7–19mm, pointed at apex, cuneate at base, margins flat, glossy green above, with a scaly indumentum. Shrub 0.7 × 0.7m in 7 years. Late April to mid May.

Nomenclatural standard: colour print provided by the registrant (WSY). Etymology: named for the mother of the registrant’s daughter-in-law

(r) **‘Winsome Winnie’**

cv. Elepidote rhododendron: ‘Dexter’s Honeydew’ (s) × ‘Virginia Richards’ × *yakushimanum*, Exbury form

H: D. Mac Mullan, G: D. Mac Mullan, N: D. Mac Mullan (2010), REG: D. Mac Mullan (2010)  
Fls 10/ball truss, funnel-shaped, c.80 × 75mm, with 5 wavy-edged lobes, ivory in bud, opening pure white throughout, with a 25mm wide scarlet marking in the centre of the throat. Lvs elliptic, 100 × 40mm, broadly acute at apex, rounded at base, margins flat, mid green above, hairless. Shrub 0.9 × 1.1m in 8 years. Mid May.

Etymology: named for Winifred Minahan, wife of Joe Minahan who is a friend of the registrant

(r) **‘Yellow Balloon’**

cv. Elepidote rhododendron: ‘Ray Kruse’s Big Cream’ × *dichroanthum* (s) × ‘Janet Blair’ × ‘Autumn Gold’

H: D. Mac Mullan, G: D. Mac Mullan, N: D. Mac Mullan (2010), REG: D. Mac Mullan (2010)  
Fls 13/ball truss, broadly funnel-shaped, c.75 × 120mm, with 6 wavy-edged lobes, deep orange/yellow in bud, opening inside moderately strong yellow in the centre, shading to light yellow at lobe edges, outside deep yellow at base, fading to mid yellow at edges; unmarked. Lvs broadly elliptic, 130 × 50mm, broadly acute at apex, rounded at base, margins downcurved, hairless. Shrub 1.5 × 1.5m in 7 years. Late May.

(r) **‘Yolanta’**

cv. Lepidote rhododendron: *hanceanum* ‘Canton Consul’ (s) × *keiskei* var. *ozawae* ‘Yaku Fairy’

H: B.N. Starling (1971), G: B.N. Starling, N: B.N. Starling, I: B.N. Starling, REG: B.N. Starling (2009)

Fls 3–4/truss, funnel-shaped, c.20 × 30mm, with 5 wavy-edged lobes, buds light greenish yellow (3D), opening pale yellow-green (4D) inside and out, unmarked. Calyx c.5–7mm long, green. Lvs ovate to elliptic, 35–50 × 20–25mm, acute to bluntly pointed at apex, narrowly rounded at base, margins flat, matt green above, with scattered (non-touching), pale scales below. Very compact, floriferous shrub, 0.6 × 0.6m in 10 years, with reddish dormant flower buds. April.  
Etymology: named after a heroine of a Tchaikovsky opera

(v) **‘Yoshi and Marge’**

cv. Lepidote rhododendron: *bellwiggii* × (*javanicum* × ‘Duchess of Edinburgh’) (s) × *christi*  
H: R.A. Mitchell (1997), G: R.A. Mitchell & Pacific Island Nursery (2004), N: R.A. Mitchell (2010), REG: R.A. Mitchell (2010)  
Fls 7/lax truss, elongated tubular funnel-shaped, 75 × 65–70mm, with 5 flat-edged lobes, opening inside and out vivid reddish orange (43A); unmarked; filaments pink, anthers black with a white tip; style red, stigma dark red. Lvs elliptic, 140 × 60mm, acuminate at apex, cuneate at base, margins slightly upcurved, glossy and dark green (135A) above, moderate yellow-green (146C) below with a light

reddish purple centre vein, no visible indumentum. Shrub 0.7 × 0.5m in 13 years. March and July.  
Etymology: named for the husband and wife (Yoshi and Marge Watanabe) who were two of the founders of the Hawaii Chapter of the American Rhododendron Society

(r) **‘Zauberregen’**

cv. Elepidote rhododendron: ‘Orangeola’ (s) × ‘Karibia’

H: H. Hesse (1998), G: H. Hesse (2008), N: H. Hesse, REG: H. Hesse (2010)

Fls 8–13/ full, rounded truss, open funnel-shaped, 40–57 × 60–80mm, with 5–6 wavy-edged lobes, inside light to pale yellow, aging to pale yellowish pink, outside light to pale yellow changing to pale yellowish pink; striking, large, red-brown blotch on the dorsal lobe, all the other lobes speckled with many red-brown dots. Calyx 3–5mm long, pale green. Lvs elliptic to narrowly elliptic, 56–116 × 23–41mm, rounded at apex, narrowed/partially rounded at base, matt to slightly glossy green above, margins decurved, hairless. Very compact shrub, 0.75 × 0.8m in 11 years. Early May.

Etymology: translates as “magic rain” which alludes to the strong speckling over all the lobes)

## Names and addresses

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

\* = deceased

Ahern, T.

Bethlehem, Pennsylvania, USA

Baines, R.

Logan Botanic Garden, Stranraer, Wigtownshire,  
Scotland

Barlup, J.

Bellevue, Washington, USA

Baumschule Neumann

Berlin, Germany

Baumschule Stöckmann

Bad Zwischenahn-Rostrup, Germany

Bayport Plant Farm,

Lunenburg County, Nova Scotia, Canada

Becales, J.

Glen Mills, Pennsylvania, USA

Bertelmann, S.

Kea'au, Hawaii, USA

Bleuzen, Mme M.T.

Briec, France

Boscawen, Hon. E.

Tregothnan, Truro, Cornwall, England

\*Bovee, R.M.

Portland, Oregon, USA

Bovees Nursery, The

Portland, Oregon, USA

Briggs Nursery

Olympia, Washington, USA

Brim, J.D.

Mount Airy, North Carolina, USA

Brouse Nursery

Norristown, Pennsylvania, USA

Brouse, F.W.

Brouse Nursery, Norristown, Pennsylvania, USA

Brown, D. & P.J.

Browns Nursery, Tauranga, New Zealand

Browns Nursery

Tauranga, New Zealand

Clancy, B.

Bentleigh, Victoria, Australia

Cockeram, E.

Florence, Oregon, USA

Coker, B.J. & H.

Christchurch, New Zealand

Coleman, J.H.

Lithonia, Georgia, USA

Colombel, M.

Menez-Rohou, Fouesnant, France

Cook, J.W.

Whiteville, North Carolina, USA

Cox, K.N.E.

Glendoick Gardens, Glencarse, Perthshire, Scotland

Cummins Garden

Marlboro, New Jersey, USA

Danneil, H.-J.

Heidelberg, Germany

Davies, Mrs S.E.

Palmerston North, New Zealand

Dosser, D.J.

Lower Wilmot, Tasmania, Australia

Dover Nursery

Sandy, Oregon, USA

Enjoy Rhododendrons

Occidental, California, USA

Falmouth, Lord & Lady

Tregothnan, Truro, Cornwall, England

Fujioka, F.

Freeland, Washington, USA

Gilbert, R.J.

Bodmin, Cornwall, England

Glendoick Gardens

Glencarse, Perthshire, Scotland

Goheen, D.W.

Camus, Washington, USA

\*Gordon, I.F.

Taihape, New Zealand

Greer Gardens

Eugene, Oregon, USA

Greer, H.E.

Eugene, Oregon, USA

Griswold Nursery

Kirkland, Washington USA

Griswold, D.A.

Griswold Nursery, Kirkland, Washington, USA

Grov, K.R.  
Stord, Norway

Hagan, K.  
Kingston Beach, Tasmania, Australia

Hazelwood Gardens  
Newcastle, Washington, USA

Heritage Museum and Gardens  
Sandwich, Massachusetts, USA

Heritage Park Rhododendron Charitable Trust  
Kimbolton, New Zealand

Hesse, H.  
Hamersen, Germany

\*Heyduck, G.  
Bellevue, Washington, USA

Hindle, J.R.  
Kannapolis, North Carolina, USA

Holbrook, R.  
Charlotte, North Carolina, USA

Hübbers, K.-H. & Mrs L.  
Kranenburg, Germany

Huisman, T.  
Dalfsen, the Netherlands

Imrie, S.T.  
Bonney Lake, Washington, USA

Jago, E.  
Stratford, New Zealand

Jørgensen, P.M.  
Det Norske Arboret, Hjellevstad, Norway

Justice, C.L.  
Vancouver, British Columbia, Canada

Justice, D.  
New Westminster, British Columbia, Canada

Kesterson, G.H.  
Freeland, Washington, USA

Klimavicz, J.F.  
Vienna, Virginia, USA

Krebs, S.L.  
The Holden Arboretum, Kirtland, Ohio, USA

Kruppa, R.F.  
Volcano, Hawaii, USA

Lake Tapps Rhododendron Nursery  
Bonney Lake, Washington, USA

Le Duigou, A. & L.  
Raden, Scaër, France

\*Leach, D.G.  
North Madison, Ohio, USA

Lehmann, C.A.  
Klampenborg, Denmark

Log Cabin Nursery  
Springfield, Oregon, USA

Logan Botanic Garden  
Stranraer, Wigtownshire, Scotland

Lord, S.  
Linkwood, Maryland, USA

Mac Mullan, D.  
Hamburg, Pennsylvania, USA

Marques, R.  
Kea'au, Hawaii, USA

Masters, G.  
Stratford, New Zealand

Millais, D.G.  
Millais Nurseries, Farnham, Surrey, England

Millais Nurseries  
Farnham, Surrey, England

Miller, L.O.  
Grove, Oklahoma, USA

Miller, W.C.  
Bethesda, Maryland, USA

Mitchell, R.A.  
Volcano, Hawaii, USA

\*Morrison, B.Y.  
Glenn Dale, Maryland, USA

Moyles, E.W.  
Oakland, California, USA

\*Moynier, W.  
Los Angeles, California, USA

Murata, H.  
Matsumoto-city, Nagano-prefecture, Japan

\*Neal, J.  
Frederick, Maryland, USA

Ohta, K.  
Shinshiro-city, Aichi-prefecture, Japan

Oldham, B.E.  
Auckland, New Zealand

Pacific Island Nursery  
Kea'au, Hawaii, USA

Parsons, J.  
Head gardener, Caerhays Castle, Cornwall, England

Petton, J.-F.  
Prat Vagagen, Scaër, France

Petton, L.  
Maisdon-sur-Sèvre, France

- Reich, W.  
Alfeld, Germany
- Reiley, H.E. & Mrs M.L.  
Woodsboro, Maryland, USA
- Roelink, G.  
Vireya Valley Nurseries, Cockatoo, Victoria, Australia
- \*Rothschild, E. de  
Exbury Gardens, Hampshire, England
- Rothschild, Leopold de  
Exbury Gardens, Hampshire, England
- Rouse, A.  
Hawthorn East, Victoria, Australia
- \*Rouse, J.L.  
Toorak, Victoria, Australia
- Saint-Jalm, J.F.  
Guémené-sur-Scorff, France
- Sande, J. van de  
Neptune, New Jersey, USA
- Sandwich Club  
Sandwich, Massachusetts, USA
- Saner, D. & M.  
North Bend, Washington, USA
- Sather, E.  
Brinnon, Washington, USA
- Schröder, T.  
Wiefelstede-Dringenburg, Germany
- \*Schrope, N.  
Hegin, Pennsylvania, USA
- \*Sifferman, K.I.  
Seattle, Washington, USA
- Singing Tree Gardens  
McKinlayville, California, USA
- Smart, D.  
Carnation, Washington, USA
- Smith, C.  
Orting, Washington, USA
- Smith, T.  
Mount Airy, Maryland, USA
- Snell, G.L.S.  
Highfields, Queensland, Australia
- Snyder, I.G.  
Stevenson, Washington, USA
- Société Bretonne du Rhododendron  
Scaër, France
- Southerland, S.B.  
Chapel Hill, North Carolina, USA
- Starling, B.N.  
Drewsteignton, Exeter, Devon, England
- Steele, R.M.  
Bayport Plant Farm, Lunenburg County, Nova Scotia, Canada
- Stent, Mrs  
England
- Suchy, Alan & Angela  
Bellingham, Washington, USA
- \*Sullivan, P.  
San Francisco, California, USA
- Thornton, J.O.  
Conyers, Georgia, USA
- Thornton, J.T.  
Franklinton, Louisiana, USA
- Trewithen Gardens  
Probus, Cornwall, England
- Viraraghavan, M.S.  
Kodaikanal, Tamil Nadu, India
- Wagstaff, S.  
Te Kuiti, New Zealand
- Wallace, D.S.  
Singing Tree Gardens, McKinlayville, California, USA
- Watson, Mrs E.M.  
Kirkland, Washington, USA
- Weekes, T.  
Oyster Bay Cove, New York, USA
- Westhoff, J.  
Bremen, Germany
- White, G.  
Kingham, Oxfordshire, England
- Williams, C.H.  
Caerhays Castle, Gorran, Cornwall, England
- \*Williams, J.C.  
Caerhays Castle, Gorran, Cornwall, England
- Winberg, J.D.  
Fall City, Washington, USA
- Winkler, W.F.  
Brunswick, Maine, USA
- Woodard, G.  
Westbury, New York, USA
- Wright, H.R.  
Courtenay, British Columbia, Canada
- Yoshimura, E.  
Okazaki-city, Aichi-prefecture, Japan

# Corrections to the International Rhododendron Register and Checklist

The following list indicates further significant additions and corrections that need to be made to the printed text of the *Register and Checklist* published in 2004 or to the five subsequent Supplements.

The Registrar is grateful to all those who have brought such errors to his notice and welcomes knowledge of further amendments that may need to be made.

Minor changes are not noted here but such changes have been made to the database and will be available in any future electronic version.

The database will also be taking into account additional information that comes to light relating to any individual entry, as well as including further entries to the Checklist.

In the following list the page number refers to the page in the *Register and Checklist* (or one of its Supplements) on which the entry concerned is to be found.

**‘Ada Brunières’** (p.19)

Following information provided from Donald Waterer this cultivar should be shown as a Koster-raised Occidentale hybrid

**‘Caparo’** (Fifth Supplement, p.6)

The pollen parent should be amended to ‘Stagecoach Frills’ × ‘Twilight Sky’, a selection that has been named ‘Arouca’

**‘Cherish’** (p.237)

This should be listed as an evergreen azalea not a rhododendron

**‘Glen Catacol’** (p.514)

The measurement of leaf dimensions should be in cm not mm

**‘Haida Gold’** (p.568)

This is now known to be a trade designation for the plant previously registered as ‘Bob’s Yellow’

**‘Ibykus’** (Fifth Supplement, p.14)

The meaning of the word kranich in German is crane not stork

**‘Koromo-shikibu’** (p.736)

For *macropetalum* read *stenopetalum* (*macrosepalum*)

**‘Madame de Cannaert d’Hamale’** (p.832)

This should be spelt ‘Madame de Cannart d’Hamale’

**‘Marabella’** (Fifth Supplement, p.19)

The pollen parent should be amended to ‘Stagecoach Frills’ × ‘Twilight Sky’, a selection that has been named ‘Arouca’

**‘Mayaro’** (Fifth Supplement, p.21)

The pollen parent should be amended to ‘Stagecoach Frills’ × ‘Twilight Sky’, a selection that has been named ‘Arouca’

*Rhododendron lochia*. Following the comments in Argent (*Rhododendrons of subgenus Vireya*, 2006) and after consultation with regional representatives for the ICRA, the Registrar has amended all entries in *The International Register and Checklist* which included mention of *R. lochia* to read *R. viriosum*, in the belief that no authentic hybrids have yet been named involving the true *R. lochia*. If anyone has evidence in individual cases that true *R. lochia* has been used in breeding please contact the Registrar.



ISBN 978-1-907057-23-6



9 781907 057236 >