

---

**Division 1****TRUMPET DAFFODIL CULTIVARS**


One flower to a stem; corona ('trumpet') as long as, or longer than the perianth segments ('petals')



---

**Division 2****LARGE-CUPPED DAFFODIL CULTIVARS**


One flower to a stem; corona ('cup') more than one-third, but less than equal to the length of the perianth segments ('petals')




---

**Division 3****SMALL-CUPPED DAFFODIL CULTIVARS**

One flower to a stem; corona ('cup') not more than one-third the length of the perianth segments ('petals')





**Division 4**

**DOUBLE DAFFODIL CULTIVARS**


One or more flowers to a stem, with doubling of the perianth segments or the corona or both

---

**Division 5**

**TRIANDRUS DAFFODIL CULTIVARS**

Characteristics of *N. triandrus* clearly evident: usually two or more pendent flowers to a stem; perianth segments reflexed





---

**Division 6**

**CYCLAMINEUS DAFFODIL CULTIVARS**

Characteristics of *N. cyclamineus* clearly evident: one flower to a stem; perianth segments significantly reflexed; flower at an acute angle to the stem, with a very sort pedicel ('neck')





**Division 7**

**JONQUILLA AND APODANTHUS  
DAFFODIL CULTIVARS**


Characteristic of Sections Jonquilla or Apodanthi clearly evident: one to five (rarely eight) flowers to a stem; perianth segments spreading or reflexed; corona cup-shaped, funnel-shaped or flared, usually wider than long; flower usually fragrant

---

**Division 8**

**TAZETTA DAFFODIL CULTIVARS**

Characteristics of Section Tazettae clearly evident: usually three to twenty flowers to a stout stem; perianth segments spreading not reflexed; flowers usually fragrant




---

**Division 9**

**POETICUS DAFFODIL  
CULTIVARS**

Characteristics of the *N. poeticus* and related species clearly evident; perianth segments pure white; corona very short or disc-shaped, not more than one-fifth the length of the perianth segments; corona usually with a green and/or yellow centre and red rim, but sometimes wholly or partly of other colours; anthers usually set at two distinct levels; flowers fragrant.






---

**Division 10****BULBOCODIUM DAFFODIL CULTIVARS**

## Characteristics of Section

Bulbocodium clearly evident: usually one flower to a stem; perianth segments insignificant compared with the dominant corona; anthers dorsifixed (ie attached more or less centrally to the filament); filament and style usually curved



---

**Division 11****SPLIT – CORONA DAFFODIL CULTIVARS**

Corona split – usually for more than half its length:

## a) Collar Daffodils

Split-corona daffodils with the corona segments opposite the perianth segments; the corona segments usually in two whorls of three

## b) Papilion Daffodils

Spilt-corona daffodils with the corona segments alternate to the perianth segments; the corona segments usually in a single whorl of six

Spilt-corona daffodils with the corona segments alternate to the perianth segments; the corona segments usually in a single whorl of six

---

**Division 12****OTHER DAFFODIL CULTIVARS**

Daffodil cultivars which do not fit the definition of any other division

---

**Division 13****DAFFODILS DISTINGUISHED SOLELY BY BOTANICAL NAME**

---