



THE ROYAL HORTICULTURAL SOCIETY

The International  
**RHODODENDRON**  
**REGISTER and**  
**CHECKLIST (2004)**  
*Fourth Supplement*



© 2008 The Royal Horticultural Society  
80 Vincent Square, London SW1P 2PE, United Kingdom  
[www.rhs.org.uk](http://www.rhs.org.uk)

International Rhododendron Registrar:  
Dr A C Leslie

All rights reserved. No part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright holder

ISBN 9781902896915

Printed and bound in the UK by the University Press, Cambridge

Cover: *Rhododendron* 'Mrs Charles E. Pearson' by Alfred John Wise

# THE INTERNATIONAL RHODODENDRON REGISTER AND CHECKLIST (2004) Fourth Supplement

The second edition of the *International Rhododendron Register and Checklist* included accounts of all plants whose names had been registered up to the end of December 2002. The first three Supplements included names registered from 1 January 2003 to 31 December 2006. This fourth Supplement adds those names registered between 1 January 2007 and 31 December 2007. A list of further corrections to the 2004 *Register and Checklist* is also included.

The layout of each entry follows the format adopted in the 2004 *Register and Checklist* and is explained there in detail on pages 7-10 of the Introduction. It may be helpful to repeat here that the following abbreviations are employed:

- (a) = azalea
- (az) = azaleodendron
- cv. = cultivar
- G = grown to first flower by...
- gp = Group
- H = hybridised by...
- I = introduced by...
- N = named by...
- (r) = rhododendron
- R = raised by...
- REG = registered by...
- S = selected by...
- (s) = seed parent
- (v) = vireya

Where possible colour references are given to the *RHS Colour Chart* (1966, 1986, 1995 & 2001) with colour descriptions being taken from *A contribution towards standardization of color names in horticulture*, by R.D. Huse and K.L. Kelly (ed. D.H. Voss), 1984.

All raisers and introducers of new rhododendrons are urged to ensure that the names of their plants have been registered and are reminded that registration needs to be effected before the plant is introduced or the name appears in print. Registration forms can be obtained from the International Rhododendron Registrar (at the address given below), from the RHS website ([www.rhs.org.uk/plants/registerpages/rhodoform.pdf](http://www.rhs.org.uk/plants/registerpages/rhodoform.pdf)) or from national or regional registrars as follows:

- |               |  |
|---------------|--|
| North America | Mrs J.W. Murray, 21 Squire Terrace, Colts Neck, New Jersey<br>07722-1021, USA    |
| Australia     | Mr K. Gillanders, Woodbank, 2040 Huon Road, Longley,<br>Tasmania 7150, Australia |
| New Zealand   | Mr B.J. Coker, 42 Massey Crescent, Christchurch 1, New Zealand                   |
| Japan         | Mr K. Fujiwara, 363-3 Yoshioka-cho, Kanonji-shi, Kagawa-ken,<br>Japan 768-0021   |

Dr A.C. Leslie, International Rhododendron Registrar, 109 York Street, Cambridge  
CB1 2PZ, England (Email: [alanleslie@rhs.org.uk](mailto:alanleslie@rhs.org.uk))

# INTERNATIONAL RHODODENDRON REGISTER AND CHECKLIST

## (r) 'Alice Boyd'

cv. Elepidote rhododendron : Olympic Lady Group (s) × *campylocarpum*. H: A. Craig-Mooney (1983), G: A. Craig-Mooney (1990), N: A. Craig-Mooney (2006), REG: A. Craig-Mooney (2007). Fls 5/lax truss, funnel-campanulate, 50 × 60mm, with 5 wavy-edged lobes, buds strong red to light pink (50 A–D), opening light purplish pink (73C) both inside and out, fading to white; two vivid purplish red (61C) blotches in the throat. Calyx green. Lvs orbicular, 60–70 × 50–60mm, with slightly wavy edges, matt green above, cordate at base, obtuse at apex, hairless. Shrub 3.0 × 2.0m in 20 years. April to May. (Nomenclatural standard: colour print provided by the registrant (WSY0100756). Etymology: named for Viscountess Boyd of Ince Castle, Cornwall, England).

## (r) 'Always April'

cv. Elepidote rhododendron : 'Mindy's Love' (s) × 'Pink Petticoats'. H: J. Barlup (1995), G: J. Barlup (2001), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 24/ball truss, saucer-shaped, c.50 × 75mm, with 5 very wavy-edged lobes, strong purplish red (61B) in bud, opening inside very pale purple (69C) with slightly deeper very pale purple (69A) edging, the dorsal lobe tinged pale yellow (11D) from base extending upwards for c.40mm; outside very pale purple (69C) flushed with darker moderate purplish pink (65A) on the dorsal lobes, and with strong reddish purple (70B) longitudinal veins on all lobes; base of the dorsal lobe with two strong red (53D) flares, solid at base and extending upwards for c.10mm and then dense, tiny spots of the same colour extending for another 20mm. Lvs elliptic, 115 × 45mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.5m in 10 years. Late April.

## (a) 'Amherst'

cv. Evergreen azalea : (un-named hybrid (CB1, fls semidouble) × 'Desiree' (Shammarello)) (s) × un-named hybrid (CB1, fls semidouble). H: Robert Stewart (1992), G: Robert Stewart (1995), N: Robert Stewart (2006), REG: Robert Stewart (2007). Fls single, 3/truss, broadly funnel-shaped, 45 × 75mm, with 6 wavy-edged lobes, white in bud, opening white inside and out, with

sparse light yellow-green (142C) spotting in the throat. Calyx 15mm long, slightly more yellow than strong yellow-green (142A). Lvs elliptic, 55 × 20mm, broadly acute at apex, cuneate at base, margins upcurved, semi-glossy and moderate olive green to moderate yellow-green (137B–C) above, hairless. Shrub 0.8 × 1.1m in 10 years. Midseason.

## (r) 'Anne-Marie Sim'

cv. Lepidote rhododendron : 'Kotuku' (the *maddenii* selection) (s) × *maddenii* Polyandrum Group. H: T.A. Garbutt (1987), G: Mrs A. Sim (2003), N: T.A. Garbutt (2007), REG: T.A. Garbutt (2007). Fls 7/lax truss, tubular funnel-shaped, 110 × 140mm, with 5 wavy-edged lobes, buds white, flushed pink, opening inside white, with a yellow throat, outside white, flushed pink; not marked; with a musky scent. Calyx 2mm long, green. Lvs lanceolate, 140 × 45mm, margins flat, cuneate at base, acute at apex, glossy green above, scaly. Shrub with an open habit, 1.5 × 1.0m in 5 years. Late November (New Zealand). (Nomenclatural standard: colour print provided by the registrant (WSY0100130). Etymology: named for a friend of the registrant who is growing the plant. Sometimes listed in error as 'Anne Marie Sim').

## (r) 'April Sunset'

cv. Elepidote rhododendron : ('Apricot Nectar' × 'Lem's Cameo') (s) × 'Lem's Cameo'. H: C. Weeks (1998), G: C. Weeks (2002), N: C. Weeks, I: Westgate Gardens Nursery (2007), REG: C. Weeks (2007). Fls 12–13/domed truss, funnel-campanulate, 65 × 75mm, with 6 wavy-edged lobes, vivid red (45A) in bud, opening inside strong red (39A) at margins, blending with brilliant yellow (8A) towards centre and then to strong red (41B) and finally moderate red (47A) at base; outside moderate red (47A) to deep yellowish pink (47C); unmarked. Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and dark yellowish green (139A) above, hairless. Shrub 1.2 × 1.2m in 9 years. Early to mid April.

## (r) 'Atrotrill'

cv. Elepidote rhododendron : 'Atroflo' (s) × 'Trilby'. H: D.B. Johnson (1986), G: D.B. Johnson (1995), N: D.B. Johnson (2006),

REG: D.B Johnson (2007). Fls 13/domed truss, funnel-shaped, 45 × 65mm, with 5 very wavy-edged lobes, dark red (59A) in bud, opening inside strong purplish red (61B), fading lighter to vivid purplish red (57D) at centre; outside similarly coloured; black spotting on the dorsal lobe. Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (137A) above, hairy all over when young, but only beneath when mature (hairs whitish at first, brownish when mature). Shrub with an open habit, 2.1 × 1.2m in 20 years. Early May.

(r) **'Betty Spady'**

cv. Elepidote rhododendron : *yakushmanum* 'Koichiro Wada' (s) × *proteoides* (Rock 147). H: C.C. Smith, G: C.C. Smith, N: H.A. Spady (2007), I: Cecil & Molly Smith Garden, REG: H.A. Spady (2007). Fls 11/ball truss, open funnel-shaped, 30 × 50mm, with 6 wavy-edged lobes, pale purplish pink (56A) in bud, opening inside nearly white, tinged with pale purplish pink (56A), the dorsal lobes heavily spotted with deep purplish pink (55A); outside opening pale purplish pink (56D), each lobe striped with pale purplish pink (56A). Calyx minute. Lvs oblong, 40–50 × 15–20 mm, broadly acute at apex, cuneate to rounded at base, with downcurved margins, dull and moderate olive green to moderate yellow-green (147A–B(–C)) above, with a floccose indumentum above at first, later hairless, the lower surface felted and persistently hairy (the hairs moderate orange-yellow (168D) at first aging to moderate reddish brown (174B)). Dense shrub, 0.9 × 0.9m in 20 years. Early.

(r) **'Billy Wilder'**

cv. Elepidote rhododendron : 'Goldsworth Orange' (s) × (('Inca Gold' × *yakushmanum*) × ('Catalgl' × *dichroanthum*)). H: D. MacMullan (1983), G: D. MacMullan (1987), N: D. MacMullan (2006), REG: D. MacMullan (2007). Fls 14/lax truss, funnel-shaped, 65 × 65mm, with 5 flat to wavy edged, reflexed lobes, vivid reddish orange (33B) in bud, opening inside light yellowish pink (29C) in the throat, shading to light yellowish pink (19B) at the margins, with a deep yellowish pink (39B) blotch at the centre of the throat; outside deep yellowish pink (39B) near the base, lobes strong orange-yellow (17A), with stripes of strong orange (25A) extending from the base to the margins. Pedicels vivid reddish orange (33B). Lvs elliptic, 125 × 40mm, broadly acute at apex, cuneate at base, margins flat, dull fir green above, hairless. Shrub 0.9 × 1.2m in 10 years. Midseason.

(r) **'Bodacious'**

cv. Elepidote rhododendron : ('Ring of Fire' × 'Lem's Cameo') (s) × 'One Thousand Butterflies'. H: J.S. Smith (1998), G: J.S. Smith (2002), N: J.S. Smith (2002), I: Roadside Rhododendrons (2005), REG: J.S. Smith (2007). Fls 32/domed truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, inside with a continuous moderate pink (51D) edge, changing abruptly to a pale greenish yellow (2D) throat; outside strong red (46A). Calyx 25mm long, strong greenish yellow (151B). Lvs lanceolate, 245 × 65mm, broadly acute at apex, oblique at base, glossy and moderate olive green (147A) above, hairless. Dense shrub, 1.8 × 1.2m in 10 years. Mid to late April.

(r) **'Brimble'**

cv. Elepidote rhododendron : *griffithianum* (s) × 'Hotei'. H: A. Craig-Mooney (1984), G: A. Craig-Mooney (1991), N: A. Craig-Mooney, REG: A. Craig-Mooney (2007). Fls 9/full, rounded truss, broadly funnel-shaped, 80mm wide, with 5 wavy-edged lobes, buds pale yellow (11C), opening pale greenish yellow (2D) both inside and out, fading to white, the throat pale yellow-green (149C); unmarked; filaments white, anthers brown; pistil yellow. Calyx unevenly lobed, coloured as the corolla. Lvs elliptic, 100 × 45mm, broadly acute at apex, broadly cuneate at base, with wavy margins, hairless. Shrub 2.25 × 1.4m in 15 years. April to May. (Nomenclatural standard: colour print provided by the registrant (WSY0100745). Etymology: named after a pet labrador dog which died in 2003 and belonged to the registrant).

(r) **'Butter Rum'**

cv. Elepidote rhododendron : 'Janet Blair' (s) × ('Nancy Evans' × (('Whopper' × 'Brandt's Tropicana') No.2)). H: J.S. Smith (1999), G: J.S. Smith (2004), N: J.S. Smith (2007), I: Roadside Rhododendrons (2007), REG: J.S. Smith (2007). Fls 12/flat truss, broadly funnel-shaped, 65 × 90mm, with 7 very wavy-edged lobes, opening inside pale greenish yellow (13D), with deep yellowish pink (47C) spotting on the two dorsal lobes; the spots show through the lobes creating an orange appearance; outside pale yellow (18C). Calyx c.15mm long, deep purplish pink (55A). Lvs elliptic, 175 × 50mm, broadly acute at apex, oblique at base, with wavy margins, semi-glossy and moderate olive green (137A) above, hairless. Dense shrub, 1.2 × 0.9m in 5 years. Mid May.

(r) **'Cascade Dawn'**

cv. Elepidote rhododendron : probably 'Hotei' or 'Odee Wright', open-pollinated. H: D.E. King (c. 1995), G: D.E. King (c. 2001),

N: D.E. King (2007), REG: D.E. King (2007). Fls 10/domed truss, broadly funnel-shaped, 55 × 75mm, with 6 wavy-edged lobes, inside of the corolla progresses with age from pink through yellow and cream to white, with a dark orange-red to red blotch in the throat; outside colour progresses with age from pink to cream; red bud scales appear on the new growth during the flowering season, adding significantly to the attractiveness of the plant. Calyx 2–3mm long, red. Lvs elliptic, 115 × 40mm, cuneate to rounded at base, acute at apex, margins wavy, dull and mid green above, hairless. Shrub 1.1 × 1.1m in 15 years. Mid April to early May.

(r) **'Catherine's Coral Delight'**

cv. Elepidote rhododendron : 'Raspberry Swirl' (s) × 'Lem's Cameo'. H: C. Weeks (1995), G: C. Weeks (2002), N: C. Weeks (2007), I: Westgate Gardens Nursery (2007), REG: C. Weeks (2007). Fls 8/domed truss, funnel-shaped, 50 × 65mm, with 5 wavy-edged lobes, strong reddish orange (31A) in bud, opening inside strong purplish pink (63C), slightly darker strong purplish red (63B) at edges and tinged yellow in the centre; outside strong purplish pink (62A) to moderate purplish pink (62B); unmarked. Calyx 20mm long, coloured as corolla. Lvs elliptic, 175–200 × 60mm, broadly acute at apex, rounded at base, margins slightly downcurved, semi-glossy and dark yellowish green (139A) above, hairless. Dense shrub, 0.6 × 1.0m. Mid to late March.

(r) **'Catherine's Cutie Pie'**

cv. Elepidote rhododendron : 'Sierra Sunrise' (s) × 'Lori Eichelser'. H: C. Weeks (1994), G: C. Weeks (1998), N: C. Weeks, I: Westgate Gardens Nursery (2007), REG: C. Weeks (2007). Fls 7/flat truss, campanulate, 40 × 40mm, with 5 wavy-edged lobes, inside moderate pink (49B), tinged light yellowish pink (159A); outside deep yellowish pink (39B); unmarked. Calyx 35mm long, coloured as the corolla. Lvs elliptic, 35 × 15mm, broadly acute at apex, cuneate at base, margins flat and downcurved, dull and moderate yellow-green (148A) above, hairless. Dense shrub, 0.2 × 0.4m in 13 years (growing only 10–20mm/year). Mid to late March.

(a) **'Chariots of Fire'**

cv. Evergreen azalea : parentage unknown. H: W.G.T. Hyde (late 1960s), G: W.G.T. Hyde (late 1970s), N: Mrs R.G. Legrand (2007), I: Mrs R.G. Legrand (2007), REG: Mrs R.G. Legrand (2007). Fls 9–19/ball truss, open trumpet-shaped, 60 × 30mm, with 5 wavy-edged lobes, vivid red (45A) in bud, opening vivid reddish orange (42A) throughout; scented. Calyx 12mm long. Lvs narrowly

elliptic, 120 × 45mm, margins decurved, cuneate at base, acute at apex, glossy green above, hairless. Shrub 1.55 × 1.5m in 30 years. May.

(r) **'Charlotte Weingarten'**

cv. Elepidote rhododendron : parentage unknown. H: C.B. Smith (1987), G: B. Weingarten (1993), N: B. Weingarten, I: Van Veen Nursery, REG: B. Weingarten (2007). Fls c.20/conical truss, open to broadly funnel-shaped, 75 × 50mm, with 5 very wavy-edged lobes, dark purple (79A) in bud, opening inside deep purplish red (71A), with a dark greyish purple (202A) blotch; outside dark purple (79A); anthers whitish. Lvs elliptic, 115 × 45mm, acute at apex, cuneate at base, margins flat, glossy and moderate olive green above, hairless. Shrub 1.5 × 1.2m in 20 years. Mid May.

(a) **'Christopher'**

cv. Evergreen azalea : parentage unknown. H: W.G.T. Hyde (1960s), G: W.G.T. Hyde, N: Mrs R.G. Legrand (2007), I: Mrs R.G. Legrand (2007), REG: Mrs R.G. Legrand (2007). Fls star-like in form, 2/truss, 39 × 15mm, with 5 wavy-edged lobes, vivid red (45C) in bud, opening inside vivid red (45C), with a hint of strong red (46A) on the three dorsal lobes, outside vivid red (45C). Calyx 5mm long, strong yellow-green (143C). Lvs narrowly elliptic, 50 × 20mm, margins decurved, cuneate at base, acute at apex, glossy green, hairless. Shrub 0.7 × 1.0m in 20 years. May. (Etymology: named for the husband of the registrant).

(r) **'Chromatic Fantasy'**

cv. Elepidote rhododendron : 'Reigate Squire' (s) × 'Peach Cobbler'. H: R.H. Gustafson (1992), G: R.H. Gustafson, N: R.H. Gustafson (2006), REG: R.H. Gustafson (2007). Fls 10/domed truss, broadly funnel-shaped, c.85 × 115mm, with 6 wavy-edged lobes, buds strong purplish pink (62A), with pale purplish pink ribs, opening inside light purplish pink (62C) at edges, shading to pale purplish pink (62D), with a moderate red (180A) throat and moderate red (180A) flare fading to light orange-yellow (16C) on the two dorsal lobes; outside light purplish pink (62C) at edges, shading to pale purplish pink (62D), with pale yellow (19C) on ribs. Calyx rim-like, light yellow-green (145B). Lvs elliptic, 125 × 65mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.5 × 1.2m in 6 years. Late May.

(r) **'Cinnarose'**

cv. Elepidote rhododendron : 'Anita Dunstan' (s) × 'Hill's Low Red'. H: J. Barlup (1997), G: J. Barlup (2001), N: J. Barlup (2006),

REG: J. Barlup (2007). Fls 15/ball truss, broadly funnel-shaped, 40 × 50mm, with 5 wavy-edged lobes, strong red (50A) in bud, opening inside pale yellowish pink (27C), with streaks of strong red (53D) on midveins and slight touches of the same colour on all the lobes; outside pale yellowish pink (27C) infused with strong red (53D), particularly on the dorsal lobes, extending to the tips, and on all the midveins; deep red (53A) basal blotch extending c.10mm from the centre. Calyx 20mm long, pale yellowish pink (27C), with strong red (53D) lobe tips and three spots, c.5mm long, of deep red (53A). Lvs elliptic, 100 × c.50mm, broadly acute at apex, rounded at base, margins downcurved, dull and moderate olive green (147A) above, hairless. Shrub 0.7 × 1.0m in 8 years. Early May.

(f) **'Claret Mist'**

cv. Elepidote rhododendron : 'Jonathan Shaw' (s) × 'Plum High'. H: J. Barlup (2000), G: J. Barlup (2004), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 17/ball truss, saucer-shaped, c.50 × 85mm, with 5 very wavy-edged lobes, inside with touches of pale yellow (11D) from the base, blending with very pale purple (76D) midribs and deepening at margins to very light purple (75C), with darker edges of light purple (75B) on the dorsal lobe; outside very pale purple (76D), with darker midveins of very light purple (75C) and the tip of the dorsal lobe light purple (75B); dark red (183A) speckled dorsal flare extending upwards c.35mm from the base. Lvs elliptic, 115 × c.60mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × c.1.0m in 5 years. Late May.

(a) **'Corley's Cardinal'**

cv. Deciduous azalea : *calendulaceum*, yellow-flowered (s) × *calendulaceum*, orange-flowered. H: R.O. Smitherman (1987), G: T.E. Corley (1992), N: R.O. Smitherman (2000), REG: R.O. Smitherman (2007). Fls single, 7–8/bud with up to three buds in a domed truss, tubular funnel-shaped, 45 × 50mm, with 5 wavy-edged lobes, vivid reddish orange (34B) in bud, opening vivid reddish orange (34B) throughout, with a vivid yellowish pink (30C) dorsal blotch. Lvs obovate, 50–70 × 30–40mm, broadly acute at apex, cuneate at base, dull and moderate olive green (137B) above, hairless. Shrub 2.4 × 2.4m in 22 years. Mid April.

(r) **'Creamcicle'**

cv. Elepidote rhododendron : 'Brandt's Tropicana' (s) × 'Apricot Fantasy'. H: W.A. Stipe (1989), G: W.A. Stipe (1993), N: S. Stipe (2007), I: Glynneden Gardens (2007),

REG: W.A. Stipe (2007). Fls 14/ball truss, broadly funnel-shaped, 55 × 65mm, with 5 wavy-edged lobes, strong red (46A) in bud, opening inside light orange-yellow (22C), with a strong red (53B) dorsal flare; outside brilliant orange (25C). Calyx 50mm long, light orange (22C), with strong red (53B) dorsal flare. Lvs oblong, 115 × 50mm, obtuse at apex, oblique at base, margins downcurved, semi-glossy and dark green (135A) above, hairless. Shrub 2.4 × 1.8m in 15 years. Early May.

(r) **'Cymbeline'**

cv. Elepidote rhododendron : parentage unknown. H: W.G.T. Hyde (1960s), G: W.G.T. Hyde (1970s), N: Mrs R.G. Legrand (2007), I: Mrs R.G. Legrand (2007), REG: Mrs R.G. Legrand (2007). Fls 16/compact ball truss, open trumpet-shaped, 50 × 30mm, with 5 very wavy-edged lobes, buds vivid purplish red (61C), opening shaded pale purplish pink (65C), with the edges tinted deep purplish pink (66C) and with delicate vivid purplish red (66B) speckles on the dorsal lobe, outside moderate purplish pink (66D). Calyx 2mm long, deep purplish pink (66C). Lvs narrowly elliptic, 150 × 55mm, margins decurved, attenuate at base, acute at apex, matt green above, with a hairy indumentum – near brownish orange (165B) but darker and more ginger in colour. Shrub 1.2 × 1.2m in 15–20 years. May. (Etymology: named as part of a collection with Shakespearean names).

(a) **'Debra Lee'**

cv. Evergreen azalea : ('Saybrook Glory' × 'Dorothy Clark') (s) × 'Saybrook Glory'. H: Robert Stewart (1996), G: Robert Stewart, N: Robert Stewart (2006), REG: Robert Stewart (2007). Fls single, 3/truss, broadly funnel-shaped, 45 × 75mm, with 5 wavy-edged lobes, slightly deeper than deep purplish pink (67C) in bud, opening inside bicoloured: with a band of light purple (75B) to very light purple (75C) at margins, changing abruptly to a paler shade of the same colour, with purplish pink dorsal spotting; outside close to deep purplish pink (67C). Calyx c.6mm long, close to brilliant yellow-green (142B). Lvs elliptic, 55 × c.25mm, broadly acute at apex, cuneate at base, margins upcurved, semi-glossy and close to moderate yellow-green (139C) above, hairless. Shrub 0.5 × 0.6m in 10 years. Midseason.

(r) **'Dick Brooks'**

cv. Elepidote rhododendron : 'Dexter's Giant Red' (s) × *yakushimanum*, Whitney dwarf selection. H: C. Patterson (1989), G: G. Hibben (1995), N: G. Hibben (2006), I: RareFind Nursery, REG: G. Hibben (2007). Fls 7–12/domed truss, broadly

funnel-shaped, c.55 × 100mm, with 5–7 wavy-edged lobes, deep to strong purplish pink (55A–B) in bud, opening inside with moderate purplish pink (62B) margins and flushed lighter pink, aging to white; with a basal blotch of deep red (53A) overlaid with dark red (187A), and a spotted dorsal flare of the same colour; outside also aging to white, after opening with strong purplish pink (62A) margins and midveins. Calyx minute, light yellow-green (145C). Lvs elliptic, 100–150 × 30–45mm, broadly acute at apex, slightly oblique at base, margins flat, dull and moderate olive green (137A) above, new leaves with hairs above and below, mature leaves with hairs only below and heaviest on the midribs (hairs white (155D) when young, aging to moderate orange-yellow (164B)). Dense shrub, 1.5 × 3.0m in 17 years. Mid to late May. (Published by S. & J. Perkins in *J.Am.Rh.Soc.* 61(2):103 (2007), with photograph).

(v) **‘Dick Shaw’**

cv. *Lepidote rhododendron* : selection from *loranthiflorum* subsp. *loranthiflorum*, collected on Papua New Guinea by L.A. Craven (RBGE accession no. 19830533. S: Royal Botanic Gardens, Edinburgh, N: G. Argent (2007), REG: Royal Botanic Garden, Edinburgh (2007). Fls 4–7/loose truss (fls semi-erect to half hanging), trumpet-shaped, 35–42 × 25–35mm, with 5 flat-edged lobes, opening white (155D) throughout, unmarked; faintly sweet-scented. Calyx c.1mm long, green. Lvs obovate, 25–60 × 15–25mm, margins decurved, tapering at base, obtuse to rounded at apex, with a scaly indumentum (scales white at first, maturing to pale brown). Shrub 1.5 × 1.0m in 10 years. Flowering at irregular intervals. AM 2007. (Nomenclatural standard: herbarium specimen of the award plant (WSY0100901). Etymology: named for a former Curator of the Royal Botanic Gardens, 1972–1987).

(a) **‘Dorset Sandy’**

cv. Evergreen azalea : parentage unknown. H: W.G.T. Hyde (late 1960s), G: W.G.T. Hyde (late 1970s), N: Mrs R.G. Legrand (2007), I: Mrs R.G. Legrand (2007), REG: Mrs R.G. Legrand (2007). Fls open trumpet-shaped, 1–2/truss, 40 × 23mm, with 5 wavy-edged lobes, strong red (39A) in bud, opening deep yellowish pink (39B) throughout, the dorsal lobe with muted strong red (53B) speckles; retains colour well. Calyx 5mm long, strong yellow-green (144C). Lvs elliptic, 25 × 13mm, margins flat, cuneate at base, obtuse at apex, glossy green above, hairless. Shrub 0.67 × 0.9m in 20 years. Late May to early June. (Etymology: named for the strong family connections with the county of Dorset).

(r) **‘Entourage’**

cv. *Elepidote rhododendron* : (‘Top Banana’ × *brachycarpum* (as subsp. *tigerstedtii*)) (s) × ‘Delicate Air’. H: A.T. & S.A. Anderson (1997), G: A.T. & S.A. Anderson (2003), N: A.T. & S.A. Anderson (2006), REG: A.T. & S.A. Anderson (2007). Fls 11/ball truss, broadly funnel-shaped, 50 × 80mm, with 5 wavy-edged lobes, light yellow (18A) in bud, opening light yellow-green (2C) throughout, with brilliant greenish yellow (6C) speckling on the dorsal lobe and along the adjacent sides of the flanking lobes. Lvs lanceolate, 140 × 55mm, acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Shrub 0.75 × 0.7m in 8 years. Early May.

(r) **‘Fifi’**

cv. *Elepidote rhododendron* : *yakushimanum* ‘Koichiro Wada’ (s) × *arboreum* ‘Tony Schilling’. H: F.S. Norman (1980), G: F.S. Norman (1990), N: F.S. Norman (2007), REG: F.S. Norman (2007). Fls c.20/finally globose truss (slightly open or loose at the top), tubular-campanulate, c.40 × c.45mm, with 5 wavy-edged lobes, bright deep pink in bud, opening inside blush pink, outside pale blush pink with deeper stripes, some relatively inconspicuous, small (c.0.5mm), deep pink spots on the lower two thirds of the dorsal lobe; flowers do not bleach to white until the corolla drops. Calyx 2.5mm long, pale greenish yellow. Lvs narrowly elliptic, 135 × 42mm, margins flat but the leaf transverse section V-shaped, matt green above, narrowly obtuse at apex, narrowly obtuse to cuneate at base, with a white indumentum when young which ages to buff. Shrub c.2.2 × 1.6m in 30 years. Mid April. (Nomenclatural standard: colour print provided by the registrant (WSY0096964). Etymology: named after the registrant’s wife).

(a) **‘Fruit Salad’**

cv. Deciduous azalea : parentage unknown. H: M.C. Pratt, S: E.G. Millais (c. 2000), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 8/truss, tubular funnel-shaped, 30 × 45–55mm, with 5 wavy-edged lobes, buds strong red (39A), opening moderate pink (49B) to strong purplish red/deep purplish pink (54A–B), shading to light purplish pink (55C); outside light purplish pink (55C), shading slightly darker at margins; dorsal lobe almost entirely vivid orange-yellow (23A), shading to brilliant orange-yellow (23B) at the margins; stamens exerted, 80mm long, filaments pale pink, anthers pale cream; style pale pink; slightly scented. Calyx 3mm long, light green. Lvs elliptic, 70 × 25mm, margins wavy, glossy



above, acute at apex. Shrub 1.5 × 1.0m in 10 years. Late June. (Selected by E.G. Millais as a plant labelled DP103 at Dunham Massey (National Trust, UK). Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100240)).

(r) '**Ghost Bomber**'

cv. *Elepidote rhododendron* : 'Ghost' (s) × 'Red Bomber'. H: A.T. & S.A. Anderson (1998), G: A.T. & S.A. Anderson (2003), N: A.T. & S.A. Anderson (2006), REG: A.T. & S.A. Anderson (2007). Fls 17–18/domed truss, saucer-shaped, 40 × 75mm, with 5 wavy-edged lobes, opening inside deep purplish pink (55A) at margins, shading to white (155D) in the centre, with faint, light yellow-green (154D) dorsal spotting; outside deep purplish pink (55A) at edges, shading to strong, light and pale purplish pink (55B–D). Lvs elliptic, 165 × 70mm, broadly acute at apex, cordate at base, with wavy margins, dull and moderate olive green (137A) above, with a light, flat coating of hairs below. Shrub 0.6 × 0.8m in 8 years. Late May.

(r) '**Ginger Peach**'

cv. *Elepidote rhododendron* : 'Nancy Evans' (s) × (((Jalisco Group × *yakushmanum*) × *yakushmanum*) × 'Joanita') × ('Goldfort' × 'Odee Wright')). H: J. Barlup (2000), G: J. Barlup (2005), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 13/ball truss, broadly funnel-shaped, c.50 × 75mm, with 5 wavy-edged lobes, deep red (53A) in bud, opening inside pale yellow (12D), with dorsal lobes brilliant yellow (12B) and tinges of light yellowish pink (38C) on lobe edges; outside similar but with 38C markings more pronounced on the dorsal lobes; deep red (53A) nectar pouches at the base. Calyx c.30mm long, brilliant yellow (12B), with light yellowish pink (38C) tips. Lvs elliptic, 135 × c.50mm, broadly acute at apex, rounded at base, margins wavy, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.8 × 1.2m in 5 years. Early May.

(r) '**Gle003**'

cv. *Elepidote rhododendron* : 'Whitney's Best Yellow' (s) × 'Lem's Cameo'. H: J. Somers (1983), G: P.A. & K.N.E. Cox (1988), N: P.A. & K.N.E. Cox, I: Glendoick Gardens, REG: P.A. & K.N.E. Cox (2007). Fls 13–15/rounded truss, funnel-campanulate, 60mm wide, with 7 wavy-edged lobes, buds light orange/light yellowish pink (29B–C), opening inside brilliant yellow/light greenish yellow (8A–B), shading to light yellowish pink (29C), outside similarly coloured but more mixed; basal blotch of vivid reddish orange (44B); slightly scented. Calyx 10–13mm

long, brilliant yellow/light greenish yellow (8A–B) shading to light yellowish pink (29C). Lvs ovate/obovate, 80–100 × 40–50mm, margins flat, glossy above, hairless. Shrub 2.0 × 2.0m in 20 years. May. Trade designation: GLENDOICK BUTTERSCOTCH. (Glendoick is a registered trademark belonging to Glendoick Gardens).

(r) '**Gle011**'

cv. *Elepidote rhododendron* : 'Goldkrone' (s) × 'Cupcake'. H: P.A. & K.N.E. Cox, G: P.A. & K.N.E. Cox, N: P.A. & K.N.E. Cox, I: Glendoick Gardens (2005), REG: P.A. & K.N.E. Cox (2007). Fls 13/open-topped truss, funnel-campanulate, 40 × 35mm, with 5 flat-edged lobes, buds light greenish yellow (4B), opening light greenish yellow (4B) throughout, with vivid red (45A) spots on the dorsal lobe. Calyx 2mm long, green. Lvs obovate 50–60 × 25–32mm, margins a little decurved, cuneate at base and apex, hairless. Shrub 1.0 × 1.0m in 10–12 years. Late April to May. Trade designation: GLENDOICK GOLD. (Glendoick is a registered trademark belonging to Glendoick Gardens).

(r) '**Gle017**'

cv. *Elepidote rhododendron* : *yakushmanum* (s) × *hemsleyanum*. H: P.A. & K.N.E. Cox (pre 2003), G: P.A. & K.N.E. Cox, N: P.A. & K.N.E. Cox, I: Glendoick Gardens (2003), REG: P.A. & K.N.E. Cox (2007). Fls 15–20/globular truss, open funnel-shaped, 50–60 × 70–80mm, with 6 wavy-edged lobes, buds white, opening pure white throughout, with yellow-green spots in the throat; vanilla-scented. Calyx vestigial. Lvs ovate, 120 × 55mm, margins decurved, matt green above, pointed at apex, with a very thin layer of hairs that is easily rubbed off. Shrub 1.5 × 2.0m in 15 years. Early to mid June. Trade designation: GLENDOICK VANILLA. (Glendoick is a registered trademark belonging to Glendoick Gardens).

(a) '**Gle022**'

cv. Evergreen azalea : 'Eisprinzessin' (s) × un-named red (no.94–3A). H: P.A. & K.N.E. Cox (1997), G: P.A. & K.N.E. Cox (2000), N: P.A. & K.N.E. Cox, I: Glendoick Gardens, REG: P.A. & K.N.E. Cox (2007). Fls semi-double ('with fused stamens'), 3–6/more-or-less rounded truss, open campanulate, 10 × 25mm, with 5 wavy-edged lobes, deep to strong purplish pink (55A–B), marked, lined and spotted deep purplish pink (55A). Calyx 2–3mm long, green. Lvs elliptic, 15 × 7–8mm, margins slightly decurved, cuneate at base, with some hairs. Shrub 0.6 × 0.8m in 10 years. May. Trade designation: GLENDOICK ROSEBUD. (Glendoick is a registered trademark owned by Glendoick Gardens).

(r) **GLENDICK BUTTERSCOTCH**  
Trade designation for 'GLE003'.

(r) **GLENDICK GOLD**  
Trade designation for 'Gle0011'.

(a) **GLENDICK ROSEBUD**  
Trade designation for 'Gle022'.

(r) **GLENDICK VANILLA**  
Trade designation for 'Gle017'

(r) **'Glenn O'Sheal'**  
cv. Elepidote rhododendron : 'Trilby' (s) × *aberconwayi*. H: D.B. Johnson (1985), G: D.B. Johnson (1993), N: D.B. Johnson (2005), REG: D.B. Johnson (2007). Fls 13/ domed truss, open funnel-shaped, 45 × 75mm, with 5 wavy-edged lobes, strong purplish red (64B) in bud, opening inside strong purplish pink (63C); outside strong purplish red (63B) at margins, shading to strong purplish pink (63C) at the base; large, white (155D), dorsal flare. Lvs elliptic, 175 × 65mm, broadly acute at apex, rounded at base, margins wavy, dull and moderate olive green (137A) above, hairless. Shrub 2.1 × 1.8m in 18 years. Early May.

(a) **'Glittering Prize'**  
cv. Deciduous azalea : parentage unknown. H: W.G.T. Hyde (late 1960s), G: W.G.T. Hyde (late 1970s), N: Mrs R.G. Legrand (2007), I: Mrs R.G. Legrand (2007), REG: Mrs R.G. Legrand (2007). Fls 10–15/compact truss, open trumpet-shaped, 70 × 37mm, with 5 wavy-edged lobes, buds brilliant orange-yellow (21B), shaded vivid reddish orange (30B), opening vivid orange-yellow (21A), shaded vivid yellowish orange (30C), outside brilliant orange-yellow (21B), shaded vivid yellowish orange (30C); with dorsal speckling which is dark greenish yellow (152C) near the centre, shading out to vivid orange-yellow (23A) towards the apex of the dorsal lobe. Calyx 15mm long, strong yellow-green (144A). Lvs narrowly elliptic, 95 × 35mm, margins flat, cuneate at base, acute at apex, glossy green above, hairless. Shrub 1.4 × 1.15m in 30 years. May to early June.

(r) **'Glynneden Dawn Chorus'**  
cv. Elepidote rhododendron : 'Nancy Evans' (s) × 'Exbury Naomi'. H: W.A. Stipe (1992), G: W.A. Stipe (1997), N: W.A. Stipe (2002), I: Glynneden Gardens (2005), REG: W.A. Stipe (2007). Fls 14/domed truss, broadly funnel-shaped, 75 × 100mm, with 5 wavy-edged lobes, dark red (59A) in bud, opening inside vivid yellow (14B), with sparse, deep yellowish pink (39B) dorsal spotting; outside light yellow (20B), with moderate yellowish pink (38B) markings. Calyx 40mm long,

moderate yellowish pink (38B). Lvs elliptic, 125 × 50mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive-green (137A) above, hairless. Shrub 1.4 × 1.1m in 9 years. Mid May.

(v) **'Golden Retriever Mandy'**  
cv. Lepidote rhododendron : ('Gardenia Odyssey' × *laetum*) (s) × ('Gardenia Odyssey' × *laetum*). H: G.L.S. Snell (2001), G: Pacific Island Nursery (2006), N: R. Kruppa (2007), I: Pacific Island Nursery, REG: R. Kruppa (2007). Fls 6–7/flat truss, tubular funnel-shaped, 57 × 65mm, with 5 wavy-edged lobes, brilliant yellow-green (149B) in bud, opening inside vivid yellow (14A), aging in 9 days to strong orange-yellow (17A); outside vivid yellow (14A), aging in 10 days to vivid orange-yellow (21A); unmarked. Calyx minute, green. Lvs elliptic to obovate, 108–120 × 35–51mm, acuminate at apex, cuneate at base, margins upcurved, semi-glossy and moderate olive green (137A) above, moderate yellow-green (138B) below, with small dark scales below (not visible to the naked eye). Fls long-lasting (8–10 days on the plant, 12–15 days as cut flowers). Mid August and perhaps also in February to March. (Known locally as "(Mr Snell's) 'Meadow Yellow' F2").

(r) **'Great Gatsby'**  
cv. Elepidote rhododendron : 'Janet Blair' (s) × 'Phipps 84'. H: W. Brack (1986), G: W. Brack (1995), N: W. Brack (2007), REG: W. Brack (2007). Fls 10/domed truss, broadly funnel-shaped, 65 × 125mm, with 6 wavy-edged lobes, pale purplish pink (56C) in bud, opening white inside, with 6 strong red (46A) flares in the throat and strong red (46A) dorsal speckling, all on a light greenish yellow (8C) background; outside white. Calyx 3mm long, strong yellow-green (144A). Lvs obovate, 140 × 65mm, obtuse at apex, rounded at base, margins flat, dull and moderate olive green (146A) above, hairless. Dense shrub, 2.1 × 1.6m in 21 years. Mid May.

(r) **'Honey Mist'**  
cv. Elepidote rhododendron : 'Invitation' (s) × ('Satin Gold' × (('Whopper' × 'Lem's Cameo') × ('Whopper' × 'Brandt's Tropicana'))). H: J. Barlup (1989), G: J. Barlup (1994), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 18/ball truss, broadly funnel-shaped, c.50 × 75mm, with 7 wavy-edged lobes, moderate purplish red (58A) in bud, opening inside pale yellow (11C), with a 10mm wide pale purplish pink (62D) edging; outside similar, with strong purplish pink (62A) midveins; deep red (53A) nectaries extending c.10mm from the base. Calyx 40mm long, pale yellow (11C), with pale purplish pink (62D) tips and two

deep red (53A) nectaries. Lvs elliptic, 100 × 45mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.0 × 1.3m in 16 years. Late April.

(r) '**Howard Kline**'

cv. *Elepidote rhododendron* : ('Coral Velvet' × *wardii* (KW4170)) (s) × 'Wizard'. H: D. MacMullan (1983), G: D. MacMullan (1987), N: D. MacMullan (2006), REG: D. MacMullan (2007). Fls 19/ball truss, broadly funnel-shaped, 90 × 75mm, with 5 wavy-edged lobes, deep orange-red in bud, opening inside moderate pink (49B) at margins, shading to strong pink (49A) at the base, with a spotted flare of vivid yellow (12A) to brilliant yellow (12B) on the dorsal lobe; outside deep pink (52B) with light orange-yellow (19A) at the base of the tube, shading outwards to deep yellowish pink (39B) at the margins: the overall effect is peach-coloured. Lvs elliptic, 125 × 40mm, broadly acute at apex, oblique at base, margins flat, dull and mid green above, hairless. Shrub 1.2 × 1.2m in 10 years. Mid to late May.

(r) '**Iona Campagnolo**'

cv. *Elepidote rhododendron* : 'Etta Burrows' (s) × 'Fusilier'. H: J. Round (1986), G: G. Wedemire, N: C. Dancer & N. Todd (2007), REG: N. Todd (2007). Fls 20/domed truss, open campanulate, 65 × 75mm, with 5 wavy-edged lobes, vivid red (46C) in bud, opening inside deep yellowish pink (46D), with a strong red (46A) basal blotch; outside deep yellowish pink (46D); anthers black. Calyx c.5mm long, strong red (46A). Lvs lanceolate, 250 × 70mm, acuminate at apex, rounded at base, margins flat, semi-glossy and dark yellowish green (139A) above, with very sparse, moderate reddish brown (166B) hairs on the midvein. Shrub 2.7 × 2.7m in 30 years. Early to midseason (March to April). (Etymology: named for the Governor General of British Columbia, Canada).

(r) '**Jane Gardner**'

cv. *Elepidote rhododendron* : 'Lem's Cameo' (s) × 'Goldfinger'. H: L.A. Hutchison (1992), G: L.A. Hutchison (1996), N: L.A. Hutchison (2006), REG: L.A. Hutchison (2007). Fls 12–13/loose ball truss, broadly funnel-shaped, 60 × 75–100mm, with 6–7 wavy-edged lobes, deep yellowish pink (47C) in bud, opening inside strong pink (48D) shading to light yellowish pink (38D) in lobe centres, with the throat below the dorsal lobes light orange-yellow (25D), outside strong red (47B) gradually shading to deep pink (48A) and to pale yellowish pink (36D) at the base; dorsal lobes with numerous, small, light orange (28C) spots; the colour both

inside and out tends to fade with maturity and is displayed better in semi-shade; style light yellow-green (150D), stigma strong yellow-green (144B). Calyx 15–20mm long, petaloid, strong pink (48C), with areas of pale yellowish pink (36D). Lvs elliptic, 100–120 × 40–50mm, margins slightly decurved, matt moderate olive green (147A) above, narrowly rounded at base, apiculate at apex, hairless. Reasonably compact shrub, 1.25 × 1.20m in 13 years. Early October to early November (New Zealand). (Nomenclatural standard: colour print provided by the registrant (WSY0096693). Etymology: named for a friend of the registrant who had admired the plant in his garden. The identity of the pollen parent ('Goldfinger') is uncertain: the pollen was obtained from a plant so-labelled in another garden in New Zealand but its location has now been lost. It seems likely to be an unrecorded use of this name as the registered 'Goldfinger' is a lepidote burmanicum hybrid).

(v) '**Jennifer Jean**'

cv. *Lepidote rhododendron* : selection from *gardenia*, collected by G. Argent in Papua New Guinea (RBGE accession no.19761327). S: Royal Botanic Garden, Edinburgh, N: G. Argent (2007), REG: Royal Botanic Garden, Edinburgh (2007). Fls 2–4/open truss, trumpet-shaped, 90–110 × 110–140mm, with 5–8 wavy-edged lobes, opening brilliant yellow-green (150C), with faint pink spots at the base of each lobe; strongly sweet-scented. Calyx 1–2mm long, green. Lvs elliptic to broadly elliptic, 80–150 × 55–90mm, margins decurved, broadly tapered to rounded at base, obtuse to rounded at apex, with a scaly indumentum (scales silvery at first, aging to brown). Shrub 1.5 × 1.0m in 10 years. Flowering at irregular intervals. AM 2007. (Etymology: named for the collector's daughter. Nomenclatural standard: herbarium specimen of the award plant (WSY0100899)).

(r) '**Jim's Black Heart**'

cv. *Elepidote rhododendron* : 'Wine Fuschia' (s) × Kilimanjaro Group. H: J.S. Smith (2000), G: J.S. Smith (2003), N: J.S. Smith, I: Roadside Rhododendrons (2007), REG: J.S. Smith (2007). Fls 21–25/tight, rigid, conical truss, funnel-campanulate, 65 × 90mm, with 5 wavy-edged lobes, waxy, opening inside dark red (59A), with prominent dark red (187A) speckling on the dorsal lobe and at the base; outside dark red (187B). Lvs oblong, 240 × 65mm, broadly acute at apex, rounded at base, with downcurved margins, glossy and moderate olive green (147A) above, hairless. Shrub 1.2 × 0.9m in 5 years. Late May.

(v) **'John Rouse'**

cv. Lepidote rhododendron : selection from *rousei*, collected on Sibuyan Island, Philippines by G. Argent (RBGE accession no. 19902332). S: Royal Botanic Garden, Edinburgh, N: G. Argent (2007), REG: Royal Botanic Garden, Edinburgh (2007). Fls 2–6/ open truss, funnel-shaped, 22–28 × 45–57mm, with 5 wavy-edged lobes, opening white (155D) throughout, unmarked; not scented. Calyx 1mm long, green. Lvs elliptic to obovate, 25–50 × 15–30mm, margins decurved, tapered at base, obtuse to rounded at apex, with a scaly indumentum (scales silver at first, maturing to brown). Shrub 0.3 × 0.25m in 5 years. Flowering irregularly but at least two times a year. PC 2007. (Etymology: named for the late John Rouse, a well-known breeder and enthusiast for vireya rhododendrons in Australia. Nomenclatural standard: herbarium specimen of the award plant (WSY0100900)).

(a) **'Karee Lynn'**

cv. Evergreen azalea : ('Mrs Nancy Dippel' × 'Satellite') (s) × 'Mrs Nancy Dippel'. H: Robert Stewart (1997), G: Robert Stewart (2000), N: Robert Stewart (2006), REG: Robert Stewart (2007). Fls double, 3/truss, broadly funnel-shaped, 40 × 65mm, with very wavy-edged lobes, deep yellowish pink (43C) in bud, opening slightly paler than deep yellowish pink (43C) throughout, with sparse, deep yellowish pink (43C) spotting. Calyx c.15mm long, brilliant yellow-green (142B). Lvs elliptic, 60 × c.25mm, broadly acute at apex, cuneate at base, margins upcurved, semi-glossy and moderate yellow-green (146B) above, hairless. Shrub 0.5 × 0.8m in 9 years. Midseason.

(r) **'Karen Gervais Beauty'**

cv. Elepidote rhododendron : 'Anna' (s) × 'Purple Lace'. H: W.A. Stipe (1991), G: W.A. Stipe (1995), N: S. Stipe (2007), I: Glynneden Gardens (2007), REG: W.A. Stipe (2007). Fls 21/domed truss, broadly funnel-shaped, 75 × 75mm, with 5 very wavy-edged lobes, dark purple (79A) in bud, opening inside strong purplish red (60B), with dark purple (79A) spots and blotch in the throat; outside deep red (60A). Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins downcurved, semi-glossy and dark yellowish green (139A) above, hairless. Shrub 1.8 × 1.5m in 15 years. Mid May.

(v) **'Kingston Blush'**

cv. Lepidote rhododendron : parentage unknown. H: unknown, G: A.B. Raper, N: K. & G. Hagan (2007), REG: K. & G. Hagan (2007). Fls 20/flat truss, tubular funnel-shaped, 70 × 115mm, with 7 slightly wavy-edged lobes, light yellow (18B) in

bud, opening moderate yellowish pink (37C) with a suffusion of light yellow (18B) towards the base which also extends up the centre of the lobes; outside light pink (49C); unmarked; strongly scented. Lvs oblong, 115 × 55mm, margins flat, glossy above, cuneate at base, acute at apex, hairless. Shrub 1.0 × 1.0m. Autumn. (Nomenclatural standard: colour print provided by the registrants (WSY0100242). Etymology: derived from the home town of the registrants (Kingston Beach) and the flower colour.)

(a) **'Kirishima-sakura'**

cv. Evergreen azalea: selection from *kiusianum*. G: H. Maeda (2003), S: H. Maeda (2004), N: H. Maeda (2005), REG: H. Maeda (2007). Fls 3–5/hemispherical truss, shortly tubular funnel-shaped, 25 × 28mm, with 5 slightly wavy-edged lobes, white in bud, opening inside very pale pink, outside pink and white, unmarked. Calyx 2–3mm long, light yellow-green. Lvs elliptic to obovate, 12 × 6mm (in spring), margins decurved, cuneate at base, obtuse at apex, sparsely hairy; summer lvs partially persistent, when they are brownish green. Small compact shrub, 40 × 45cm in 7 years. Late May.

(r) **'Kristie Lynn'**

cv. Elepidote rhododendron : 'Karen Triplett' (s) × 'Big Deal'. H: G. Woodard (1999), G: G. Woodard (2002), N: K. Sisson (2007), I: G. Woodard, REG: G. Woodard (2007). Fls 12–15/conical truss, broadly funnel-shaped, 50 × 50mm, with 6–7 wavy-edged lobes, pale yellow (11C) in bud, opening pale yellow (11C) throughout; unmarked. Calyx c.15mm long, strong yellowish green (141C). Lvs elliptic, 175 × 65mm, broadly acute at apex, cuneate at base, margins downcurved, glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 5 years. Midseason.

(r) **'Leggs'**

cv. Elepidote rhododendron : 'Arthur Bedford' (s) × 'La Bar's White'. H: A.T. & S.A. Anderson (1981), G: A.T. & S.A. Anderson (1990), N: A.T. & S.A. Anderson (2006), REG: A.T. & S.A. Anderson (2007). Fls 19/conical truss, broadly funnel-shaped, 45 × 70mm, with 5 wavy-edged lobes, vivid purple (81B) in bud, opening inside light purple (80D) and very pale purple (76D), with a prominent dorsal blotch of dark purple (79A) spots, outside light purple (80C). Lvs elliptic, 125 × 50mm, acute at apex, cuneate at base, margins flat, dull and moderate olive green (137A) above, hairless. Shrub 1.5 × 1.2m in 10 years. Late May.

(r) '**Lemon Zinger**'

cv. Elepidote rhododendron : 'Original Vagabond' (s) × 'Fashion Plate'. H: R.H. Gustafson (2001), G: R.H. Gustafson (2005), N: R.H. Gustafson (2006), REG: R.H. Gustafson (2007). Fls 16/domed truss, broadly funnel-shaped, c.75 × 100mm, with 6 wavy-edged lobes, light greenish yellow (3C) in bud, opening inside brilliant greenish yellow (1B), with the three dorsal lobes deepening to brilliant greenish yellow (1A) in the throat and with a strong purplish red (61B) marking deep in the throat; outside light greenish yellow (3C), fading to light greenish yellow (3D) at edges; strong greenish yellow (153B) spotting on the dorsal lobes; slightly scented. Calyx c.1.5mm long, light yellow-green (144B). Lvs oblong, 175 × 75mm, obtuse at apex, oblique at base, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.2 × 0.9m in 5 years. Mid May.

(r) '**Little Susie Q**'

cv. Elepidote rhododendron : ('Scintillation' × Inamorata Group) (s) × (yellow-flowered selection from *fortunei* × (*dichroanthum* × *wardii*)). H: R. Murcott (1965), G: R. Murcott (1971), N: S. Baricilli (2005), REG: S. Baricilli (2007). Fls 13/ball truss, broadly funnel-shaped, 50 × 90mm, with 7 wavy-edged lobes, deep purplish pink (54B) in bud, opening inside pale yellowish pink (36D), with pale yellow (11D) deep in the base; outside moderate yellowish pink (38B); unmarked. Calyx 1mm long, strong yellowish green (134B). Lvs elliptic, 100 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and vivid yellowish green (140A) above, hairless. Shrub 1.8 × 1.8m in 12 years. Midseason.

(r) '**Loch Linnhe**'

cv. Elepidote rhododendron : 'Nancy Evans' (s) × 'Percy Wiseman'. H: P.A. & K.N.E. Cox, G: P.A. & K.N.E. Cox, N: P.A. & K.N.E. Cox, I: Glendoick Gardens, REG: P.A. & K.N.E. Cox (2007). Fls 13/truss (truss full, but slightly wider than high), 50 × 80mm, with 5(-6) wavy-edged lobes, buds strong red (53B), opening inside deep purplish pink (54B) on the lobes shading to light yellow (14D) in the throat and with a lobed ring of deep purplish pink (54B) right at the base of the throat; outside similarly coloured. Calyx 3mm long. Lvs narrowly elliptic, 80-110 × 30-45mm, rounded at base, cuneate at apex, hairless. Shrub 1.0 × 1.3m in 7 years. April to May.

(r) '**Lockington Idyll**'

cv. Elepidote rhododendron : 'Miss B.L. Jones' (s) × 'Petra's Debut'. H: D.J. Dosser (1999), G: D.J. Dosser (2005), N: D.J. Dosser (2006), REG: D.J. Dosser (2007). Fls 14/

ball truss, funnel-shaped, 50 × 100mm, with 5 wavy-edged and partially recurved lobes, light purplish pink (73C) in bud and throughout on opening, with a strong pink (50C), spotted, dorsal blotch; slightly scented. Lvs elliptic, 150 × 50mm, margins decurved, glossy above, truncate at base, broadly acute at apex, hairless. Shrub 1.0 × 1.0m in 7 years. November (Australia). (Nomenclatural standard: colour print provided by the registrant (WSY0096514)).

(r) '**Lover's Blush**'

cv. Elepidote rhododendron : 'Raspberry and Custard', open-pollinated. H: M. Sanders (1999), G: M. Sanders (2003), N: M. Sanders (2007), I: Sanders Nursery, REG: M. Sanders (2007). Fls 12-18/domed truss, open funnel-shaped, 50 × 65mm, with 5 wavy-edged lobes, white (155D) in bud, opening white (155D) and darkening gradually over a 10 day period to strong purplish pink (68B), with a dorsal flare of discrete, vivid purplish red (66A) speckles; slightly scented. Lvs elliptic, 130 × 60mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and mid green above, hairless. Shrub 0.8 × 0.9m in 8 years. Mid April.

(r) '**Mac's Magic**'

cv. Elepidote rhododendron : ('Kruse's Big Cream' × *dichroanthum*) (s) × ('Janet Blair' × 'Seattle Gold'). H: D. MacMullan (1972), G: D. MacMullan (1976), N: D. MacMullan (2006), REG: D. MacMullan (2007). Fls 16/ ball truss, broadly funnel-shaped, with 5 wavy-edged lobes, light yellowish pink (19B) to light orange-yellow (19A) in bud, opening inside brilliant greenish yellow (2B) in centre, shading to vivid greenish yellow (2A) at the margins; outside brilliant greenish yellow (2B) to light yellow-green (2C); unmarked. Lvs oblong, 200 × 90mm, obtuse at apex, rounded at base, margins flat, glossy and mid green above, hairless. Dense shrub, 1.5 × 1.5m in 10 years. Midseason.

(r) '**Mac's Solar Surprise**'

cv. Elepidote rhododendron : 'Janet Blair' (s) × ('Pacific Gold' × 'Phipps Yellow'). H: D. MacMullan (1980), G: D. MacMullan (1985), N: D. MacMullan (2005), REG: D. MacMullan (2007). Fls 9/ball truss, broadly funnel-shaped, 125 × 115mm, with 5 very wavy-edged lobes, opening inside deep yellow/orange in the centre, shading to lighter yellow towards the lobe margins, with light orange spotting near the edge; outside shades of light yellow. Calyx c.13mm long, light yellow. Lvs elliptic, 150 × 75mm, obtuse at apex, rounded at base, margins flat, semi-glossy and mid fir green above, hairless. Shrub 1.2 × 1.5m in 10 years. Midseason.

(r) **'Maisie'**

cv. Lepidote rhododendron : *moupinense* (s) × *chrysodoron*. H: J. Parsons (1999), G: F.J. Williams & J. Parsons (2002), N: J. Parsons (2007), REG: C.H. Williams & J. Parsons (2007). Fls 4/open truss, funnel-shaped, 50 × c.60mm, with 5 wavy-edged lobes, buds brilliant yellow-green (154C), becoming tinged with deep pink (52B) at apex, opening inside pale greenish yellow (2D), slightly tinged deep to strong pink (52C–D) at the tips of the dorsal lobes, outside pale greenish yellow (2D), flushed deep to strong pink (52C–D) on the dorsal lobes and on the dorsal side of the tube; large, strong to moderate yellowish pink (31C–D) spots in the dorsal throat and just extending onto the base of the dorsal lobe; style yellowish white, just tinged pink towards the apex, stigma pale green; anthers pinkish brown, filaments yellowish white, hairy at the base; ovary densely scaly. Calyx c.2mm long, green at base, pinkish above and with a hyaline, ciliate margin, with numerous, tiny, pale scales. Lvs elliptic, up to 100 × 38mm, margins flat to slightly downcurved, cuneate to broadly cuneate at base, acute and apiculate at apex, with numerous, non-overlapping, pale brown scales beneath, much less scaly above. Shrub 0.6m in 7 years. Mid March. (Nomenclatural standard: herbarium specimen provided by the registrants (WSY0100146). Etymology: named in memory of the grandmother of the hybridiser. The pinkish tinge to the flowers is sometime poorly developed.)

(r) **'Margaret S. Coe'**

cv. Elepidote rhododendron : 'Parker's Pink' (s) × 'Martha Phipps'. H: R. Murcott (1990), G: R. Murcott (1997), N: V. Simeone, REG: R. Murcott (2007). Fls 9/ball truss, broadly funnel-shaped, c.70 × 100mm, with 6 wavy-edged lobes, inside edge strong purplish pink (55B), shading to pale purplish pink (55D), with light yellow (11B) speckled markings in the centre; outside pale purplish pink (55D). Calyx 3mm long. Lvs elliptic, 140 × 65mm, obtuse at apex, rounded at base, margins flat, dull and moderate yellow-green (137C) above, hairless. Dense shrub, 1.5 × 2.4m in 16 years. Mid May. (Etymology: named for a member of the family who donated the land that became Planting Fields Arboretum (Oyster Bay, New York, USA)).

(a) **'Martha Ann Lucy'**

cv. Evergreen azalea : parentage unknown. H: W.G.T. Hyde (1960s), G: W.G.T. Hyde (1970s), N: Mrs R.G. Legrand (2007), I: Mrs R.G. Legrand (2007), REG: Mrs R.G. Legrand (2007). Fls 1–2/truss, open trumpet-shaped, 44 × 20mm, with 5 flat-edged lobes, light reddish purple (74C) in bud and throughout

on opening; with strong purplish red (63A) speckles on the three dorsal lobes. Calyx 5mm long, moderate yellow-green (146D). Lvs ovate, 55 × 25mm, margins flat, cuneate at base, acute at apex, hairless. Shrub 0.6 × 0.9m in 30 years. May. (Etymology: named in memory of the stillborn baby of the registrant's god-daughter).

(a) **'Midsummer Mermaid'**

cv. Deciduous azalea : *viscosum* hybrid, grown from seed supplied as *viscosum* by the American Rhododendron Society. G: E.G. Millais (1995), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 7/small, domed truss, long tubular funnel-shaped, with 5 smooth-edged lobes, 30 × 35mm, strong red (51A) in bud, inside strong pink (54C) to moderate purplish pink (54D), with the dorsal lobe brilliant yellow (13C) in the centre fading to pale orange-yellow (16D) near the edges; outside strong red (51A) on tube extending as stripes on lobes which have light purplish pink (55C) edges; stamens 5, exerted, filaments, like the style, deep purplish pink (54B) towards the apex, paler below; tube glandular hairy. Calyx 2mm long, pale yellow-green, ciliate and hairy. Lvs obovate to oblanceolate, 30–40 × 12–14mm, margins decurved, matt blue-green, attenuate at base, obtuse to broadly acute at apex, apiculate, ciliate and hairy on the main vein above and below. Shrub 1.2 × 1.0m in 10 years. Late June. (Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100733). This is the plant listed by Ted Millais as "VBF" in his article on *Rhododendron viscosum* and late-flowering azaleas in *Rhododendrons with Camellias and Magnolias* (2005)).

(v) **'Mino'**

cv. Lepidote rhododendron : *konori* (s) × ('Doctor Hermann Sleumer' × (*macgregoriae* × *aurigeranum*)). H: R.A. Mitchell (1999), G: R.A. Mitchell (2006), N: R.A. Mitchell (2006), REG: R.A. Mitchell (2007). Fls 5/flat truss, tubular funnel-shaped, 115 × 120mm, with 7 wavy-edged lobes, opening inside light purplish pink (68B); outside light purplish pink (68D) at margins, changing to strong purplish red (58B) going down the tube. Calyx 10mm long. Lvs elliptic, 115 × 65mm, broadly acute at apex, cuneate at base, margins flat, semi-glossy and dark yellowish green (139A) above, without obvious indumentum. Shrub with an open habit, 1.2 × 0.9m in 7 years. Late January.

(r) **'Miss Diane'**

cv. Elepidote rhododendron : 'Whitney Late Orange' (s) × (*sutchuenense* × Prelude Group) × 'Ted's Yellow'. H: D.B. Johnson (1995), G: D.B. Johnson (2002), N: D.B. Johnson

(2006), REG: D.B. Johnson (2007). Fls 14/ domed truss, open funnel-shaped, 65 × 90mm, with 7 wavy-edged lobes, light pink (50D) in bud, opening inside pale yellow-green (4D), shading to light greenish yellow (4C) at base; outside similarly coloured; unmarked. Lvs obovate, 150 × 60mm, obtuse at apex, cuneate at base, margins downcurved, dull and moderate olive green (137A) above, hairless. Shrub 1.8 × 1.5m in 12 years. Early May.

(v) **'Mitch Mitchell'**

cv. *Lepidote rhododendron* : 'Cephas' (s) × *zoelleri* (Bovee's V300). H: R.A. Mitchell (2001), G: S. Bertelmann (2005), N: S. Bertelmann (2006), I: Pacific Island Nursery (c. 2008), REG: S. Bertelmann (2007). Fls 5/flat truss, tubular funnel-shaped, 100 × 110mm, with 5 wavy-edged lobes, inside deep yellowish pink (43C) surrounding a large, central, star-shaped marking of pale yellowish green (4D), shading to light greenish yellow (4B) in the throat; outside of tube light greenish yellow (8C), with broad, deep pink (48A) margins; slightly scented. Lvs elliptic, 140 × 65mm, obtuse at apex, cuneate at base, margins upcurved, semi-glossy and dark yellowish green (139A) above. Shrub 1.1 × 0.6m in 5 years.

(r) **'Moonlite Champagne'**

cv. *Elepidote rhododendron* : *maccabeanum* (s) × ('Yaku Sunrise' × ('C.I.S.' × 'Jingle Bells')). H: W.A. Stipe (1995), G: W.A. Stipe (2000), N: W.A. Stipe (2006), I: Glynneden Gardens (2007), REG: W.A. Stipe (2007). Fls 12/domed truss, funnel-campanulate, 65 × 65mm, with 7 flat-edged lobes, deep red (60A) in bud, opening pale greenish yellow (2D) throughout, with a dark red (59A) dorsal blotch. Lvs elliptic, 150 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairy below and on midrib (hairs greyish yellow-green (195B) at first, aging to moderate yellow (161B)). Shrub 1.2 × 1.5m in 7 years. Mid May.

(r) **'Nadine Cook'**

cv. *Lepidote rhododendron* : selection from *augustinii*. G: E.J. Greig (1970?), S: A.R. Cook (2003), N: A.R. Cook (2003), REG: A.R. Cook (2007). Fls 2–6/truss, 40 × 50mm, with 5 flat-edged lobes, light violet (92B), shading to very pale purple (92D), unmarked. Lvs oblong-lanceolate, 65 × 25mm, acuminate at apex, cuneate at base, margins flat, dull green above, scaly. Dense shrub, 4.5 × 2.4m in 25 years. Mid April to May.

(r) **'Olivia'**

cv. *Elepidote rhododendron* : parentage unknown. H: W.G.T. Hyde (1960s), G:

W.G.T. Hyde (1970s), N: Mrs R.G. Legrand (2007), I: Mrs R.G. Legrand (2007), REG: Mrs R.G. Legrand (2007). Fls 7–9/domed truss, open trumpet-shaped, 55 × 30mm, with 7 wavy-edged lobes, buds deep pink (51B), opening white (155A) throughout, the centre shaded strong greenish yellow (151A) and with speckles of dark greenish yellow (152C) extending up to 30mm from the centre; scented. Calyx 4mm long, moderate yellow-green (148B). Lvs narrowly elliptic, 135 × 65mm, margins decurved, obtuse at base, acute at apex, matt green above, hairless. Shrub 3.0 × 3.0m in 30 years. May. (Etymology: named as part of a collection with Shakespearian names).

(a) **'Opossum'**

cv. Evergreen azalea : 'Purpurtraum' (s) × 'Blue Danube'. H: P.A. & K.N.E. Cox (1997), G: P.A. & K.N.E. Cox (1999), N: P.A. & K.N.E. Cox (2007), I: Glendoick Gardens, REG: P.A. & K.N.E. Cox (2007). Fls 3–5/truss, funnel-campanulate, 30 × 40–50mm, buds between deep purplish red and vivid reddish purple (71A & 74A), opening between deep purplish red and vivid reddish purple (71A & 74A) throughout, with darker spots on the dorsal lobe. Calyx 3mm long, green. Lvs narrowly obovate, 10–15mm long, margins flat to slightly upcurved, cuneate at base, more-or-less obtuse at apex, with a few hairs. Shrub 0.6 × 0.8m in 10 years. May. Very similar to 'Blue Danube' but overall more compact.

(r) **'Orange Lemonade'**

cv. *Elepidote rhododendron* : 'Goldsworth Orange', open-pollinated. H: D.B. Johnson (1985), G: D.B. Johnson (1991), N: D.B. Johnson (2006), REG: D.B. Johnson (2007). Fls 12/domed truss, funnel-shaped, 50 × 90mm, with 5 wavy-edged lobes, moderate reddish orange (35B) in bud, opening inside and out light yellowish pink (29C) at margins, shading to moderate reddish orange (35B) at base; unmarked. Lvs elliptic, 150 × 50mm, broadly acute at apex, oblique at base, margins wavy, dull and moderate olive green (137B) above, hairless. Shrub 1.5 × 1.4m in 15 years. May.

(r) **'Orange 2003'**

Syn. of 'Orange Prelude'.

(r) **'Orange Prelude'**

cv. *Elepidote rhododendron* : 'Lem's Tangerine' (s) × ('Lem's Cameo' × (('Lem's Cameo' × 'Ken Janek') × 'Brandt's Tropicana') × ('Whopper' × 'Brandt's Tropicana')). H: J. Barlup (1998), G: J. Barlup (2003), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 9/ball truss, saucer-shaped, c.60 × 90mm, with 7 very wavy-edged lobes, vivid reddish orange (44C) in bud, opening inside light

orange-yellow (16B) in centre, blending after c.25mm to deep yellowish pink (39B) lobes which have strong red (39A) tips; outside light orange-yellow (16B) at base extending to c.20mm, blending to strong red (39A) on all lobes, with strong reddish orange (42B) veining on all midribs; unmarked. Calyx 25mm long, vivid reddish orange (44C), blending to strong red (39A). Lvs elliptic, 150 × 65mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub c.1.0m × 1.8m in 8 years. Early May. Syn: 'Orange 2003'.

(v) '**Paddy Woods**'

cv. *Elepidote rhododendron* : selection from *macgregoriae*, collected as Woods 3072 in Papua New Guinea by Paddy Woods (RBGE accession no. 19682641). S: Royal Botanic Garden, Edinburgh, N: G. Argent (2007), REG: Royal Botanic Garden, Edinburgh (2007). Fls 10–15/umbellate truss, shortly tubular campanulate, 10–25 × 20–30mm, with 5 flat-edged lobes, opening brilliant yellow (8A) both inside and out, unmarked; anthers brown. Calyx 1mm long, green. Lvs elliptic to ovate-elliptic, blades 45–65 × 20–30mm, shortly acuminate at apex, with a scaly indumentum (scales silver when young, maturing to pale brown); petioles c.10mm long. Shrub 1.0 × 0.5m in 5 years. Flowering at irregular intervals, mostly in spring. AM 2007. (Etymology: named after the collector of the plant. Nomenclatural standard: herbarium specimen of the award plant (WSY0100903)).

(a) '**Patsy's Pink**'

cv. Deciduous azalea : *colemanii* nom. ined. (s) × 'Gibraltar'. H: R.O. Smitherman (1987), G: T.E. Corley (1992), N: R.O. Smitherman (2007), REG: R.O. Smitherman (2007). Fls 12–14/domed or ball truss, open funnel-shaped, 50 × 55mm, with 5 very wavy-edged lobes, deep purplish pink (66C) in bud, opening the same colour throughout, with a prominent blotch of strong orange (24A) in the basal central portion of the dorsal lobe; cinnamon/clove-scented. Calyx c.3mm long, moderate yellow-green (138C). Lvs elliptic, 55–65 × 30–40mm, acute at apex, cuneate at base, margins flat, glossy and moderate olive green (137B) above, with sparse, moderate olive green (137B) hairs along the midvein below. Shrub 2.1 × 2.4m in 20 years. Mid April to early May.

(r) '**Peach Flambé**'

cv. *Elepidote rhododendron* : 'Reigate Squire' (s) × 'Peach Cobbler'. H: R.H. Gustafson (1992), G: R.H. Gustafson, N: R.H. Gustafson (2007), REG: R.H. Gustafson (2007). Fls 11/truss, broadly funnel-shaped, 70 × 100mm,

with 6 wavy-edged lobes, deep pink (48B) in bud, opening inside strong pink (54C) at edges, shading to light pink (50D), with a light yellowish pink (19B) flare deep in the throat; outside strong pink (54C) at edges, shading to deep pink (48B) at base, with deep pink (48A) midribs; slightly scented. Calyx rim-like, brilliant yellow-green (149B). Lvs elliptic, 130 × 65mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (146A) above, hairless. Shrub 1.7 × 1.5m in 7 years. Mid May.

(r) '**Peal**'

cv. *Elepidote rhododendron* : *griffithianum* (s) × 'Vanessa Pastel'. H: A. Craig-Mooney (1984), G: A. Craig-Mooney (1992), N: A. Craig-Mooney (2006), REG: A. Craig-Mooney (2007). Fls 8/lax truss, funnel-shaped, 60 × 100mm, with 5 wavy-edged lobes, buds vivid red (52A), opening strong pink (49C–D) both inside and out, with suffusions of deep purplish pink (55A) especially around the edges of the lobes and over much of the dorsal lobe. Calyx pink. Lvs lanceolate, 160 × 60mm, acute at apex, cuneate at base, matt green above, hairless. Shrub 2.0 × 1.5m in 20 years. May. (Nomenclatural standard: colour print provided by the registrant (WSY0100750). Etymology: peal is a local name for the sea trout, the colour of whose flesh resembles the colour of the flower).

(a) '**Pine Marten**'

cv. Evergreen azalea : 'Eisprinzessin' (s) × un-named seedling (no.94–3A). H: P.A. & K.N.E. Cox, G: P.A. & K.N.E. Cox, N: P.A. & K.N.E. Cox, I: Glendoick Gardens, REG: P.A. & K.N.E. Cox (2007). Fls single, 4/truss, open campanulate, 20–30 × 60–70mm, with 5 wavy-edged lobes, buds strong purplish pink (55B), opening strong purplish pink (55B) throughout; dorsal lobe with numerous, small, strong red (51A) spots, some spots also on adjacent lobes; stamens and style pink. Calyx 3mm long, green. Lvs ovate, c.20–30 × 10mm, margins slightly upcurved, cuneate at base, acute at apex, fringed with hairs. Shrub 0.6 × 0.6m in 8 years. May.

(r) '**Plum Passion**'

cv. *Elepidote rhododendron* : 'Plum High' (s) × 'Jonathan Shaw'. H: J. Barlup (1997), G: J. Barlup (2000), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 22/ball truss, saucer-shaped, c.55 × 90mm, with 5 wavy-edged lobes, deep purplish red (71A) in bud, opening inside light purple (77D) at base and deepening over c.20mm to light purple (77C) and then to deep reddish purple (77A) on outer 25mm of all lobes; outside light purple



(77C), deepening to strong purple (77B), with deep reddish purple (77A) midribs; with a conspicuous, dark purple (79A), spotted blotch extending upwards c.35mm from the base; anthers prominent, white. Lvs elliptic, 150 × 60mm, broadly acute at apex, rounded at base, margins flat, dull and moderate olive green (147A) above, hairless. Shrub 1.0 × 1.1m in 8 years. Mid May.

(r) **'Plum Satin'**

cv. Elepidote rhododendron : 'Jonathan Shaw' (s) × 'Frosted Plum'. H: J. Barlup (1997), G: J. Barlup (2001), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 21/ball truss, broadly funnel-shaped, 40 × 70mm, with 7 very wavy-edged lobes, moderate purplish red (59C) in bud, opening inside light purple (80C), with slightly lighter light purple (78C) margins; outside similar, with strong purplish red (71B) midveins; conspicuous dorsal blotch of dark red (187A) speckles extending 25mm from the base. Lvs elliptic, 120 × 55mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.9 × 1.2m in 9 years. Early May.

(r) **'Professor Steen Olsen'**

cv. Elepidote rhododendron : *bureavii*, open-pollinated. G: S.A. Askjaer (1991), N: S.A. Askjaer (2000), I: S.A. Askjaer (2007), REG: S.A. Askjaer (2007). Fls 7–10/shallowly domed truss, funnel-shaped, 40–50 × 50–60mm, with 6–7 slightly wavy-edged lobes, buds moderate to pale purplish pink (65A–D), opening inside white (155D), with the basal half of the three dorsal lobes with a dark red (59A) blotch breaking up apically into dark red spots, outside mostly white (155D), with the internal colour showing through on the dorsal lobes; stamens 10–12, filaments 25–30mm long, white, pubescent in the basal quarter, anthers black; style 30–35mm long, white, hairless, stigma pale yellow (160D); ovary pubescent. Calyx 2–3mm long, 6–7-lobed, strong purplish red (60C) with a faint brownish tint. Lvs elliptic, 80–120 × 40–50mm, margins flat, glossy green above, cuneate at base, acute at apex, with a thin indumentum below, the hairs dark greyish yellow (199D) at first, aging to light yellowish brown (199C). Shrub 2.0 × 2.0m in 20 years. Mid to late May. (Nomenclatural standard: colour print provided by the registrant (WSY0096629). Etymology: named for the late Professor Steen Olsen, an expert in renal pathology, and a former colleague of the registrant. Grown from seeds supplied to the ARS seed exchange (ARS no.24/1978) by Mrs F. Darts in 1978.).

(r) **'Prom Queen'**

cv. Elepidote rhododendron : 'Yaku Sunrise' (s) × ('Janet Blair' × ((*wardii* × *hemsleyanum*) × 'Autumn Gold')). H: A.T. & S.A. Anderson (1996), G: A.T. & S.A. Anderson (2003), N: A.T. & S.A. Anderson (2006), REG: A.T. & S.A. Anderson (2007). Fls 10/ball truss, broadly funnel-shaped, 50 × 90mm, with 6–7 wavy-edged lobes, strong purplish pink (55B) in bud, opening inside pale purplish pink (56A) at edges, shading to pale purplish pink (56D) towards the centre, with a brilliant yellow-green (154C) blotch surrounding the ovary; outside pale purplish pink (56A–B). Lvs elliptic, 115 × 50mm, acute at apex, rounded at base, margins flat, semi-glossy and moderate yellow-green (147B) above, hairless. Shrub 0.85 × 0.75m in 10 years. Mid May.

(r) **'Purple Twilight'**

cv. Elepidote rhododendron : 'Jonathan Shaw' (s) × 'Plum High'. H: J. Barlup (2000), G: J. Barlup (2004), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 17/ball truss, broadly funnel-shaped, 50 × 75mm, with 5 wavy-edged lobes, dark red (59A) in bud, opening inside light purple (77D) at base, shading to strong purple (77B) and then deep reddish purple (77A) at edges; outside strong purple (77B) with deep reddish purple (77A) veining on midribs and tips of lobes; prominent dorsal flare of dark red (187A) spots extending c.40mm upwards from the base, with a very pale purple (76D) area extending c.15mm around the top of the flare. Lvs elliptic, 155 × 60mm, broadly acute at apex, rounded at base, margins upcurved, semi-glossy and moderate olive green (147A) above, hairless. Shrub 0.7 × 0.9m in 5 years. Mid May.

(r) **'Radiant Frills'**

cv. Elepidote rhododendron : 'Marquis of Waterford' (s) × 'One Thousand Butterflies'. H: R.H. Gustafson (2001), G: R.H. Gustafson (2004), N: R.H. Gustafson (2006), REG: R.H. Gustafson (2007). Fls 15/ball truss, broadly funnel-shaped, 65 × 75mm, with 5 very wavy-edged lobes, strong red (53C) in bud, opening inside strong purplish red (54A) at margins, shading to pale purplish pink (56C); outside strong purplish red (54A) at margins and on midribs, remainder light purplish pink (55C); brilliant yellow (10A) dorsal flare. Calyx rim-like, strong yellow-green (143C). Lvs elliptic, 125 × 55mm, acute at apex, rounded at base, margins flat, dull and moderate olive green (137A) above, hairless. Shrub 0.9 × 0.6m in 6 years. Late May.

(r) **'Ray Jefferis'**

cv. *Elepidote rhododendron* : *yakushmanum* (s) × unknown. H: R.P. Jefferis (c. 1960–1962), G: P. Livingston (c. 1975), REG: S. Jefferis (2007). Fls 8/domed truss, open funnel-shaped, 65 × 90mm, with 5 wavy-edged lobes, deep pink, almost red, in bud, opening inside pale pink with a darker pink throat; outside darker pink; darker pink markings on midveins outside. Calyx 30mm long, strong yellow. Lvs elliptic, 115–140 × 40–50mm, broadly acute at apex, cuneate at base, margins flat to slightly downcurved, semi-glossy and dark green above, hairy below (the hairs orange at first, aging to yellow-brown). Shrub 0.7 × 0.9m in 20 years. Late May. (Possibly named by Bob Wilkinson (c.1987)).

(r) **'Rose Dew'**

cv. *Elepidote rhododendron* : 'Percy Wiseman' (s) × 'Recital'. H: J. Barlup (1996), G: J. Barlup (2001), N: J. Barlup (2006), REG: J. Barlup (2007). Fls 15/ball truss, broadly funnel-shaped, c.55 × 85mm, with 5 wavy-edged lobes, strong red (53B) in bud, opening inside pale yellow-green (4D), changing to pale purplish pink (56C) and then to strong purplish pink (55B) c.15mm from the edge; outside pale purplish pink (56C) with strong purplish pink (55B) midribs and edges; dark red (59A) flares on all nectaries, c.10mm long on ventral lobes and c.25mm long on dorsal lobes; flares of discrete, dark red (59A) spots extending c.25mm from the base on the dorsal lobes. Lvs elliptic, 110 × 50mm, broadly acute at apex, rounded at base, margins flat, semi-glossy and moderate olive green (147A) above, hairless. Shrub 1.0 × 1.2m in 9 years. Late May.

(a) **'Sea of Stars'**

cv. Deciduous azalea : selection from *viscosum*, grown from American Rhododendron Society seed supplied in 1991. G: E.G. Millais (1995), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 7–12/small, domed, upright truss, long tubular funnel-shaped, up to 50 × 45mm, with 5 slightly wavy-edged lobes, buds deep pink (51B), progressing to pale yellow (12D), opening inside white (155D); outside white (155D), initially with a strong purplish pink (55B) midrib which fades to white; unmarked; stamens 5, exerted, filaments white; style white, tinged pinkish at apex, stigma dark. Outside of the corolla with dense, short, white, eglandular hairs and scattered longer hairs with a dark purplish gland at the tip. Calyx up to 4mm long, variable in length, pale yellow-green, glandular ciliate and glandular hairy. Lvs narrowly oblanceolate, up to 52 × 16mm, margins decurved, not wavy, attenuate at

base, acute and apiculate at apex, grey-blue, ciliate and with sparse, pale, eglandular hairs on the midvein below. Shrub 1.5 × 1.0m in 10 years, with very pale green young shoots, pedicels and petioles. Late June to early July. (Etymology: named in allusion to the effect of the white star-like flowers over a plant with sea blue foliage. Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100735)).

(v) **'Sheila Collette'**

cv. *Lepidote rhododendron* : selection from *polyanthemum*, collected as Collette 73/79 in Sabah by S. Collette (RBGE accession no. 19792883). S: Royal Botanic Garden, Edinburgh, N: G. Argent (2007), REG: Royal Botanic Garden, Edinburgh (2007). Fls 25–30/domed truss, funnel-shaped, 30–35 × 40–48mm, with 5 flat-edged lobes, opening strong reddish orange (32B) throughout, unmarked; scented. Calyx 1mm long, green. Lvs elliptic to broadly elliptic, 70–150 × 40–90mm, margins flat, broadly tapering to rounded at base, broadly tapering at apex, with a scaly indumentum (scales silver at first, maturing to brown). Shrub 1.5 × 1.0m in 10 years. March to April. FCC 2007. (Etymology: named after the collector of the plant. Nomenclatural standard: herbarium specimen of the award plant (WSY0100898)).

(r) **'Sheila McLeod'**

cv. *Elepidote rhododendron* : 'Lem's Cameo' (s) × 'Alice Hennessy'. H: T.A. Garbutt (1998), G: S. McLeod (2000), N: T.A. Garbutt (2007), REG: T.A. Garbutt (2007). Fls 13/ball truss, campanulate, 60 × 110mm, with 7 wavy-edged lobes, buds cream flushed pink, opening creamy lemon inside and out, unmarked; stamens 14; not scented. Calyx 2mm long, green. Lvs lanceolate, 170 × 60mm, margins decurved, glossy green and distinctly bullate above, acute at apex, narrowly cordate at base, hairless. Shrub 1.5 × 0.75m in 5 years. Mid October (New Zealand). (Nomenclatural standard: colour print provided by the registrant (WSY0100131). Etymology: named for a friend of the registrant who grew the plant from a seedling given to her by the registrant).

(v) **'Sherla Bertelmann'**

cv. *Lepidote rhododendron* : (*zoelleri* × *leucogigas*) (s) × *commonae* (red-flowered form). H: R.A. Mitchell (1996), G: R.A. Mitchell, N: R.A. Mitchell (2007), REG: R.A. Mitchell (2007). Fls 7/flat truss, tubular funnel-shaped, 65 × 65mm, with 5 wavy-edged lobes, inside strong red (53B), outside strong red (53C). Lvs obovate, obtuse at apex, cuneate at base, margins decurved,

semi-glossy and moderate olive green (137A) above. Shrub 1.8 × 0.9m in 7 years. Late June to July.

(r) **'Silk Road'**

cv. Elepidote rhododendron : (*brachycarpum* (as subsp. *tigerstedtii*) × 'Phipps Yellow') (s) × 'Nelda Peach'. H: A.T. & S.A. Anderson (1997), G: A.T. & S.A. Anderson (2003), N: A.T. & S.A. Anderson (2006), REG: A.T. & S.A. Anderson (2007). Fls 16/conical truss, broadly funnel-shaped, 65 × 105mm, with 6–7 wavy-edged lobes, strong pink (48C) in bud, opening light pink (49C), with a pale yellow (11C) dorsal flare without any spotting or marking; outside deep to strong pink (48B–C). Lvs obovate, 150 × 50mm, broadly acute at apex, rounded at base, margins flat, dull and mid green above, hairless. Shrub 0.6 × 0.8m in 9 years. Late May.

(r) **'Sister Betty'**

cv. Elepidote rhododendron : Calstocker Group (s) × 'Gill's Crimson'. H: C. Weeks (1995), G: C. Weeks (2003), N: C. Weeks (2007), I: Westgate Gardens Nursery (2007), REG: C. Weeks (2007). Fls 14–15/domed truss, funnel-campanulate, 60 × 75mm, with 5 wavy-edged lobes, darker shades of red group (54) in bud, opening inside strong purplish red (63A) at margins, paling to strong purplish red (63B), with darker red deep in the centre; outside similar but tinged red (51A); some strong purplish red (63A) spotting on the dorsal lobe and edges of the adjacent lobes. Lvs elliptic to oblong, 175–200 × 60mm, broadly acute at apex, cordate at base, margins downcurved, dull and dark yellowish green (139A) above, hairless. Shrub 1.2 × 1.2m. Mid to late March.

(r) **'Speckled Honey'**

cv. Elepidote rhododendron : 'Janet Blair' (s) × ('Apritan' × 'Seattle Gold'). H: D. MacMullan (1978), G: D. MacMullan (1982), N: D. MacMullan (2006), REG: D. MacMullan (2007). Fls 7/loose ball truss, broadly funnel-shaped, 110 × 110mm, with 5 wavy-edged, reflexed lobes, strong reddish orange (31A–B) in bud, opening inside light to pale yellowish pink (27A–D), with a reddish orange basal blotch and prominent flares of vivid red (45A) to vivid reddish orange (43B) speckles on the dorsal lobe; outside light orange (28C) at the base, changing to light yellowish pink (27A) at the margins. Pedicels c.13mm long, strong reddish orange (31A). Calyx c.3mm long, vivid reddish orange (33A). Lvs elliptic, 175 × 75mm, broadly acute at apex, rounded at base, margins flat, semi-glossy above, hairless. Shrub 1.8 × 1.8m in 10 years. Midseason.

(r) **'Sugar Puff'**

cv. Lepidote rhododendron : *racemosum* (s) × *minus* Carolinianum Group. H: F. Read, G: F. Read, N: Briggs Nursery (2006), I: Briggs Nursery (2006), REG: L. Caton (2007). Fls from 7–8 inflorescence buds/terminal shoot producing 14–20 flowers in what appears to be one large domed truss, open funnel-shaped, 25 × 30mm, with 5 wavy-edged lobes, the greenish bud is light yellow-green (145B) at the base and deep purplish pink (66C) at the apex, opening white (155C) throughout, with moderate purplish pink (66D) veining outside; unmarked. Lvs lanceolate, 40 × 25mm, broadly acute at apex, rounded at base, margins flat, dull and moderate yellow-green (137C) above, scaly. Shrub 0.75 × 0.6m in 3 years. Late March to early April.

(r) **'Summer Dawn'**

cv. Elepidote rhododendron : (Ladybird Group × Inamorata Group) (s) × 'Lascaux'. H: E.G. Millais (1987), G: E.G. Millais (1995), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 9/lax truss, campanulate, 50 × 75mm, with 5 very wavy-edged lobes, buds strong yellowish pink (32C), opening inside light yellow (18B), with lime green speckling, outside brilliant to pale yellow (13C–18C), flushed pink at the lobe sinuses; stamens 11, filaments pale yellow, anthers brown; style pale green; ovary green, with white hairs; bracts reddish pink. Calyx 30mm long, irregular, pale yellow (18C). Lvs oblanceolate, 85–110 × 35–45mm, margins flat, matt green above, hairless. Shrub 1.5 × 1.5m in 10 years. Late June. (Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100241)).

(a) **'Summer Sorbet'**

cv. Deciduous azalea : parentage unknown. H: M.C. Pratt, S: E.G. Millais (c. 2000), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 13–15/domed truss, long tubular funnel-shaped, 40 × 60mm, with 5 prominent star-shaped lobes (the dorsal lobe very wavy-edged the others less so), buds deep yellowish pink (47C) to vivid red (52A), striped light greenish yellow (3D), opening inside pale purplish pink (55D), paler towards the base, dorsal lobe vivid yellow (16A) in centre, shading to light orange-yellow (16B) towards edges, outside pale purplish pink (55D) to almost white, with strong red (47B) or slightly pinker stripes; stamens prominent, 70mm long, filaments pale pink, anthers pale yellow; style flushed pink; scented. Calyx 3mm long, light green, ciliate, hairy. Lvs oblanceolate, 55 × 20mm, margins wavy and slightly decurved, glossy green above,

acute at base and apex. Shrub 1.5 × 1.0m in 10 years. Late June. (Selected by E.G. Millais as plant labelled DP121A at Dunham Massey (National Trust, UK). Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100238)).

(a) **'Summer Sunshine'**

cv. Deciduous azalea : parentage unknown. H: M.C. Pratt, S: E.G. Millais (c. 2000), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 11/very slightly domed truss, tubular funnel-shaped, 55 × 55mm, with 5 very wavy-edged lobes, buds light greenish yellow (1C), opening inside light yellow (12C), fading to pale yellow (11C), dorsal lobe entirely vivid yellow (15A) to vivid orange-yellow (21A); outside pale yellow (18C), with deep pink (more-or-less vivid red, 52A) midveins and with gland-tipped hairs; stamens exerted, filaments creamy yellow, with white hairs on the basal part; style flushed pink, stigma green; ovary green, hairy; strongly scented. Calyx 2mm long, very pale green. Lvs oblanceolate, 70 × 25mm, margins wavy and decurved, glossy above, acute at apex. Shrub 1.5 × 1.0m in 10 years. Late June. (Selected by E.G. Millais as a plant labelled DP122A at Dunham Massey (National Trust, UK). Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100237)).

(r) **'Sun Dust'**

cv. Elepidote rhododendron : 'Nancy Evans' (s) × 'Janet Blair'. H: A.T. & S.A. Anderson (1997), G: A.T. & S.A. Anderson (2002), N: A.T. & S.A. Anderson (2006), REG: A.T. & S.A. Anderson (2007). Fls 9–10/ball truss, open funnel-shaped, 50 × 80mm, with 6 wavy-edged lobes, opening inside moderate yellowish pink (38B) at margins, shading to a blend of light yellowish pink (19B) and pale yellow (20C), with vivid reddish orange (33A) or strong reddish orange (42B) speckling on the dorsal lobe; outside moderate yellowish pink (38B). Calyx c.6mm long. Lvs oblong, 90 × 45mm, broadly acute at apex, rounded at base, margins flat, dull and mid green above, hairless. Shrub 0.9 × 0.75m in 9 years. Mid May.

(a) **'Tage Buus'**

cv. Deciduous azalea : ('Brazil' × 'Ilam Yellow' seedling) (s) × *atlanticum*. H: H. Wise (1982), G: V. Søndergaard (1987), N: S.A. Askjaer (2006), I: S.A. Askjaer (2006), REG: S.A. Askjaer (2007). Fls single, 7–10/lax truss, open tubular funnel-shaped, 40–55 × 35–45mm, with 5 flat-edged lobes, buds deep purplish pink (54B), opening inside to very light rose to nearly white (pale yellowish pink, 36D), more coloured along the margins and towards the base (near strong

purplish pink, 55B), outside darker than inside (light yellowish pink, 36C), strong purplish pink (55B) along the midrib and deep purplish pink (54B) at the base; about two-thirds of the dorsal lobe covered by a vivid orange-yellow (21A) blotch; stamens 40–50mm long, filaments very light rose to white, pubescent in the basal half; style 50–60mm long, hairless, rose; ovary strongly pubescent; strongly and pleasantly scented; outside of corolla sticky, with many short bristles. Calyx 1.5–2.0mm long, mid green, bristly. Lvs obovate, 40–50 × 20–30mm, margins flat, matt green above, cuneate at base, broadly acute at apex, hairless. Shrub 1.75 × 1.75m in 15 years. Late May to mid June. (Nomenclatural standard: colour print provided by the registrant (WSY0096630). Etymology: named for the late deputy mayor of Brønderslev (in the northern part of Jutland, Denmark) who was one of the promoters in creating there a large park with c.10,000 rhododendrons).

(r) **'Tairi Sunset'**

cv. Elepidote rhododendron : 'Hotei' (s) × ('Lem's Cameo' × 'Colonel Coen'). H: L.A. Hutchison (1991), G: L.A. Hutchison (2000), N: L.A. Hutchison (2006), REG: L.A. Hutchison (2007). Fls 13–15/rather open-topped ball truss, funnel-shaped, 40 × 75–80mm, with 5–6 wavy-edged lobes, buds strong purplish red (60C) at apex, merging halfway down to moderate orange-yellow (167D), opening inside with rim varied from strong purplish red to strong purplish pink (63B–C), with paler shadings, then changing to light orange-yellow (22B) on the two dorsal lobes and to pale orange-yellow (23D) on the other lobes, outside similar but with a wide strip of varied purplish red/pink down the lobe centres; bright red eye at the base of the dorsal lobes. Calyx lobes variable in length (5–15mm), petaloid, light yellowish pink (159A), with shadings of strong yellowish pink (43D), the two larger dorsal lobes with a butterfly-shaped solid blotch of bright red. Lvs oblanceolate to elliptic, 60–75 × 24–27mm, margins slightly wavy-edged, glossy dark yellowish green (139A) above, cuneate to narrowly rounded at base, obtuse at apex, hairless. Shrub 0.85 × 1.20m in 13 years. October (New Zealand). (Nomenclatural standard: colour print provided by the registrant (WSY0096690). Etymology: named for a district to the south of the city of Dunedin, New Zealand).

(r) **'The Marquis of Lansdowne'**

cv. Elepidote rhododendron : parentage uncertain, but may have been 'Canary' × 'Meg Merrilees'. H: The Marquis of Lansdowne (1992), G: The Marquis of Lansdowne (1998), N: The Marquis of Lansdowne (2007), REG:

The Marquis of Lansdowne (2007). Fls 18–22/spherical, slightly open-topped truss, funnel-campanulate, c.65 × c.65mm, with 6–7 wavy-edged lobes, inside pale yellow (11D) at rim, deepening to light yellow (11B) in the throat; outside pale yellow (11D) at the rim, deepening in colour near the calyx; strong red (53B) blotch at base of dorsal throat; stamens 14, not exerted, filaments pale yellowish green, anthers reddish brown; style pale yellowish green. Pedicels reddish. Lvs elliptic to lanceolate or oblanceolate, 120 × 60mm, margins upcurved, cuneate at base, acute at apex, slightly glossy and with yellowish main veins above, hairless. Compact, upright shrub, 2.0 × 1.8m in 15 years. May. (Nomenclatural standard: colour print provided by the registrant (WSY0100734)).

(f) '**Tholasma**'

cv. *Elepidote rhododendron* : *yakushmanum* (Exbury form) (s) × *lanatum* (RSF 333/1976). H: H. Kuhn, G: S.A. Askjaer (1992), N: S.A. Askjaer (2003), I: S.A. Askjaer (2007), REG: S.A. Askjaer (2007). Fls 7–10/shallowly domed truss, funnel-campanulate, 45–60 × 50–65mm, with 5(–6) slightly wavy edged lobes, deep pink (52B) in bud, opening inside yellowish white (158D), with numerous, small, strong red (53D) spots on the basal two-thirds of the three dorsal lobes (the spots changing to moderate red (179A) with age), outside yellowish white (158D) with the spots on the inside showing through the dorsal lobes; stamens 10, filaments 15–25mm long, white, basal half pubescent, anthers dark brown to black; style 35–45mm long, white, hairless, stigma light greenish yellow (3C). Calyx up to 1mm long, very light green. Lvs oblong, 60–80 × 30–40mm, margins decurved, matt green above, rounded at base, broadly acute at apex, with an indumentum below which is light orange-yellow (25D) at first, aging to moderate orange-yellow (168D). Shrub 2.0 × 1.75m in 20 years. Mid to late May. (Nomenclatural standard: colour print provided by the registrant (WSY0096632). Etymology: epithet created from the letters of the names of the registrant's grandchildren: Thomas, Lasse and Marie.)

(r) '**Tia**'

cv. *Elepidote rhododendron* : 'Tweedy Bird' (s) × ('Bambi' × (*yakushmanum* × *proteoides*)). H: W.W. Thompson (1985), G: Thompson's Nursery (1990), N: J. Barlup (2006), I: Thompson's Nursery (2000), REG: J. Barlup (2007). Fls 14/ball truss, funnel-campanulate, 45 × 60mm, with 6 wavy-edged lobes, deep red (53A) in bud, opening inside and out light greenish yellow (8C), with strong red (53B) tips and

on the 4 dorsal lobes softening to strong red (53D) along the margin for a distance of c.15mm; two, strong red (53B), basal, dorsal flares, c.20mm long, surmounted by spotting of the same colour extending for a total of c.25mm; anthers very dark red. Calyx c.15mm long, light greenish yellow (8C), with touches of strong red (53B). Lvs elliptic, 110 × 50mm, broadly acute at apex, rounded at base, margins upcurved, glossy and moderate olive green (147A) above, hairless. Shrub 1.0 × 1.2m in 9 years. Mid April.

(v) '**Tom Grieve**'

cv. *Lepidote rhododendron* : selection from *macgregoriae*, collected by Paddy Woods in Papua New Guinea (RBGE accession no.19875220). S: Royal Botanic Garden, Edinburgh, N: G. Argent (2007), REG: Royal Botanic Garden, Edinburgh (2007). Fls 10–15/umbellate truss, shortly tubular campanulate, 10–25 × 20–30mm, with 5 flat-edged lobes, opening vivid yellow (16A) throughout, unmarked; anthers dark brown. Calyx 1mm long, green. Lvs ovate-elliptic, 50–140 × 25–50mm, margins decurved, broadly tapering at base, shortly acuminate at apex, with a scaly indumentum (scales silver at first, maturing to brown). Shrub 0.8 × 0.5m in 5 years. Flowering irregularly, mostly in spring. AM 2007. (Etymology: named for a member of the staff at the Royal Botanic Garden, Edinburgh. Nomenclatural standard: herbarium specimen of the award plant (WSY0100902)).

(a) '**Tropical Salad**'

cv. *Deciduous azalea* : parentage unknown. H: M.C. Pratt, S: E.G. Millais (c. 2000), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 9–10/domed truss, tubular funnel-shaped, 30 × 70mm, 5-lobed (the dorsal lobe very wavy-edged, the others less so), buds deep yellowish pink (39B), opening inside to strong pink (48C), fading to strong purplish pink (55B), outside deep purplish pink (55A), fading to strong purplish pink (55B); dorsal lobe prominent, brilliant orange-yellow to strong orange (23B–25A), fading to strong orange-yellow (24B); stamens prominently exerted, 80mm long, filaments pale pink, anthers pale yellow; style pink, stigma bright green; slightly scented. Calyx 4mm long, light green. Lvs elliptic, 60 × 20mm, margins wavy, glossy green above, acute at base and apex. Shrub 1.5 × 1.0m in 10 years. Late June. (Selected by E.G. Millais as an un-numbered plant from a planting of Denny Pratt azaleas at Dunham Massey (National Trust, UK). Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100239)).

(a) **'Water Baby'**

cv. Deciduous azalea : *viscosum* hybrid × 'Water Girl'. H: E.G. Millais (1997), G: E.G. Millais (2000), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 5–6/small upright truss, long tubular funnel-shaped, up to 46 × 30mm, with 5–6 flat-edged lobes which become slightly reflexed, buds strong red (53B) on tube, deep purplish pink (54B) on the lobes, opening inside moderate purplish pink (68C); outside strong red (53C), with this colour extending up the midribs, edges of lobes paler (moderate purplish pink (68C)); unmarked; stamens 5, exerted, prominent, filaments red; style deep pink (52B), stigma dark. Corolla with moderately dense, short, white, eglandular hairs and numerous longer, pale, gland-tipped hairs (the glands pale). Calyx up to 5mm long, unequally lobed, pale yellow-green, glandular ciliate and shortly hairy. Lvs oblanceolate, 32 × 12mm, margins decurved, not wavy, attenuate at base, broadly acute to obtuse at apex, apiculate, ciliate and with pale, eglandular hairs on the midveins above and below. Shrub 1.0 × 0.75m in 10 years. June to July. (Etymology: named for the "baby pink" flowers on a swamp honeysuckle azalea. Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100736)).

(a) **'Water Girl'**

cv. Deciduous azalea : *viscosum* hybrid, grown from seed supplied by the American Rhododendron Society 1991. G: E.G. Millais (1995), N: D.G. Millais (2007), I: Millais Nurseries, REG: D.G. Millais (2007). Fls 10/loose, small, domed truss, long tubular funnel-shaped, 48 × 35mm, with 5 flat-edged lobes, strong red (53C) in bud, inside strong purplish pink (55B) to light purplish pink (55C), outside strong red (53D) extending up tube and on lobe midribs, with light purplish pink (55C) margins to the lobes; inconspicuous light yellow (20B) markings on the dorsal lobe; stamens 5, exerted, filaments white at base, deep purplish pink (55A) at apex; style white at base, deep pink (52B) at apex, stigma dark; scented. Corolla and pedicels with some short, pale, eglandular hairs and some glandular hairs, these often with red glands and pale shafts. Calyx up to 2.0mm long, lobes irregular, pale yellow-green, glandular ciliate. Lvs oblanceolate, up to 50 × 20mm, margins decurved, not wavy, matt blue-grey, tinged purplish, attenuate at base, acute (to shortly acuminate) at apex, apiculate, ciliate and with very sparse, pale, eglandular hairs on

the midvein below. Shrub 1.2 × 0.75m in 10 years. July. (Etymology: regarded as a later, larger version of 'Water Baby'. Nomenclatural standard: herbarium specimen provided by the registrant (WSY0100738)).

(r) **'Winter Intruder'**

cv. *Elepidoterhododendron*: chance seedling found underneath a plant of *arboreum* subsp. *delavayi*. S: F.J. Williams (1960s), G: F.J. Williams & P. Tregunna (1960s), N: F.J. Williams, REG: C.H. Williams & J. Parsons (2007). Fls 9–11/flattened hemispherical truss, broadly tubular campanulate, 36–42 × c.40mm, with 5 wavy-edged lobes, vivid red (57A) in bud and on opening, paling slightly on the tube both inside and out, the basal part of the dorsal lobe and most of the dorsal section of the throat with numerous dark red (59A) spots, sometimes extending on to parts of the adjacent lobes and accompanied by a few paler spots, base of the corolla with small areas of dark staining, best developed at the base of the dorsal side; style and filaments white; stigma deep red-purple; ovary pale green, with numerous, short, gland-tipped hairs and sparse, long, eglandular hairs. Calyx c.2mm long, very pale whitish green, with numerous, very short-stalked, purplish glands which extend down the pale greenish white pedicels. Lvs mostly oblanceolate, up to 140 × 47mm, margins flat, dark matt green above, truncate at base, obtuse and mucronate at apex, lower surface on mature leaves with sparse, very tiny, pale or slightly brownish hairs. Shrub to at least 3.8–4.5 × 3.8–4.5m. January. (Nomenclatural standard: herbarium specimen provided by the registrants (WSY0100137)).

(r) **'Winter's Tale'**

cv. *Elepidote rhododendron* : parentage unknown. H: W.G.T. Hyde (1960s), G: W.G.T. Hyde (1970s), N: Mrs R.G. Legrand (2007), I: Mrs R.G. Legrand (2007), REG: Mrs R.G. Legrand (2007). Fls 9/ truss, open trumpet-shaped, 44 × 30mm, with 7 wavy-edged lobes, buds deep purplish pink (58D) at base, pale yellow (18C) at apex, opening white (155D) throughout; delicately scented. Calyx 4mm long, brilliant yellow-green (150C). Lvs broadly oblanceolate, 90 × 37mm, margins decurved, narrowly obtuse at base, acute at apex, hairy (the hairs moderate yellow-green (146D)), with a slight hint of moderate yellow (163C)). Shrub 1.2 × 2.3m in 40 years. May. (Etymology: named as part of a collection with Shakespearian names).

# NAMES AND ADDRESSES

This list comprises the names and abbreviated addresses of all individuals, nurseries and estates mentioned in this Supplement as playing a role in the development and registration of the cultivars listed.

\* = deceased

- Anderson, A.T. & S.A.  
Franklin Lakes, New Jersey, USA
- Argent, G.  
Royal Botanic Garden, Edinburgh,  
Scotland
- Askjaer, S.A.  
Spøttrup, Denmark
- Baricilli, S.  
Sea Cliff, New York, USA
- Barlup, J.  
Bellevue, Washington, USA
- Bertelmann, S.  
Kea'au, Hawaii, USA
- Brack, W.  
St James, New York, USA
- Briggs Nursery  
Olympia, Washington, USA
- Caton, L.  
Briggs Nursery, Olympia, Washington, USA
- Cook, A.R.  
Vancouver, British Columbia, Canada
- Corley, T.E.  
Auburn, Alabama, USA
- Cox, P.A. & K.N.E.  
Glendoick Gardens, Glencarse,  
Perthshire, Scotland
- Craig-Mooney, A.  
Buckland Monachorum, Devon, England
- Dancer, C.  
Victoria, British Columbia, Canada
- Dosser, D.J.  
Lower Wilmot, Tasmania, Australia
- Garbutt, T.A.  
Oamaru, North Otago, New Zealand
- Glendoick Gardens  
Glencarse, Perthshire, Scotland
- Glynneden Gardens  
Greenbank, Washington, USA
- Greig, E.J.  
Vancouver Island, British Columbia,  
Canada
- Gustafson, R.H.  
West Windsor, New Jersey, USA
- Hagan, K. & G.  
Kingston Beach, Tasmania, Australia
- Hibben, G.  
Lincoln Massachusetts, USA
- Hutchison, L.A.  
Mosgiel, Otago, New Zealand
- \*Hyde, W.G.T.  
Ferndown, Dorset, England
- Jefferis, R.P.  
Media, Pennsylvania, USA
- Jefferis, S.  
Wayne, Pennsylvania, USA
- Johnson, D.B.  
Mountain Rest, South Carolina, USA
- King, D.E.  
Issaquah, Washington, USA
- Kruppa, R.  
Volcano, Hawaii, USA
- Kuhn, H.  
Muttontown, New York, USA
- Lansdowne, The Marquis of  
Bowood House, Wiltshire, England
- Legrand, Mrs R.  
West Moors, Dorset, England
- Livingston, P.  
Narberth, Pennsylvania, USA
- MacMullan, D.  
Hamburg, Pennsylvania, USA
- Maeda, H.  
Kirishima-city, Kagoshima-prefecture,  
Japan
- McLeod, S.  
Oamaru, North Otago, New Zealand

- Millais, \*E.G. & D.G.  
Millais Nurseries, Farnham, Surrey, England
- Millais Nurseries  
Farnham, Surrey, England
- Mitchell, R.A.  
Volcano, Hawaii, USA
- Murcott, R.  
East Norwich, New York, USA
- Norman, F.S.  
Bexleyheath, Kent, England
- Pacific Island Nursery  
Kea'au, Hawaii, USA
- Parsons, J.  
Head gardener, Caerhays Castle, Gorran,  
Cornwall, England
- Patterson, C.  
Norwell, Massachusetts, USA
- \*Pratt, M.C.  
Pulborough, West Sussex, England
- Raper, A.B.  
The Patch, Victoria, Australia
- RareFind Nursery  
Jackson, New Jersey, USA
- Read, F.  
Read Nursery, Ramapo, New York, USA
- Roadside Rhododendrons,  
Mapleton, Oregon, USA
- Round, J.  
Vancouver, British Columbia, Canada
- Royal Botanic Garden, Edinburgh  
Edinburgh, Scotland
- Sanders, M.  
Roseburg, Oregon, USA
- Sanders Nursery  
Roseburg, Oregon, USA
- Sim, Mrs A.  
Oamaru, North Otago, New Zealand
- Simeone, V.  
Oyster Bay, New York, USA
- Sisson, K.  
Brookville, New York, USA
- \*Smith, C.C.  
St Paul, Oregon, USA
- Smith Garden, Cecil & Molly  
St Paul, Oregon, USA
- Smith, J.S.  
Roadside Rhododendrons, Mapleton,  
Oregon, USA
- Smitherman, R.O.  
Auburn, Alabama, USA
- Snell, G.L.S.  
Highfields, Queensland, Australia
- Søndergaard, V.  
Herning, Denmark
- Spady, H.A.  
Silverton, Oregon, USA
- Stewart, Robert  
Springfield, Virginia, USA
- Stipe, W.A.  
Formerly of Meerkerk Rhododendron  
Gardens, Greenbank, Washington, USA
- Thompson, W.W.  
Waldport, Oregon, USA
- Thompson's Nursery  
Waldport, Oregon, USA
- Todd, N.  
Victoria, British Columbia, Canada
- Tregunna, P.  
Formerly head gardener, Caerhays Castle,  
Gorran, Cornwall, England
- Van Veen Nursery  
Portland, Oregon, USA
- Wedemire, G.  
Maple Ridge, British Columbia, Canada
- Weeks, C.  
Westgate Gardens Nursery, Eureka,  
California, USA
- Weingarten, B.  
Washougal, Washington, USA
- Westgate Gardens Nursery  
Eureka, California, USA
- Williams, C.H.  
Caerhays Castle, Gorran, Cornwall, England
- Williams, F.J.  
Caerhays Castle, Gorran, Cornwall, England
- Wise, H.  
Charleston, West Virginia, USA
- Woodard, G.  
Westbury, New York, USA



# CORRECTIONS TO THE INTERNATIONAL RHODODENDRON REGISTER AND CHECKLIST

The following list indicates further significant additions and corrections that need to be made to the printed text of the *Register and Checklist* published in 2004 or to the three subsequent Supplements. The Registrar is grateful to all those who have brought such errors to his notice and welcomes knowledge of further amendments that may need to be made. Minor changes are not noted here but such changes have been made to the database and will be available in any future electronic version. The database will also be taking into account additional information that comes to light relating to any individual entry, as well as including further entries to the Checklist.

In the following list the page number refers to the page in the *Register and Checklist* (or one of its Supplements) on which the entry concerned is to be found.

Akbar Group/'Akbar' (p.28). Change parent 'Loderi King George' to 'King George'.

'Capistrano' (p.203). Insert two extra brackets in parentage before *catawbiense* and one after Fabia Group.

Esmeralda Group (p.418). Change parent 'Loderi King George' to 'King George'.

'Frederik d'Honneur' (p.473). This should be 'Frederic d'Honneur'.

'Glendoick Crimson' (p.514). This should be treated as a trade designation (GLENDICK CRIMSON) for 'Gle004'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Dream' (p.514). This should be treated as a trade designation (GLENDICK DREAM) for 'Gle005'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Ermine' (p.514). This should be treated as a trade designation (GLENDICK ERMINE) for 'Gle006'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Frolic' (First Supplement, p.15). This should be treated as a trade designation (GLENDICK FROLIC) for 'Gle007'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Garnet' (p.514). This should be treated as a trade designation (GLENDICK GARNET) for 'Gle008'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Gem' (p.515). This should be treated as a trade designation (GLENDICK GEM) for 'Gle002'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Goblin' (p.515). This should be treated as a trade designation (GLENDICK GOBLIN) for 'Gle010'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Honeydew' (First Supplement, p.15). This should be treated as a trade designation (GLENDICK HONEYDEW) for 'Gle012'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Ice Cream' (First Supplement, p.15). This should be treated as a trade designation (GLENDICK ICECREAM) for 'Gle013'. Glendoick is a registered trademark belonging to Glendoick Gardens. The use of ICECREAM as one word in the trade designation is deliberate.

'Glendoick Jewel' (p.515). This should be treated as a trade designation (GLENDICK JEWEL) for 'Gle020'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Mystique' (First Supplement, p.15). This should be treated as a trade designation (GLENDICK MYSTIQUE) for 'Gle014'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Petticoats' (First Supplement, p.15). This should be treated as a trade designation (GLENDICK PETTICOATS) for 'Gle015'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Ruby' (p.515). This should be treated as a trade designation (GLENDICK RUBY) for 'Gle016'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'Glendoick Velvet' (p.515). This should be treated as a trade designation (GLENDICK VELVET) for 'Gle018'. Glendoick is a registered trademark belonging to Glendoick Gardens.

'J. Hutton Edgar' (p.675). The reference to Cullen (1980) should read Philipson and Philipson (1975).

'Lemon Dream' (p.773). This was supplied with an incorrect parentage at the time of registration. The hybridiser now confirms that this should read: 'Creamy Chiffon' × *yakushmanum*. Exbury form.

'Red Arrow' (p.1139). The parentage should be corrected to read: 'Dopey' × 'Royal Blood'.


'Rubinfeuer' (p.1198). This was raised by A.W. Backhus c.1947–1948 and named by W. Schmalscheidt in 1988.

'Sanko-no-tsuki' (p.1219). Add: Satsuki.

'Silver Jubilee' (p.1262). The parentage should be corrected to read *yakushmanum* × 'Coronation Day'.

General correction: With the agreement of Mr & Mrs George of Hydon Nurseries all the cultivars of rhododendrons and azaleas that have been bred, named and/or introduced at the nursery have had the roles in their entries in the Register and Checklist standardised to read: H: A.F. George, G: Hydon Nurseries, N: Mrs A.M. George, I: Hydon Nurseries and REG: Hydon Nurseries. The only exception to this is 'Anne George' which was named by Mr George for his wife.

ISBN 978-1-902896-91-5



9 781902 896915 >